

PROCEEDINGS Pub ID
31756
for the Colorado Meeting on
Fundamental Optical Processes in Semiconductors

FOPS 2004
August 8-13, 2004
Stanley Hotel, Estes Park CO

Carrier transfer between InGaAs/GaAs quantum dots observed by differential transmission spectroscopy

K. L. Silverman and R. P. Mirin


*National Institute of Standards and Technology,
Boulder, Colorado 80305*

silverma@boulder.nist.gov

S. T. Cundiff

*JILA, National Institute of Standards and Technology
and University of Colorado, Boulder, Colorado 80309*

We observe carrier transfer between self-assembled InGaAs/GaAs quantum dots (QDs) at temperatures above 230 K using differential transmission spectroscopy (DTS). The sample is a single layer of QDs embedded in a semiconductor waveguide to increase the interaction length of the weakly absorbing QDs. Pump and probe light is generated by an optical parametric oscillator synchronously pumped by a mode-locked Ti:sapphire laser. The probe signal is isolated from the pump by high frequency modulation and detection with an RF lockin amplifier.



The figure above contains spectrally resolved DT on the ground state of the QDs at various delays between pump and probe. For this measurement the pump beam is spectrally filtered to a bandwidth of ~ 2 nm with a zero-dispersion pulse shaper and the probe beam is spectrally resolved with a monochromator after interaction with the QDs. This allows us to observe the evolution of the DT signal for QDs resonant and non-resonant with the pump excitation. At small pump-probe time delays, the DT signal indicates a spectral hole much larger than the pump bandwidth. We believe that the width of this initial signal (12 nm FWHM) is representative of the homogeneous linewidth of the QDs. As the delay is increased, a signal is observed at wavelengths not covered by the initial spectral hole. After analyzing the time dependence of these two contributions to the differential transmission signal, it was determined that the decay time of the spectral hole and the rise time of the non-resonant signal were almost identical (~ 32 ps). This is clear indication that carriers initially generated in resonant QDs escape, and are recaptured by non-resonant QDs. Previous DT measurements have shown this initial recovery of the absorption bleaching, but were not able to resolve the transfer of carriers to non-resonant QDs.