



ELSEVIER

Physica A 306 (2002) 230–242

PHYSICA A

www.elsevier.com/locate/physa

Statistical physics and liquid water: “What matters”

H.E. Stanley^{a,*}, S.V. Buldyrev^a, N. Giovambattista^a,
E. La Nave^{a,b}, A. Scala^{a,b}, F. Sciortino^b, F.W. Starr^c

^aCenter for Polymer Studies and Department of Physics, Boston University, Boston, MA 02215, USA

^bDipartimento di Fisica Università di Roma La Sapienza, Istituto Nazionale di Fisica della Materia and INFN Center for Statistical Mechanics and Complexity, Piazzale Aldo Moro 2, 00185 Roma, Italy

^cPolymers Division and Center for Theoretical and Computational Materials Science, National Institute of Standards and Technology, Gaithersburg, MD 20899, USA

Abstract

We present an overview of recent research applying ideas of statistical mechanics to try to better understand the statics and especially the dynamic puzzles regarding liquid water. The take-home message for the static aspects is that what seems to “matter” more than previously appreciated is local tetrahedral order, so that liquid water has features in common with SiO₂ and P, as well as perhaps Si and C. For the dynamic aspects, what may “matter” is the number of *diffusive* directions in the potential energy landscape. © 2002 Published by Elsevier Science B.V.

PACS: 65.20.+w; 61.20.-p; 61.50.-f; 64.10.+h

Keywords: Water; Local tetrahedral order; Potential energy landscape; Instantaneous normal modes; Dynamic heterogeneities

1. The goal: understanding “what matters”

Many physicists are attracted to physics because of the focus on understanding just enough of a subject to comprehend the key features that really matter. As soon as some physicists feel they understand “what matters”, insatiable appetites for novelty force attention to new puzzles. At the 1968 Kyoto *STATPHYS-8* conference, among the most exciting new developments was the degree to which the principle of scale

* Corresponding author.

E-mail address: hes@bu.edu (H.E. Stanley).

invariance provided increased understanding of “what matters” near a critical point. The key point was that what matters near a critical point is the correlation length for statics, and the correlation time for dynamics. An exciting question these days is “what matters” in understanding the statics and dynamics of liquid water, and important clues are emerging when one focuses on behavior in the deeply supercooled region, especially just above—and just below—the “critical” temperature T_{MCT} predicted by mode coupling theory (MCT).

2. Statics: “what matters” is local tetrahedral geometry

2.1. Introduction

Liquid water is not a typical liquid. However, some progress has occurred in understanding its highly anomalous equilibrium and dynamical properties [1–4]. Water is a space-filling hydrogen bond network, as expected from continuum models of water. However when we focus on the well-bonded molecules, we find that water can be regarded as having certain clustering features—the clusters being not isolated “icebergs” in a sea of dissociated liquid (as postulated in mixture models dating back to Röntgen) but rather patches of well-bonded molecules embedded in a highly connected network or “transient gel” [5–11]. Similar physical reasoning applies if we generalize the concept of well-bonded molecules to molecules with a smaller than average energy [12] or to molecules with a more ordered than average “local structure” [13].

2.2. Liquid–liquid phase transition hypothesis

A few years ago, Poole, Sciortino, and Essmann made computer simulations of the ST2 model of water, with the goal of exploring in detail what might happen in the low-temperature region [14]. What they discovered in computer water was the apparent existence of a second critical point C' , below which the liquid phase separates into two distinct phases—a low-density liquid (LDL) and a high-density liquid (HDL).

Although in experiments we cannot get closer than 5–15°, C' nonetheless exerts a strong effect in the experimentally accessible region of the phase diagram. If we have a singularity in our phase diagram at a well-defined critical point, it is going to have an effect on an entire region around it—a “critical region”. The size of this critical region depends on the material, but it is usually somewhere between 10% and 100% above the critical temperature and pressure that response functions are significantly affected (e.g., an ideal gas has an inverse compressibility given by T , while an interacting gas has an inverse compressibility that deviates significantly from T by a factor of two above the critical point). It is not required that the system is exactly *at* its critical point for the system to exhibit remarkable behavior, such as the phenomenon of critical opalescence discovered and correctly explained in 1869 by Andrews [15] in terms of increased fluctuations away from (but close to) the critical point. It is for this reason that critical phenomena are particularly interesting.

2.3. Experimental work

When liquid water is supercooled below the homogeneous nucleation temperature T_H (-38°C at $P=1$ atm), crystal phases nucleate homogeneously, and the liquid freezes spontaneously to the crystalline phase. Mishima creates 1 cm^3 high-pressure ices in a piston–cylinder apparatus, decompresses the sample at a constant rate of 0.2 GPa/min , and—because melting is endothermic—observes melting transitions of the ice polymorphs using a thermocouple to detect a change in the sample temperature during the decompression [16,17]. He then determines melting pressures at different temperatures. The melting curves he obtains agree with previously reported data for stable melting lines [18,19], and extend our knowledge of the location of metastable melting lines to much lower temperatures.

The Gibbs potential G of the ice polymorphs is known. Since G is identical in coexisting phases, locating the melting lines of the ice polymorphs is sufficient to learn G for water along these lines. By interpolating data for G obtained along these melting lines, one can find the approximate experimental G for a wide range of temperatures and pressures in the no-man’s land below T_H [16]). After finding G as a function of pressure P and temperature T , one can find by differentiation the volume as a function of P and T . Volume as a function of T is just what we want—this is the equation of state of liquid water. The P – V – T relation found is consistent with the existence of a line of first-order liquid–liquid transitions which continues from the line of low-density amorphous transition to high-density amorphous transitions and terminates at an apparent critical point C' . The P – V – T relation is also consistent with other known experimental data [20–25] and also with simulation results [14,26–37].

2.4. Theoretical work

The most natural response to the concept of a second critical point in a liquid is bafflement—such a thing just does not make sense. To make the concept more plausible, we offer the following remarks. Consider a typical member of the class of intermolecular potentials that go by the name of core-softened potentials [38–40]. Recently such potentials have been re-visited [27,41–54]; they are attractive to study because they can be solved analytically in one dimension and are tractable to study using approximation procedures (and simulations) in higher dimensions. They are also more realistic than one might imagine at first sight, and indeed may reflect “what matters” in water–water interactions, since the repulsive soft core mimics the effect of the small number (4) of nearest neighbors in liquids with a local tetrahedral structure. Although such a picture may seem to be oversimplified, it is consistent with neutron data [22–25]. Also, simulation results are in good accord with neutron results (see, e.g., Ref. [55]), and Sasai relates these two distinct local structures to dynamic properties [56].

One can characterize (or at least “caricature”) the local structural heterogeneities that appear in liquid water by simulations. Specifically, Canpolat and collaborators [57] considered state points of liquid water at different pressures—especially near its phase boundaries with ice I_h and with ice VI. In the spirit of the “Walrafen

pentamer”, they develop a model of interacting water pentamers, and find two distinct local energy minima which they identify with two well-defined configurations of neighboring pentamers. The “Walrafen pentamer” is defined by four water molecules located at the corners of a tetrahedron that are hydrogen-bonded to a central molecule—see, e.g., Refs. [58,59]; the corner molecules are separated from the central molecule by 2.8 Å, corresponding to the first peak in the oxygen–oxygen radial distribution function. Canpolat and collaborators advance the hypothesis that these configurations may be related to the local “high-density” and “low-density” structural heterogeneities occurring in liquid water.

2.5. Outlook

Before concluding this brief discussion of statics, we ask “What is the requirement for a liquid to have such a second critical point?” By the arguments presented above, some other liquids should display second critical points, namely systems that at low temperature and low pressure have anticorrelated entropy and specific volume fluctuations. Thus a natural extension to our work is to consider other tetrahedrally coordinated liquids. Since other tetrahedral liquids have that similar features, we might anticipate that similar critical points occur on the liquid free energy surface of these liquids. Evidence in favor of this possibility has been reported for SiO₂ [60,61], amorphous GaSb [62,63], C [64,65], and Si [66]. Recently, clear experimental evidence for a liquid–liquid phase transition has been reported in phosphorus, where the low-density liquid phase is a molecular liquid of tetrahedral P₄ “molecules” [67,68]. With a change in pressure, the low-pressure, low-density molecular liquid transforms to a high-pressure, high-density polymeric liquid. During the transformation, two forms of liquid coexist, showing that phosphorus has a first-order liquid–liquid phase transition.

3. Dynamics on the potential energy landscape: “what matters” is the number of diffusive directions

3.1. Introduction

The study of the dynamics in supercooled liquids is receiving great interest [69] due to novel experimental techniques [70,71], detailed theoretical predictions [72], and by the opportunity to follow the microscopic dynamics via computer simulation [73,74]. MCT [72] quantitatively predicts the time evolution of correlation functions and the dependence on temperature T of characteristic correlation times. Unfortunately, the temperature region in which MCT is able to make such predictions for the long time dynamics is limited to weakly supercooled states. Parallel with the development of MCT, theoretical work [75–79] has called attention to thermodynamic approaches to the glass transition, and to the role of configurational entropy in the slowing down of dynamics [80–82]. These theories, which build on ideas put forward some time ago [83–85], stress the relevance of the topology of the potential energy landscape (PEL) explored in supercooled states. Detailed studies of the PEL may provide insights into

the slow dynamics of liquids, and new ideas for extending the present theories to the deep supercooling regime.

3.2. *Instantaneous normal modes and the topology of the PEL*

One approach to understanding the role of the PEL is to study the connectivity between different local configurations using the instantaneous normal mode (INM) formalism [86]. Analogous to the standard normal mode theory for solids, an INM is the eigenfunction of the Hessian, which is the matrix of the second derivatives of the potential energy with respect to all $6N$ atomic coordinates. In a liquid state, the eigenvalues of the Hessian matrix are not all generally positive; the negative eigenvalues indicate a downward curvature of the PEL, i.e., indicate unstable directions for the system. Previous studies using the INM formalism indicate that the number of directions with negative curvature is reduced on cooling, motivating theories relating diffusion in liquids to the INM density of states [87,88]. Low-temperature liquid dynamics involve the superposition of fast oscillations around quasi-equilibrium positions (intra-basin motion) and the rearrangement of the system between these positions (inter-basin motion). The typical oscillation period is much shorter than the typical time needed by the system to rearrange itself, i.e., the structural relaxation time. INM theories for diffusion relate the diffusion of the system in configuration space to activated processes of inter-basin motion. In this respect, the unstable modes are considered representative of the barriers crossed when the system changes basins.

One approach [89,90] among many [91,92] for separating the diffusive modes (basin changes in configuration space) from the non-diffusive modes (no basin changes) is classifying the modes according to their potential energy profile (Fig. 1), and partition those unstable modes into two groups: (i) unstable normal modes due to the anharmonicities (shoulder modes) and (ii) modes along which the system is crossing a saddle (double-well modes) [91]. In order to distinguish between shoulder and double-well modes, the potential energy profile is calculated along straight paths that follow the direction of the eigenvector. Furthermore, to distinguish the false and true double wells, we calculate the steepest descent trajectories starting from the opposite sides of the saddle. A mode represents true double well, and this is called a diffusive mode if these trajectories end up in two distinct local minima.

3.3. *Results*

Next we discuss the numerical relationship between D and the number of diffusive modes f_{diff} in the vicinity of the fragile-to-strong crossover temperature T_{\times} . We review recent work on two different models of tetrahedral liquids, the SPC/E extended simple point charge model for water [90,93] and the BKS model of silica [94]. For silica (Fig. 2), the fragile-to-strong transition temperature T_{\times} coincides numerically [94–96] with the critical temperature T_{MCT} identified by MCT. For both models, it appears that D depends on T and P only through f_{diff} —the analog of the magnetization $M(H, T)$ of a ferromagnet depending on magnetic field H and temperature T only through the




Fig. 1. Schematic sketch of the possible shapes of the PEL associated with imaginary eigenvalues. Unstable modes are first separated into shoulder and double-well modes. Furthermore, double-well modes are split into diffusive and non-diffusive ones. Adapted from Ref. [90].

correlation length ξ . Specifically, for both models it appears that D follows a general power-law relation of the form

$$D/T \sim (f_{\text{diff}})^{\alpha} \quad (1)$$

for roughly two decades in f_{diff} and three decades in D/T . For the water model, $\alpha \approx 2$ while for the silica model it appears that $\alpha \approx 1.3$. In the case of silica, the identical functional form describes the relationship between D and f_{diff} both above and below T_{\times} , showing that while the T dependence of both D and f_{diff} is sensitive to the microscopic mechanisms controlling the dynamics, the fragile-to-strong transition does not affect the relation between D and f_{diff} . The exponent value $\alpha = 2$ found for water has recently been theoretically interpreted [97].

In summary, then, two different dynamical mechanisms affect the slowing down of the dynamics in supercooled states [90]:

- (i) In the weakly supercooled region, the slowing down of the dynamics arises from the progressive reduction in the number of directions where free exploration of configuration space is possible. The system is always located close to a multi-dimensional ridge between different basins, and the time scale of




Fig. 2. Arrhenius plot of (a) the diffusion constant D for Si atoms in SiO₂ and (b) f_{diff} . The crossover to the straight line Arrhenius behavior below T_x represents the fragile-to-strong crossover for silica. (c) shows The parametric relation D/T vs f_{diff} in a log–log scale. The data are smooth through the “mode-coupling” crossover temperature T_x . Adapted from Ref. [94].

the long-time dynamics is set by the time required to probe one of the free directions. In this range of T , the diffusion is not limited by the presence of energy barriers that must be overcome by thermally activated processes, but is controlled by the limited number of directions leading to different basins along almost constant potential energy paths. Furthermore, the number of free directions completely determines the value of D , independent of the thermodynamic parameters T and ρ .

- (ii) Close to T_{MCT} , the system starts to sample regions of configuration space that have no free directions. The change in the dynamics above and below T_{MCT} can be viewed as a change in the properties of the PEL sampled in equilibrium, from configurations always close to a ridge of progressively lower and lower dimension to configurations far from any ridge [95,96]. Below T_{MCT} , the system must go close to the ridge and then select the right direction. The search for the ridge below T_{MCT} , i.e., the search for a rare event, can be probably described as an activated process, which corresponds to Arrhenius behavior of the diffusion constant.
- (iii) The relation between connectivity and number of local minima in the PEL—which can be calculated in theoretical models as recently done for the random energy model [98]—may help build on the existing ideas bridging thermodynamics and dynamics [99].

4. Dynamics below the MCT: “what matters” is cooperative motion

4.1. Introduction

As a supercooled liquid is cooled toward the glassy state, the system is increasingly found near local potential energy minima, called inherent structure configurations [83]. In this description, in the glassy state, the system is localized in one of the potential energy basins [95,100–102]. While such a picture of liquid dynamics is difficult to verify experimentally, computer simulation offers an excellent opportunity to explore these ideas. For a pre-defined liquid potential, a liquid trajectory can be generated via molecular dynamics simulation and the local potential energy minima can be evaluated by an energy minimization method [83]. With this procedure, the motion in phase space is converted into a minimum-to-minimum trajectory, or *inherent structure trajectory*. A general picture of the system moving among a set of basins surrounding the multitude of local minima has evolved. More specifically, simulations have shown that both the depth of the minima sampled by the system, as well as the number of these minima, decrease on cooling [102,103].

The description of the real motion of the system as an inherent structure trajectory becomes a powerful way of separating the vibrational contribution, responsible for the thermal broadening of instantaneous measurements from the slow structural component [104]. Such an approach becomes even more powerful below T_{MCT} , since most of the instantaneous configurations are far from saddles, making correlation functions calculated from the inherent structure trajectory fully account for the α -relaxation dynamics [100].

4.2. Results

Recent results [105] are based on molecular dynamics simulations of the SPC/E model [106] of water for 216 molecules, at fixed density $\rho = 1 \text{ g/cm}^3$. The numerical procedure is described in Ref. [107]. The trajectories are analyzed at $T = 180 \text{ K}$, and the mode coupling temperature for this density is $T_{\text{MCT}} = 193.6 \text{ K}$ [107], so the system




Fig. 3. (a) Displacement of each of the 216 molecules during the course of a transition from one inherent structure to another. (b) Distribution of displacements u of the oxygen atoms between inherent structure changes, $P(u)$, sampled along a 30 ns trajectory in 20,000 inherent structure changes. The exponential tail of $P(u)$, with a characteristic length of about 0.02 nm, is mostly due to the highly mobile molecules, while the power law with exponent 2.5 would correspond to an “elastic” response of the system to these highly mobile molecules [115]. Adapted from Ref. [105].

is in the deep supercooled liquid state. At this temperature, the diffusion coefficient is four orders of magnitude smaller than its value at $T=300$ K and only a few molecules move significantly (with displacements larger than 0.025 nm) at each simulation time step.

To aid in understanding the distribution of the displacements during the IS changes, Fig. 3(a) shows the displacements u of all 216 individual molecules for a typical inherent structure transition. In fact, there is a relatively small set of molecules with a large displacement. A snapshot of the eight molecules with the largest displacement is shown in Fig. 4. Interestingly, this set of molecules forms a cluster of bonded molecules. Indeed, for all cases studied, the set of molecules which displace most forms a cluster of bonded molecules. The observed clustering phenomenon characterizes the IS transitions in water and can be interpreted as the analog of the string-like motion observed in simple atomistic liquids [100], connected to the presence of dynamical heterogeneities [108–111]. Similar results were found by Ohmine et al. using the TIP4P and TIPS2 models for water [112].

To characterize the distribution of individual molecular displacements between different inherent structures more carefully, Fig. 3(b) shows the distribution of displacements u of the oxygen atoms $P(u)$. Note that $P(u)$ was previously studied by Schröder et al. for a binary Lennard–Jones mixture [100].




Fig. 4. Snapshot of the system in one IS. Only the eight molecules with displacement larger than 0.025 nm [Fig. 3(a)] are shown here. Hydrogen-bonded molecules are connected by tubes. Note that all 8 molecules are nearby and form a cluster, which unlike the Lennard–Jones case, are bounded and less string-like. Adapted from Ref. [105].

Analysis of the changes in hydrogen bond connectivity associated with inherent structure changes reveals that these transitions are associated with the breaking and reformation of hydrogen bonds.

Giovambattista et al. [105] further shows that the transitions associated with an increase in the energy correspond to the breaking of linear bonds and to the simultaneous formation of bifurcated bonds [113,114]. Similarly, the transitions associated with a decrease in the energy correspond to the breaking of bifurcated bonds and to the simultaneous formation of linear bonds. This result supports the hypothesis that the linear to bifurcated transition can be considered as an elementary step in the rearrangement of the hydrogen bond network.

Acknowledgements

We thank M. Barbosa, G. Franzese, S.C. Glotzer, T. Keyes, W. Kob, G. Malescio, S. Mossa, P. Netz, I. Ohmine, G. Ruocco, R. Sadr, S. Sastry, T.B. Schröder, A. Skibinsky and M. Yamada for helpful discussions, and NSF Grant CHE-0096892 for support. FS acknowledges support from MURST (PRIN 2000) and INFN (PRA-HOP

and Initiative Parallel Computing), and the conference organizers for inviting this work to be presented at Statphys-22.

References

- [1] P.G. Debenedetti, *Metastable Liquids*, Princeton University Press, Princeton, NJ, 1996.
- [2] O. Mishima, H.E. Stanley, *Nature* 396 (1998) 329.
- [3] P. Ball, *Life's Matrix: A Biography of Water*, Farrar Straus and Giroux, New York, 2000.
- [4] V. Brazhkin, S.V. Buldyrev, V. Ryzhov, H.E. Stanley (Eds.), *New Kinds of Phase Transition Phenomena*, Proceedings of the Volga River NATO Advanced Research Workshop, Kluwer, Dordrecht, 2002.
- [5] H.E. Stanley, *J. Phys. A* 12 (1979) L329.
- [6] H.E. Stanley, J. Teixeira, *J. Chem. Phys.* 73 (1980) 3404.
- [7] H.E. Stanley, J. Teixeira, A. Geiger, R.L. Blumberg, *Physica A* 106 (1981) 260.
- [8] A. Geiger, H.E. Stanley, *Phys. Rev. Lett.* 49 (1982) 1749.
- [9] H.E. Stanley, R.L. Blumberg, A. Geiger, *Phys. Rev. B* 28 (1983) 1626.
- [10] H.E. Stanley, R.L. Blumberg, A. Geiger, P. Mausbach, J. Teixeira, *J. Phys.* 45 (1984) C7[3].
- [11] L. Bosio, J. Teixeira, H.E. Stanley, *Phys. Rev. Lett.* 46 (1981) 597.
- [12] R.L. Blumberg, H.E. Stanley, A. Geiger, P. Mausbach, *J. Chem. Phys.* 80 (1984) 5230.
- [13] E. Shiratani, M. Sasai, *J. Chem. Phys.* 104 (1996) 7671.
- [14] P.H. Poole, F. Sciortino, U. Essmann, H.E. Stanley, *Nature* 360 (1992) 324.
- [15] T. Andrews, *Philos. Trans.* 159 (1869) 575.
- [16] O. Mishima, H.E. Stanley, *Nature* 392 (1998) 164.
- [17] O. Mishima, *Phys. Rev. Lett.* 85 (2000) 334.
- [18] P.W. Bridgman, *Proc. Am. Acad. Arts Sci.* 47 (1912) 441.
- [19] L.F. Evans, *J. Appl. Phys.* 38 (1967) 4930.
- [20] R.S. Smith, B.D. Kay, *Nature* 398 (1999) 788.
- [21] K.P. Stevenson, G.A. Kimmel, Z. Dohnalek, R.S. Smith, B.D. Kay, *Science* 283 (1999) 1505.
- [22] M.-C. Bellissent-Funel, *Europhys. Lett.* 42 (1998) 161.
- [23] M.C. Bellissent-Funel, L. Bosio, *J. Chem. Phys.* 102 (1995) 3727.
- [24] A.K. Soper, M.A. Ricci, *Phys. Rev. Lett.* 84 (2000) 2881.
- [25] M.A. Ricci, A.K. Soper, *Physica A* 304 (2002) 43.
- [26] P.H. Poole, F. Sciortino, T. Grande, H.E. Stanley, C.A. Angell, *Phys. Rev. Lett.* 73 (1994) 1632.
- [27] C.F. Tejero, M. Baus, *Phys. Rev. E* 57 (1998) 4821.
- [28] F.W. Starr, S. Sastry, E. La Nave, A. Scala, H.E. Stanley, F. Sciortino, *Phys. Rev. E* 63 (2001) 041201.
- [29] A. Scala, F.W. Starr, E. La Nave, H.E. Stanley, F. Sciortino, *Phys. Rev. E* 62 (2000) 8016.
- [30] M. Yamada, S. Mossa, H.E. Stanley, F. Sciortino, *cond-mat/0202094*.
- [31] S. Sastry, P.G. Debenedetti, F. Sciortino, H.E. Stanley, *Phys. Rev. E* 53 (1996) 6144.
- [32] E. Shiratani, M. Sasai, *J. Chem. Phys.* 108 (1998) 3264.
- [33] H. Tanaka, *J. Chem. Phys.* 105 (1996) 5099.
- [34] O. Mishima, *J. Chem. Phys.* 100 (1994) 5910.
- [35] H. Tanaka, *Nature* 380 (1996) 328.
- [36] P.A. Netz, F.W. Starr, H.E. Stanley, M.C. Barbosa, *J. Chem. Phys.* 115 (2001) 344.
- [37] F.W. Starr, C.A. Angell, H.E. Stanley, *cond-mat/9903451*.
- [38] P.C. Hemmer, G. Stell, *Phys. Rev. Lett.* 24 (1970) 1284.
- [39] G. Stell, P.C. Hemmer, *J. Chem. Phys.* 56 (1972) 4274.
- [40] C.K. Hall, G. Stell, *Phys. Rev. A* 7 (1973) 1679.
- [41] M.R. Sadr-Lahijany, A. Scala, S.V. Buldyrev, H.E. Stanley, *Phys. Rev. Lett.* 81 (1998) 4895.
- [42] M.R. Sadr-Lahijany, A. Scala, S.V. Buldyrev, H.E. Stanley, *Phys. Rev. E* 60 (1999) 6714.
- [43] A. Scala, M.R. Sadr-Lahijany, N. Giovambattista, S.V. Buldyrev, H.E. Stanley, *Phys. Rev. E* 63 (2001) 041202.

- [44] A. Scala, M. Reza Sadr-Lahijany, N. Giovambattista, S.V. Buldyrev, H.E. Stanley, *J. Statist. Phys.* 100 (2000) 97.
- [45] E.A. Jagla, *Phys. Rev. E* 58 (1998) 1478.
- [46] E.A. Jagla, *J. Chem. Phys.* 111 (1999) 8980.
- [47] E.A. Jagla, *Phys. Rev. E* 63 (2001) 061509.
- [48] G. Franzese, G. Malescio, A. Skibinsky, S.V. Buldyrev, H.E. Stanley, *Nature* 409 (2001) 692.
- [49] G. Malescio, G. Pellicane, *Phys. Rev. E* 63 (2001) 020501.
- [50] F.H. Stillinger, T. Head-Gordon, *Phys. Rev. E* 47 (1993) 2484.
- [51] F.H. Stillinger, D.K. Stillinger, *Physica A* 244 (1997) 358.
- [52] T. Head-Gordon, F.H. Stillinger, *J. Chem. Phys.* 98 (1993) 3313.
- [53] N. Guisoni, V.B. Henriques, *J. Chem. Phys.* 115 (2001) 5238.
- [54] S.V. Buldyrev, G. Franzese, N. Giovambattista, G. Malescio, M.R. Sadr-Lahijany, A. Scala, A. Skibinsky, H.E. Stanley, *Physica A* 304 (2002) 23.
- [55] F.W. Starr, M.-C. Bellissent-Funel, H.E. Stanley, *Phys. Rev. E* 60 (1999) 1084.
- [56] M. Sasai, *Physica* 285 (2000) 315.
- [57] M. Canpolat, F.W. Starr, M.R. Sadr-Lahijany, A. Scala, O. Mishima, S. Havlin, H.E. Stanley, *Chem. Phys. Lett.* 294 (1998) 9.
- [58] G.E. Walrafen, *J. Chem. Phys.* 40 (1964) 3249.
- [59] W.B. Monosmith, G.E. Walrafen, *J. Chem. Phys.* 81 (1984) 669.
- [60] P.H. Poole, M. Hemmati, C.A. Angell, *Phys. Rev. Lett.* 79 (1997) 2281.
- [61] I. Saika-Voivod, F. Sciortino, P.H. Poole, *Phys. Rev. E* 63 (2001) 011202.
- [62] E.G. Ponyatovskii, *JETP Lett.* 66 (1997) 281.
- [63] E.G. Ponyatovskii, O.I. Bakalov, *Mater. Sci. Rep.* 8 (1992) 147.
- [64] M. Togaya, *Phys. Rev. Lett.* 79 (1997) 2474.
- [65] J. Glosli, F.H. Ree, *Phys. Rev. Lett.* 82 (1999) 4659.
- [66] S. Sastry, C.A. Angell, 2002, preprint.
- [67] Y. Katayama, T. Mizutani, W. Utsumi, O. Shimomur, M. Yamakata, K.-i. Funakoshi, *Nature* 403 (2000) 170.
- [68] G. Ruocco, et al., 2002, preprint.
- [69] S. Sastry, P.G. Debenedetti, F.H. Stillinger, *Nature* 393 (1998) 554.
- [70] F. Sette, M.H. Krish, C. Masciovecchio, G. Ruocco, G. Monaco, *Science* 280 (1998) 1550.
- [71] P. Lunkenheimer, A. Pimenov, A. Loidl, *Phys. Rev. Lett.* 78 (1997) 2995.
- [72] W. Götzke, *J. Phys.: Condens. Matter* 11 (1999) A1.
- [73] K. Binder, et al., in: M. Rubí, C. Perez-Vicente (Eds.), *Complex Behavior of Glassy Systems*, Springer, Berlin, 1997.
- [74] W. Kob, *J. Phys.: Condens. Matter* 11 (1999) R85.
- [75] M. Mézard, G. Parisi, *J. Phys.: Condens. Matter* 11 (1999) A157.
- [76] R. Speedy, *J. Chem. Phys.* 110 (1999) 54559.
- [77] R. Speedy, *J. Chem. Phys. B* 103 (1999) 4060.
- [78] M. Shultz, *Phys. Rev. B* 57 (1998) 11319.
- [79] D.C. Wallace, *Phys. Rev. E* 56 (1997) 4179.
- [80] P.G. Debenedetti, F.H. Stillinger, *Nature* 410 (2001) 259.
- [81] I. Saika-Voivod, P.H. Poole, F. Sciortino, *Nature* 412 (2001) 514.
- [82] S. Sastry, *Nature* 409 (2001) 164.
- [83] F.H. Stillinger, T.A. Weber, *Phys. Rev. A* 28 (1983) 2408.
- [84] R.O. Davies, G.O. Jones, *Adv. Phys.* 2 (1953) 370.
- [85] M. Goldstein, *J. Chem. Phys.* 51 (1969) 3728.
- [86] T. Keyes, *J. Phys. Chem. A* 101 (1997) 2921.
- [87] W. Li, T. Keyes, *J. Chem. Phys.* 111 (1999) 5503.
- [88] T. Keyes, *J. Chem. Phys.* 101 (1994) 5081.
- [89] J.D. Gezelter, E. Rabani, B.J. Berne, *J. Chem. Phys.* 107 (1997) 4618.
- [90] E. La Nave, A. Scala, F.W. Starr, H.E. Stanley, F. Sciortino, *Phys. Rev. E* 64 (2001) 036102.
- [91] S. Bembek, B. Laird, *Phys. Rev. Lett.* 74 (1995) 936.
- [92] W. Li, T. Keyes, F. Sciortino, *J. Chem. Phys.* 108 (1998) 252.

- [93] E. La Nave, A. Scala, F.W. Starr, F. Sciortino, H.E. Stanley, *Phys. Rev. Lett.* 84 (2000) 4605.
- [94] E. La Nave, H.E. Stanley, F. Sciortino, *Phys. Rev. Lett.* 88 (2002) 035501.
- [95] L. Angelani, G. Ruocco, A. Scala, F. Sciortino, *Phys. Rev. Lett.* 85 (2000) 5356.
- [96] F. Sciortino, P. Tartaglia, *Phys. Rev. Lett.* 78 (1997) 2385.
- [97] M. Sasai, *Proceedings of the International Conference on Slow Dynamics and Glass Transition*, Bangalore, India, 6–10 January 2002.
- [98] T. Keyes, *Phys. Rev. E* 62 (2000) 7905.
- [99] G. Adam, J.H. Gibbs, *J. Chem. Phys.* 43 (1965) 139.
- [100] T.B. Schröder, S. Sastry, J.C. Dyre, S.C. Glotzer, *J. Chem. Phys.* 112 (2000) 9834.
- [101] A. Heuer, *Phys. Rev. Lett.* 78 (1997) 4051.
- [102] F. Sciortino, W. Kob, P. Tartaglia, *Phys. Rev. Lett.* 83 (1999) 3214.
- [103] A. Scala, F.W. Starr, E. La Nave, F. Sciortino, H.E. Stanley, *Nature* 406 (2000) 166.
- [104] I. Ohmine, H. Tanaka, *Chem. Rev.* 93 (1993) 2545.
- [105] N. Giovambattista, F.W. Starr, F. Sciortino, S.V. Buldyrev, H.E. Stanley, *Phys. Rev. E* 65 (2002) 041502.
- [106] H.J.C. Berendsen, J.R. Grigera, T.P. Straatsma, *J. Phys. Chem.* 91 (1987) 6269.
- [107] F.W. Starr, F. Sciortino, H.E. Stanley, *Phys. Rev. E* 60 (1999) 6757.
- [108] M. Hurley, P. Harrowell, *Phys. Rev. E* 52 (1995) 1694.
- [109] W. Kob, C. Donati, S.J. Plimpton, P.H. Poole, S.C. Glotzer, *Phys. Rev. Lett.* 79 (1997) 2827.
- [110] C. Donati, J.F. Douglas, W. Kob, S.J. Plimpton, P.H. Poole, S.C. Glotzer, *Phys. Rev. Lett.* 80 (1998) 2338.
- [111] B. Doliwa, A. Heuer, *Phys. Rev. Lett.* 80 (1998) 4915.
- [112] I. Ohmine, S. Saito, *Accel. Chem. Res.* 32 (1999) 741.
- [113] F. Sciortino, A. Geiger, H.E. Stanley, *Nature* 354 (1991) 218.
- [114] F. Sciortino, A. Geiger, H.E. Stanley, *J. Chem. Phys.* 96 (1992) 3857.
- [115] J.C. Dyre, *Phys. Rev. E* 59 (1999) 2458.