

Atmospheric Retention of Man-made CO₂ Emissions

Bert W. Rust¹

¹National Institute of Standards and Technology
Gaithersburg, Maryland 20899-8910, USA
e-mail: bert.rust@nist.gov

Keywords: CO₂ Emissions, Atmospheric CO₂, Global Temperatures, Global Warming

Abstract. *Rust and Thijsse [9, 11] have shown that global annual average temperature anomalies $T(t_i)$ vary linearly with atmospheric CO₂ concentrations $c(t_i)$. The $c(t_i)$ can be related to man-made CO₂ emissions $F(t_i)$ by a linear regression model whose solution vector gives the unknown retention fractions $\gamma(t_i)$ of the $F(t_i)$ in the atmosphere. Gaps in the $c(t_i)$ record make the system underdetermined, but the constraints $0 \leq \gamma(t_i) \leq 1$ make estimation tractable. The $\gamma(t_i)$ are estimated by two methods: (1) assuming a finite harmonic expansion for $\gamma(t)$, and (2) using a constrained least squares algorithm [8] to compute average values of $\gamma(t)$ on suitably chosen time subintervals. The two methods give consistent results and show that $\gamma(t)$ declined non-monotonically from ≈ 0.6 in 1850 to ≈ 0.4 in 2000.*

1 Atmospheric CO₂ and Global Temperatures

The upper plot in Figure 1 shows an optimal regression spline [11] fit $c(t)$ to the record of atmospheric CO₂ concentrations obtained by combining atmospheric measurements at the South Pole [5] with reconstructions from Antarctic ice cores [1, 7]. Although the latter display larger random variations than the former, the two records are consistent in the years where they overlap. The spline $c(t)$ was used to model the Climatic Research Unit's record [4] of annual average global surface temperature anomalies shown in the lower plot. The solid curve was obtained by fitting the model

$$T(t) = T_0 + \eta [c(t) - 277.04] + A \sin \left[\frac{2\pi}{\tau} (t + \phi) \right],$$

with free parameters T_0 , η , A , τ , and ϕ . The constant 277.04 ppmv is the preindustrial CO₂ concentration estimated by averaging ice-core measurements for 1647-1764. The corresponding temperature anomaly, estimated by the fit, was $\hat{T}_0 = (-0.507 \pm .016)^\circ\text{C}$. The sinusoid, with $\hat{\tau} = (71.5 \pm 2.2)$ yr and $\hat{A} = (0.099 \pm .012)^\circ\text{C}$, represents the oscillation discovered by Schlesinger and Ramankutty [10]. It accounts for $\approx 8\%$ of the variance in the record. The baseline $T_0 + \eta [c(t) - 277.04]$, with $\hat{\eta} = (0.01039 \pm .00042)^\circ\text{C/ppmv}$, accounts for $\approx 77\%$ of the variance. It indicates a *linear relationship between global warming and increasing atmospheric CO₂*. The total warming since 1856 has been $\approx 0.9^\circ\text{C}$, and *that warming is accelerating*.




Figure 1: The relationship between atmospheric carbon dioxide and global temperatures.

2 Man-made Emissions and Atmospheric CO₂ Concentrations

Annual total man-made CO₂ emissions $F(t_i)$, for the years 1850-2000, are shown in the lower plot of Figure 2. These totals are the sums of annual fossil fuel emissions [6] and emissions due to changes in land use [3]. Taking $t_0 = 1850.0$ gives, for any later time t_i ,

$$c(t_i) = c_0 + \int_{t_0}^{t_i} \gamma(\tau) F(\tau) d\tau + \delta S(t_i),$$

where $c_0 = c(t_0)$, $\gamma(\tau)$ is the fraction remaining in the atmosphere, and $S(t)$ is a ramp function representing the Mt. Pinatubo eruption on June 15, 1991. $S(t)$ is 0 on $[1850.0, 1991.54]$, increases linearly to 1 on $[1991.54, 1993.54]$, and remains 1 thereafter. The amplitude constant δ turns out to be negative [2].

One way to estimate $\gamma(\tau)$ is to assume a harmonic expansion of the form

$$\gamma(t) = A_0 + B_0 t + \sum_{k=1}^{n_h} [A_k \cos(2\pi kt/150) + B_k \sin(2\pi kt/150)], \quad 1850 \leq t \leq 2000,$$

with n_h chosen so that $2n_h + 4$ is less than the number of observed $c(t_i)$. Substituting the expansion into the above integral leads to linear least squares estimates for c_0 , δ , and the A_k and B_k . Choosing n_h too large produces implausibly oscillating estimates which violate the constraints $0 \leq \gamma(t) \leq 1$. The estimate for $n_h = 2$ is plotted as a smooth curve in Figure 3. The corresponding estimate for $c(t)$ is shown as a dashed curve in the upper plot of Figure 2.

Another approach, which seeks a vector approximation $\gamma(\tau_j)$, is to approximate the integral using a rectangular quadrature rule with $\Delta\tau = 1$ year. This gives a linear regression model

$$\mathbf{c}(t_i) = [\mathbf{1}, \mathbf{F}(t_i, \tau_j), \mathbf{s}] \begin{bmatrix} c_0 \\ \gamma(\tau_j) \\ \delta \end{bmatrix} + \boldsymbol{\epsilon}(t_i), \quad \boldsymbol{\epsilon}(t_i) \sim N(\mathbf{0}, \boldsymbol{\Sigma}^2),$$




Figure 2: Man-made CO₂ emissions and CO₂ concentration in the atmosphere.

where $\mathbf{F}(t_i, \tau_j)$ is a matrix of columns formed from zeroes and the values $F(\tau_j)$, \mathbf{s} is a vector representation of $S(t_i)$, and $\boldsymbol{\epsilon}(t_i)$ is a vector of measurement errors. The covariance matrix $\boldsymbol{\Sigma}^2$ was estimated by assuming constant variance for the ice core records and a different constant variance for the atmospheric measurements. These two constants were estimated from deviations of the measurements from optimal regression spline fits [11] to the two sets of data.

Because of gaps in the $c(t_i)$ record, the matrix $[\mathbf{1}, \mathbf{F}(t_i, \tau_j), \mathbf{s}]$ has more columns than rows, but estimation is possible because $0 \leq \gamma(\tau_j) \leq 1$. Even so, the estimates $\hat{\gamma}(\tau_j)$ oscillate wildly between those bounds, so it was necessary to estimate average values of $\gamma(\tau)$ on various time subintervals of [1850, 2000]. O’Leary’s BRAKET-LS algorithm [8] was used to compute 95 % confidence intervals for 6 nonoverlapping 25-year subintervals shown in Figure 3. The pre-1925 uncertainties are large, but the bounds give good agreement with the $\hat{\gamma}(t)$ estimated from the harmonic expansion. Combining the fit of the corresponding $\hat{c}(t)$ to the measurements in Figure 2 with the results in Figure 1 suggests that *man-made CO₂ emissions are a major contributor to global warming*.

REFERENCES

- [1] D. M. Etheridge, L. P. Steele, R. L. Langerfelds, R. J. Francey, J. M. Barnola and V. I. Morgan. *Historical CO₂ Records from the Law Dome DE08, DE08-2, and DSS Ice Cores*. Trends: A Compendium of Data on Global Change, CDIAC, Oak Ridge, TN, (1998), <http://cdiac.ornl.gov/trends/co2/lawdome.html>.
- [2] L. Gu, D. D. Baldocchi, S. C. Wofsy, J. W. Munger, J. J. Michalsky, S. P. Urbanski, and T. A. Boden. *Response of a Deciduous Forest to the Mount Pinatubo Eruption: Enhanced Photosynthesis*. Science, Vol. 299, (2003), pp. 2035-2038.
- [3] R. A. Houghton. *Carbon Flux to the Atmosphere From Land-Use Changes: 1850-2005*. Trends: A Compendium of Data on Global Change, CDIAC, Oak Ridge, TN, (2008),




Figure 3: Fraction of CO₂ emissions remaining in the atmosphere.

<http://cdiac.ornl.gov/trends/landuse/houghton/houghton.html>.

- [4] P. D. Jones, M. New, D. E. Parker, S. Martin, and I. G. Rigor. *Surface Air Temperature and its Variations Over the Last 150 Years*. Reviews of Geophysics, Vol. 37, (1999), pp. 173-199, <http://www.cru.uea.ac.uk/cru/data/temperature/>.
- [5] C. D. Keeling and T. P. Whorf. *Atmospheric CO₂ Concentrations Derived from Flask Air Samples at Sites in the SIO Network*. Trends: A Compendium of Data on Global Change, CDIAC, Oak Ridge, TN, (2004), <http://cdiac.ornl.gov/trends/co2/sio-keel-flask/sio-keel-flaskspo.html>.
- [6] G. Marland, T. A. Boden, and R. J. Andres. *Global, Regional and National CO₂ Emissions*. Trends: A Compendium of Data on Global Change, CDIAC, Oak Ridge, TN, (2008), http://cdiac.ornl.gov/trends/emis/tre_glob.html.
- [7] A. Neftel, H. Friedli, E. Moor, H. Lötscher, H. Oeschger, U. Singenthaler, and B. Stauffer. *Historical CO₂ Record from the Siple Station Ice Core*. Trends: A Compendium of Data on Global Change, CDIAC, Oak Ridge, TN, (1994), <http://cdiac.ornl.gov/trends/co2/siple.html>.
- [8] D. P. O’Leary and B. W. Rust. *Confidence Intervals for Inequality-Constrained Least Squares Problems, with Applications to Ill-Posed Problems*. SIAM Journal of Scientific and Statistical Computing, Vol. 7, (1986), pp. 473-489.
- [9] B. W. Rust and B. J. Thijsse. *Proceedings of the 2007 International Conference on Scientific Computing*. H. R. Arabnia, J. Y. Yang and M. Q. Yang (eds.), CSREA Press, Las Vegas, NV, (2007), pp. 10-16.
- [10] M. E. Schlesinger and N. Ramankutty. *An Oscillation in the Global Climate System of Period 65-70 Years*. Nature, Vol. 367, (1994), pp. 723-726.
- [11] B. J. Thijsse and B. W. Rust. *Freestyle Data Fitting and Global Temperatures*. Computing in Science & Engineering, Vol. 10, (2008), pp.49-59.