



## Short communication

## STR sequence analysis for characterizing normal, variant, and null alleles

Margaret C. Kline\*, Carolyn R. Hill, Amy E. Decker<sup>1</sup>, John M. Butler

National Institute of Standards and Technology, 100 Bureau Drive, M/S 8312, Gaithersburg, MD 20899, USA

## ARTICLE INFO

## Article history:

Received 5 April 2010

Received in revised form 23 July 2010

Accepted 8 September 2010

## Keywords:

Short tandem repeat

STR typing

DNA sequencing

Allele dropout

Null allele

Variant allele

## ABSTRACT

DNA sequence variation is known to exist in and around the repeat region of short tandem repeat (STR) loci used in human identity testing. While the vast majority of STR alleles measured in forensic DNA laboratories worldwide type as “normal” alleles compared with STR kit allelic ladders, a number of variant alleles have been reported. In addition, a sequence difference at a polymerase chain reaction (PCR) primer binding site in the DNA template can cause allele drop-out (i.e., a “null” or “silent” allele) with one set of primers and not with another. Our group at the National Institute of Standards and Technology (NIST) has been sequencing variant and null alleles supplied by forensic labs and cataloging this information on the NIST STRBase website for the past decade. The PCR primer sequences and strategy used for our STR allele sequencing work involving 23 autosomal STRs and 17 Y-chromosome STRs are described along with the results from 111 variant and 17 null alleles.

Published by Elsevier Ireland Ltd.

## 1. Introduction

Short tandem repeat (STR) loci are the primary genetic markers used for human identity testing and are typically examined with multiplex polymerase chain reaction (PCR) assays that measure PCR product length relative to sequenced allelic ladders [1]. A common set of about 20 autosomal STRs is routinely examined in various combinations as part of commercially available kits [2]. While the vast majority of the millions of STR alleles examined worldwide each year fall within the sizing bins used to correlate allelic ladder repeat number to the tested sample's STR repeat length, occasionally variant or “off-ladder” alleles are observed [3–9]. Our group at NIST has provided variant and null allele sequence analysis as a service to the forensic DNA community for the past decade in addition to cataloging this information on the STRBase website [59]; <http://www.cstl.nist.gov/biotech/strbase/>. As of February 2010, there were 528 variants cataloged. In addition, DNA sequence variation is known to exist in the repeat region and the flanking regions of most of the widely-used STRs [10,11]. A mismatch due to sequence variation in the DNA template can cause allele dropout when it falls within PCR primer binding sites. The resulting “null alleles” can be detected when concordance studies are performed using sets of PCR primers with different annealing positions [12–37].

In order to understand at the molecular level the reason for a variant or null allele, the DNA sample needs to be sequenced at the STR locus. If the locus in question is heterozygous and contains two distinguishable alleles, these alleles must be separated from one another prior to sequencing. Sets of new PCR primers have been developed that anneal outside all known STR kit primer positions to enable examination of the nucleotide sequences underlying these primer binding sites [Supplemental Table S1]. While there have been other reports with DNA sequence information from a few STR loci [38–58], this article describes comprehensive coverage for all the common human identity testing STR markers.

## 2. Materials and methods

## 2.1. DNA samples

DNA samples used in this study came from our in-house set of more than 650 U.S. population samples described previously [60] or were variant alleles provided by outside DNA testing laboratories. Samples with previously confirmed or suspected null alleles [26] were typically provided as DNA extracts or blood stains. All the samples were completely stripped of any personal identifiers and made anonymous prior to testing at NIST in accordance with our laboratory's institutional review board policy. DNA extractions from the bloodstains on paper were performed with Chelex as previously described [61] or by DNA IQ extraction (Promega Corp., Madison, WI) following manufacturer's protocol. If limited in size, bloodstains supplied as FTA punches were purified by manufacturer's protocol (Whatman Inc., Piscataway, NJ) and the purified punch used in the initial PCR.

\* Corresponding author. Tel.: +301 975 3134; fax: +301 975 8505.

E-mail address: [margaret.kline@nist.gov](mailto:margaret.kline@nist.gov) (M.C. Kline).<sup>1</sup> Current address: Armed Forces DNA Identification Laboratory, 1413 Research Blvd., Rockville, MD 20850, United States.

## 2.2. Primer design for PCR and sequencing primers

PCR primers were designed against each reference sequence using web-based Primer3 [62]. Typically the default parameters were used, including melting temperature ( $T_m$ ) values in the range of 57–63 °C. The region encompassing known kit primer sequences [63] was highlighted to ensure that it would be included within the PCR product upon the sequencing primer design.

Unlabeled oligonucleotides were purchased from Qiagen Operon (Alameda, CA). Table S1 contains a list of the DNA sequencing primers used in these experiments. These primers were quality controlled prior to use via UV absorbance at 260 nm using a Cary 100 UV-Vis spectrophotometer (Varian, Palo Alto, CA). All the primers were used in 1.0  $\mu\text{mol/L}$  concentrations and stored at 4 °C.

## 2.3. PCR amplification

PCR amplifications were performed as described previously [36] with approximately 5–7 ng extracted DNA amplified with locus-specific annealing temperature (see Table S1) in the range of 50–60 °C.

## 2.4. Post-amplification quantification

PCR products were evaluated for specificity and quantity following amplification using 1.2% or 2.2% agarose FlashGel System, with FlashGel QuantLadder, (Lonza Rockland Inc., Rockland, ME). For each sample, 3  $\mu\text{L}$  of product was examined.

## 2.5. Gel electrophoresis for STR allele isolation

Allele separation for heterozygote allele samples was performed on acrylamide based gels prepared to be in the range of 9–12% total acrylamide with 3% cross link of bis-acrylamide in a 50 mmol/L Tris-formate buffer pH 9.0, (50 mmol/L of formate obtained using formic acid and titrating the solution to pH9.0 using Tris Base). The gel matrix used was dependent on the size of the PCR product to be separated. Large PCR amplicons (>500 bp) were separated on the 9% gels while smaller amplicons <300 bp were separated on 12% gels. Homozygous allele samples do not require gel separation. Details of the gel casting, electrophoresis and silver staining can be found in Supplemental Protocols 1 and 2.

Using a scalpel or razor blade, the desired silver stained allele bands were excised from the gel and placed overnight in 50–100  $\mu\text{L}$  of  $\text{TE}^{-4}$  buffer (10 mmol/L Tris-HCl, 0.1 mmol/L EDTA, pH 8.0). The volume of  $\text{TE}^{-4}$  buffer was dependent on the color/concentration of the DNA obtained from the gel cut.

## 2.6. Reamplification of separated heterozygous alleles

After the gel-separated alleles from heterozygote samples sat overnight in  $\text{TE}^{-4}$  buffer, a 5  $\mu\text{L}$  aliquot was reamplified using the same PCR conditions as described [36].

## 2.7. DNA sequencing reactions

All the STR alleles from homozygous individuals and reamplified gel-separated alleles from heterozygous individuals were processed using Big Dye v 3.1 as described previously [36] to the analysis of the sequencing product.

## 2.8. Analysis on the ABI PRISM 3130xl Genetic Analyzer

Purified cycle sequencing samples were prepared for analysis on the ABI PRISM 3130xl Genetic Analyzer (Applied Biosystems) by

adding 1  $\mu\text{L}$  of sample to 12  $\mu\text{L}$  of Hi-Di formamide in a 96 well reaction tray. Although a variety of instrument configurations have been used over the years, we have found the 80 cm array and POP7 polymer (Applied Biosystems) to work optimally. Samples were injected onto the capillaries with 22 s of 1 kV and electrophoretically separated at 15 kV.

## 2.9. Data analysis

DNA sequences were analyzed using Sequence Analysis version 5.2 (Applied Biosystems) and LaserGene SeqMan Version 7.1.0 (DNASTAR inc., Madison WI). Sequence information for each STR locus was aligned against a GenBank reference sequence with the number of repeat units listed in Table S1. Reference sequences for most of the STR loci used are available at [http://www.cstl.nist.gov/biotech/strbase/seq\\_ref.htm](http://www.cstl.nist.gov/biotech/strbase/seq_ref.htm).

## 3. Results and discussion

The steps involved in our STR allele sequencing strategy are noted in Fig. 1. First, a sample is identified that is in need of sequencing to understand the molecular basis for allele dropout or to characterize the repeat region or flanking region surrounding an STR marker. The null allele samples are typically identified as part of a concordance study [28,33] where a heterozygote is observed with one primer set while a homozygote is seen with a different primer set. Variant alleles are often “off-ladder” alleles observed by a DNA testing laboratory and provided to NIST as DNA extracts or bloodstain punches for further evaluation.

The DNA samples of interest are PCR-amplified with forward and reverse sequencing primers. These sequencing primers have been designed to anneal far enough away from the STR repeat in order to generate PCR products that encompass the primer binding regions for all the known kit primers surrounding commonly used STR loci. The DNA sequencing PCR primers listed in Table S1 were all tested with homozygous samples to verify that clean sequence information could be obtained for each of the 23 autosomal and 17 Y-chromosome STR loci listed. Locus-specific annealing temperatures were used to obtain optimal results.

Following the generation of DNA sequence information using the steps outlined in Fig. 1, the forward (F) and reverse (R)





Fig. 1. Summary of the steps used in sequencing STR alleles.



**Fig. 2.** Gel separation of several DYS385 PCR amplified samples with sequencing and typing results for a 12,15 sample.

sequences are aligned with one another and then compared to a reference sequence. Differences between the generated sequences from a putative variant allele were carefully examined and then compared to other sequenced alleles as well as the reference sequence, which originates from the GenBank accession numbers noted on STRBase for each locus: [http://www.cstl.nist.gov/biotech/strbase/str\\_fact.htm](http://www.cstl.nist.gov/biotech/strbase/str_fact.htm). Fig. 2 illustrates a gel separation of the alleles present in several DYS385 PCR-amplified samples with the sequence information for each allele in a 12,15 type.

From the hundreds of normal and variant alleles sequenced over the past decade, 111 alleles from 19 autosomal and 4 Y-STR loci are included in Table S2. This information represents only a subset of more than 500 variant alleles cataloged on STRBase: [http://www.cstl.nist.gov/biotech/strbase/var\\_tab.htm](http://www.cstl.nist.gov/biotech/strbase/var_tab.htm). In addition, 17 null alleles characterized following detection through concordance studies are reported in Table S3. Further information on null alleles is also available on STRBase at <http://www.cstl.nist.gov/biotech/strbase/NullAlleles.htm>.

In Tables S2 and S3, the nomenclature for reporting variation in the flanking region follows that recommended by the International Society for Forensic Genetics (ISFG) DNA Commission [64]. For example, a deletion of a “T” 14 bp downstream of the repeat region is listed as “D14Tdel”.

#### 4. Conclusions

DNA sequence analysis is helpful to understand the molecular basis for an allele that sizes “off-ladder” or one that fails to be amplified with a specific set of PCR primers. Through funding from the National Institute of Justice, NIST has provided STR allele sequence analysis free-of-charge to the human identity testing community. The supplemental files with this article provide the DNA sequencing primers used and a selection of results obtained with normal, variant, and null alleles.

#### Acknowledgments

Funding for this work was provided by the National Institute of Justice through an interagency agreement 2008-DN-R-121 with the NIST Office of Law Enforcement Standards. Points of view in this document are those of the authors and do not necessarily represent the official position or policies of the U.S. Department of Justice. Samples used for this study were kindly provided by numerous forensic and clinical DNA laboratories from around the world. Jill Appleby helped with some early experiments on this project, and Richard Schoske designed some of the Y-chromosome STR sequencing primers. Jan Redman aided in cataloging the variant alleles in STRBase.

#### Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at [doi:10.1016/j.fsigen.2010.09.005](http://doi:10.1016/j.fsigen.2010.09.005).

#### References

- [1] J.M. Butler, *Fundamentals of Forensic DNA Typing*, Elsevier Academic Press, San Diego, 2010.
- [2] J.M. Butler, Genetics and genomics of core STR loci used in human identity testing, *J. Forensic Sci.* 51 (2006) 253–265.
- [3] S. Furedi, J. Woller, Z. Padar, M. Angyal, Y-STR haplotyping in two Hungarian populations, *Int. J. Legal Med.* 113 (1999) 38–42.
- [4] R. Klein, G. Braunschweiger, A. Junge, P. Wiegand, A very long ACTBP2 (SE33) allele, *Int. J. Legal Med.* 117 (2003) 235–236.
- [5] E.M. Dauber, G. Dorner, S. Wenda, E.M. Schwartz-Jungl, B. Glock, W. Bär, Mayr, F.W.R., Unusual FGA and D19S433 off-ladder alleles and other allelic variants at the STR loci D8S1132, vWA, D18S51 and ACTBP2 (SE33), *Forensic Sci. Int. Genet. Suppl. Ser. 1* (2008) 109–111.
- [6] V. Onofri, M. Pesaresi, F. Alessandrini, C. Turchi, A. Tagliabracci, D16S539 micro-variant or D2S1338 off-ladder allele? A case report about a range overlapping between two loci, *Forensic Sci. Int. Genet. Suppl. Ser. 1* (2008) 123–124.
- [7] A. Tamura, M. Iwata, K. Tsuboi, T. Miyazaki, H. Nishio, K. Suzuki, Sequence analysis of two de novo mutation alleles at the Y-STR locus, *Sci. Int. Genet. Suppl. Ser. 1* (2008) 250–251.
- [8] E.M. Dauber, E.M. Schwartz-Jungl, S. Wenda, G. Dorner, B. Glock, W.R. Mayr, Further allelic variation at the STR-loci ACTBP2 (SE33), D3S1358, D8S1132, D18S51 and D21S11, *Forensic Sci. Int. Genet. Suppl. Ser. 2* (2009) 41–42.
- [9] A. Morales-Valverde, S. Silva-De La Fuente, G.G. Nuñez-Rivas, M. Espinoza-Esquivel, Characterisation of 12 new alleles in the STR system D18S51, *Forensic Sci. Int. Genet. Suppl. Ser. 2* (2009) 43–44.
- [10] B. Egyed, S. Furedi, M. Angyal, L. Boutrand, A. Vandenberghe, J. Woller, Z. Padar, Analysis of eight STR loci in two Hungarian populations, *Forensic Sci. Int.* 113 (2000) 25–27.
- [11] H. Oberacher, F. Pitterl, G. Huber, H. Niederstätter, M. Steinlechner, W. Parson, Increased forensic efficiency of DNA fingerprints through simultaneous resolution of length and nucleotide variability by high-performance mass spectrometry, *Hum. Mutat.* 29 (2008) 427–432.
- [12] L. Gusmão, A. Amorim, M.J. Prata, L. Pereira, M.V. Lareu, A. Carracedo, Failed PCR amplifications of MBP-STR alleles due to polymorphism in the primer annealing region, *Int. J. Legal Med.* 108 (1996) 313–315.
- [13] M.C. Kline, B. Jenkins, S. Rodgers, Non-amplification of a vWA allele, *J. Forensic Sci.* 43 (1998) 250–251.
- [14] R.W. Allen, J. Pritchard, J. Harmon, S. Floyd, Phenotypic differences at the HUMvWA locus amplified with different STR kits, *J. Forensic Sci.* 45 (2000) 1166–1167.
- [15] E.A. Cotton, R.F. Allsop, J.L. Guest, R.R. Frazier, P. Koumi, I.P. Callow, A. Seager, R.L. Sparkes, Validation of the AMPFISTR SGM plus system for use in forensic casework, *Forensic Sci. Int.* 112 (2000) 151–161.
- [16] S.A. Greenspoon, P.J. Lytle, S.A. Turek, J.M. Rolands, M.A. Scarpetta, C.D. Carr, Validation of the PowerPlex 1.1 loci for use in human identification, *J. Forensic Sci.* 45 (2000) 677–683.
- [17] C. Alves, A. Amorim, L. Gusmão, L. Pereira, VWA STR genotyping: further inconsistencies between Perkin-Elmer and Promega kits, *Int. J. Legal Med.* 115 (2001) 97–99.
- [18] B. Budowle, A. Masibay, S.J. Anderson, C. Barna, L. Biega, S. Brenneke, B.L. Brown, J. Cramer, G.A. DeGroot, D. Douglas, B. Duceaman, A. Eastman, R. Giles, J. Hamill, D.J. Haase, D.W. Janssen, T.D. Kupferschmid, T. Lawton, C. Lemire, B. Llewellyn, T. Moretti, J. Neves, C. Palaski, S. Schueler, J. Sgueglia, C. Sprecher, C. Tomsey, D. Yet, STR primer concordance study, *Forensic Sci. Int.* 124 (2001) 47–54.
- [19] B. Budowle, C.J. Sprecher, Concordance study on population database samples using the PowerPlex 16 kit and AmpFISTR Profiler Plus kit and AmpFISTR COfiler kit, *J. Forensic Sci.* 46 (2001) 637–641.
- [20] G.R. Han, E.S. Song, J.J. Hwang, Non-amplification of an allele of the D8S1179 locus due to a point mutation, *Int. J. Legal Med.* 115 (2001) 45–47.
- [21] C.S. Tomsey, M. Kurtz, F. Kist, M. Hockensmith, P. Call, Comparison of PowerPlex 16, PowerPlex 1.1/2.1, and ABI AmpFISTR Profiler Plus/COfiler for forensic use, *Croat. Med. J.* 42 (2001) 239–243.
- [22] V.K. Kashyap, S. Guha, R. Trivedi, Concordance study on 15 STR loci in three major populations of Himalayan State Sikkim, *J. Forensic Sci.* 47 (2002) 1163–1167.
- [23] M.S. Nelson, E.N. Levedakou, J.R. Matthews, B.E. Early, D.A. Freeman, C.A. Kuhn, C.J. Sprecher, A.S. Amin, K.C. McElfresh, J.W. Schumm, Detection of a primer-binding site polymorphism for the STR locus D16S539 using the Powerplex 1.1 system and validation of a degenerate primer to correct for the polymorphism, *J. Forensic Sci.* 47 (2002) 345–349.
- [24] P. Grubwieser, R. Muhlmann, W. Parson, New sensitive amplification primers for the STR locus D2S1338 for degraded casework DNA, *Int. J. Legal Med.* 117 (2003) 185–188.
- [25] A. Amorim, C. Alves, L. Pereira, L. Gusmão, Genotyping inconsistencies and null alleles using AmpFISTR Identifier and PowerPlex 16 kits, *Int. Congress Ser. (Progress in Forensic Genetics 10)* 1261 (2004) 176–178.
- [26] T.M. Clayton, S.M. Hill, L.A. Denton, S.K. Watson, A.J. Urquhart, Primer binding site mutations affecting the typing of STR loci contained within the AMPFISTR SGM Plus kit, *Forensic Sci. Int.* 139 (2004) 255–259.

- [27] M. Delamoye, C. Duverneuil, K. Riva, M. Leterreux, S. Taieb, P. De, Mazancourt, False homozygosities at various loci revealed by discrepancies between commercial kits: implications for genetic databases, *Forensic Sci. Int.* 143 (2004) 47–52.
- [28] J. Drabek, D.T. Chung, J.M. Butler, B.R. McCord, Concordance study between miniplex assays and a commercial STR typing kit, *J. Forensic Sci.* 49 (2004) 859–860.
- [29] S.W. Forrest, T.D. Kupferschmid, B.C. Hendrickson, T. Judkins, D.J. Petersen, T. Scholl, Two rare novel polymorphisms in the D8S1179 and D13S317 markers and a method to mitigate their impact on human identification, *Croat. Med. J.* 45 (2004) 457–460.
- [30] S.A. Greenspoon, J.D. Ban, L. Pablo, C.A. Crouse, F.G. Kist, C.S. Tomsey, A.L. Glessner, L.R. Mihalacki, T.M. Long, B.J. Heidebrecht, C.A. Braunstein, D.A. Freeman, C. Sobersalski, B. Nathan, A.S. Amin, E.K. Douglas, J.W. Schumm, Validation and implementation of the PowerPlex 16 BIO System STR multiplex for forensic casework, *J. Forensic Sci.* 49 (2004) 71–80.
- [31] B.C. Hendrickson, B. Leclair, S. Forrest, J. Ryan, B.E. Ward, D. Petersen, T.D. Kupferschmid, T. Scholl, Accurate STR allele designations at the FGA and vWA loci despite primer site polymorphisms, *J. Forensic Sci.* 49 (2004) 250–254.
- [32] P. Kuzniar, R. Ploski, STR data for the PowerPlex 16 loci in a population from Central Poland, *Forensic Sci. Int.* 139 (2004) 261–263.
- [33] C.R. Hill, M.C. Kline, J.J. Mulero, R.E. Lagacé, C.W. Chang, L.K. Hennessy, J.M. Butler, Concordance study between the AmpFISTR MiniFiler PCR amplification kit and conventional STR typing kits, *J. Forensic Sci.* 52 (2007) 870–873.
- [34] U. Ricci, G. Melean, C. Robino, M. Genuardi, A single mutation in the FGA locus responsible for false homozygosities and discrepancies between commercial kits in an unusual paternity test case, *J. Forensic Sci.* 52 (2007) 393–396.
- [35] E.M. Dauber, B. Glock, W.R. Mayr, Two examples of null alleles at the D19S433 locus due to the same 4 bp deletion in the presumptive primer binding site of the AmpFISTR Identifiler kit, *Forensic Sci. Int. Genet. Suppl. Ser. 1* (2008) 107–108.
- [36] C.R. Hill, M.C. Kline, M.D. Coble, J.M. Butler, Characterization of 26 miniSTR loci for improved analysis of degraded DNA samples, *J. Forensic Sci.* 53 (2008) 73–80.
- [37] C.R. Hill, D.L. Duewer, M.C. Kline, C.J. Sprecher, R.S. McLaren, D.R. Rabbach, B.E. Krenke, M.G. Ensenberger, P.M. Fulmer, D.R. Storts, J.M. Butler, Concordance and population studies along with stutter and peak height ratio analysis for Powerplex ESX 17 and ESI 17 systems, *Forensic Sci. Int. Genetic* (2010), doi:10.1016/j.fsigen.2010.30.014.
- [38] M.A. Marino, J.M. Devaney, J.K. Smith, J.E. Girard, Sequencing using capillary electrophoresis of short tandem repeat alleles separated and purified by high performance liquid chromatography, *Electrophoresis* 19 (1998) 108–118.
- [39] L. Boutrand, B. Egyed, S. Furedi, N. Mommers, G. Mertens, A. Vandenbergh, Variations in primer sequences are the origin of allele drop-out at loci D13S317 and CD4, *Int. J. Legal Med.* 114 (2001) 295–297.
- [40] K. Lazaruk, J. Wallin, C. Holt, T. Nguyen, P.S. Walsh, Sequence variation in humans and other primates at six short tandem repeat loci used in forensic identity testing, *Forensic Sci. Int.* 119 (2001) 1–10.
- [41] S. Hering, R. Nixdorf, J. Dressler, Sequence variations in the primer binding regions of the highly polymorphic STR system SE33, *Int. J. Legal Med.* 116 (2002) 365–367.
- [42] C. Alves, L. Gusmão, L. Pereira, A. Amorim, Multiplex STR genotyping: comparison study, population data and new sequence information, *Int. Congress Ser. (Progress in Forensic Genetics 9)* 1239 (2003) 131–135.
- [43] T. Bhoopat, C. Hohoff, H. Steger, Identification of DYS385 Allele variants by using shorter amplicons and northern Thai haplotype data, *J. Forensic Sci.* 48 (2003) 1108–1112.
- [44] C. Leibelt, B. Budowle, P. Collins, Y. Daoudi, T. Moretti, G. Nunn, D. Reeder, R. Roby, Identification of a D8S1179 primer binding site mutation and the validation of a primer designed to recover null alleles, *Forensic Sci. Int.* 133 (2003) 220–227.
- [45] C. Alves, L. Gusmão, A. Damasceno, B. Soares, A. Amorim, Contribution for an African autosomic STR database (AmpFISTR Identifiler and PowerPlex 16 System) and a report on genotyping variations, *Forensic Sci. Int.* 139 (2004) 201–205.
- [46] M. Heinrich, M. Müller, S. Rand, B. Brinkmann, C. Hohoff, Allelic drop-out in the STR system ACTBP2 (SE33) as a result of mutations in the primer binding region, *Int. J. Legal Med.* 118 (2004) 361–363.
- [47] K. Thangaraj, A.G. Reddy, L. Singh, Mutation in the STR locus D21S11 of father causing allele mismatch in the child, *J. Forensic Sci.* 49 (2004) 99–103.
- [48] C. Turchi, M. Pesaresi, F. Alessandrini, V. Onofri, A. Arseni, A. Tagliabracchi, Unusual association of three rare alleles and a mismatch in a case of paternity testing, *J. Forensic Sci.* 49 (2004) 260–262.
- [49] P. Grubwieser, R. Mühlmann, H. Niederstätter, M. Pavlic, W. Parson, Unusual variant alleles in commonly used short tandem repeat loci, *Int. J. Legal Med.* 119 (2005) 164–166.
- [50] M. Heinrich, H. Felske-Zech, B. Brinkmann, C. Hohoff, Characterisation of variant alleles in the STR systems D2S1338, D3S1358 and D19S433, *Int. J. Legal Med.* 119 (2005) 310–313.
- [51] S. Hering, R. Nixdorf, J. Dressler, Identification of more sequence variations in the D8S1179 locus, *Forensic Sci. Int.* 149 (2005) 275–278.
- [52] D. Becker, K. Bender, J. Edelmann, F. Götz, L. Henke, S. Hering, C. Hohoff, K. Hoppe, M. Klintschar, M. Mucbe, B. Rolf, R. Szibor, V. Weirich, M. Jung, W. Brabert, New alleles and mutational events at 14 STR loci from different German populations, *Forensic Sci. Int. Genet.* 1 (2007) 232–237.
- [53] B. Glock, E.M. Dauber, W. Parson, E.M. Schwartz-Jungl, W.R. Mayr, A DYS438 null allele observed in two generations of a large family, *Forensic Sci. Int. Genet. Suppl. Ser. 1* (2008) 206–207.
- [54] T. Lederer, G. Braunschweiger, B. Dunkelmann, P. Betz, Characterization of two unusual allele variants at the STR locus ACTBP2 (SE33), *Forensic Sci. Med. Pathol.* 4 (2008) 164–166.
- [55] N. Mizuno, T. Kitayama, K. Fujii, H. Nakahara, K. Yoshida, K. Sekiguchi, N. Yonezawa, M. Nakano, K. Kasai, A D19S433 primer binding site mutation and the frequency in Japanese of the silent allele it causes, *J. Forensic Sci.* 53 (2008) 1068–1073.
- [56] N. Zahra, L.A.A. Sallam, S. Hadi, W. Goodwin, The analysis of UAE populations using AmpFISTR® Y Filer™: identification of novel and null alleles, *Forensic Sci. Int. Genet. Suppl. Ser. 1* (2008) 255–256.
- [57] G. Mertens, S. Rand, J. Butler, M. Kline, E. Jehaes, N. Mommers, E. Cardoen, I. De Bruyn, G. Leijnen, K. Van Brussel, W. Jacobs, Non-exclusion maternity case with two genetic incompatibilities, a mutation and a null allele, *Forensic Sci. Int. Genet. Suppl. Ser. 2* (2009) 224–225.
- [58] A.M. Divne, H. Edlund, M. Allen, Forensic analysis of autosomal STR markers using Pyrosequencing, *Forensic Sci. Int. Genet.* 4 (2010) 122–129.
- [59] J.M. Butler, New resources for the forensic genetics community available on the NIST STRBase website, *Forensic Sci. Int. Genet. Suppl. Ser. 1* (2008) 97–99.
- [60] J.M. Butler, R. Schoske, P.M. Vallone, J.W. Redman, M.C. Kline, Allele frequencies for 15 autosomal STR loci on U.S. Caucasian African American, and Hispanic populations, *J. Forensic Sci.* 48 (2003) 908–911.
- [61] M.C. Kline, D.L. Duewer, J.W. Redman, J.M. Butler, D.A. Boyer, Polymerase chain reaction amplification of DNA from aged blood stains: quantitative evaluation of the “suitability for purpose” of four filter papers as archival media, *Anal. Chem.* 74 (2002) 1863–1869.
- [62] S. Rozen, H. Skaletsky, Primer3 on the WWW for general users and for biologist programmers, in: S. Krawetz, S. Misener (Eds.), *Bioinformatics Methods and Protocols in the series Methods in Molecular Biology*, Humana Press, Totowa, NJ, 2000, pp. 365–386, code available at <http://fokker.wi.mit.edu/primer3/>.
- [63] B.E. Krenke, A. Tereba, S.J. Anderson, E. Buel, S. Culhane, C.J. Finis, C.S. Tomsey, J.M. Zchetti, A. Masibay, D.R. Rabbach, E.A. Amiot, C.J. Sprecher, Validation of a 16-locus fluorescent multiplex system, *J. Forensic Sci.* 47 (2002) 773–785.
- [64] L. Gusmão, J.M. Butler, A. Carracedo, P. Gill, M. Kayser, W.R. Mayr, N. Morling, M. Prinz, L. Roewer, C. Tyler-Smith, P.M. Schneider, DNA Commission of the International Society of Forensic Genetics (ISFG): an update of the recommendations on the use of Y-STRs in forensic analysis, *Forensic Sci. Int.* 157 (2006) 187–197.