

Home Search Collections Journals About Contact us My IOPscience

Feshbach enhanced *s*-wave scattering of fermions: direct observation with optimized absorption imaging

This content has been downloaded from IOPscience. Please scroll down to see the full text. 2016 New J. Phys. 18 013001 (http://iopscience.iop.org/1367-2630/18/1/013001) View the table of contents for this issue, or go to the journal homepage for more

Download details:

IP Address: 129.6.107.133 This content was downloaded on 17/12/2015 at 16:39

Please note that terms and conditions apply.

New Journal of Physics

The open access journal at the forefront of physics

Deutsche Physikalische Gesellschaft DPG

Published in partnership with: Deutsche Physikalische Gesellschaft and the Institute of Physics

PAPER

OPEN ACCESS

CrossMark

RECEIVED 30 September 2015

REVISED 20 November 2015

ACCEPTED FOR PUBLICATION

24 November 2015
PUBLISHED

17 December 2015

Content from this work may be used under the terms of the Creative Commons Attribution 3.0 licence.

Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.



Feshbach enhanced s-wave scattering of fermions: direct observation with optimized absorption imaging

D Genkina^{1,4}, L M Aycock^{1,2,4}, B K Stuhl¹, H-I Lu¹, R A Williams³ and I B Spielman¹

¹ Joint Quantum Institute, National Institute of Standards and Technology, and University of Maryland, Gaithersburg, MD 20899, USA

- ² Physics Department, Cornell University, Ithaca, NY 14850, USA
- $^{\scriptscriptstyle 3}$ $\,$ National Physical Laboratory, Hampton Road, Teddington, Middlesex, TW11 0LW, UK $\,$
 - These authors contributed equally to the work.

E-mail: ian.spielman@nist.gov

Keywords: quantum gases, atomic physics, Feshbach resonances, absorption imaging

Abstract

We directly measured the normalized *s*-wave scattering cross-section of ultracold ⁴⁰K atoms across a magnetic-field Feshbach resonance by colliding pairs of degenerate Fermi gases (DFGs) and imaging the scattered atoms. We extracted the scattered fraction for a range of bias magnetic fields, and measured the resonance location to be $B_0 = 20.206(15)$ mT with width $\Delta = 1.0(5)$ mT. To optimize the signal-to-noise ratio (SNR) of atom number in scattering images, we developed techniques to interpret absorption images in a regime where recoil induced detuning corrections are significant. These imaging techniques are generally applicable to experiments with lighter alkalis that would benefit from maximizing SNR on atom number counting at the expense of spatial imaging resolution.

1. Introduction

Feshbach resonances are widely used for tuning the interaction strength in ultracold atomic gases. In degenerate Fermi gases (DFGs), the tunability of interactions provided by Feshbach resonances has allowed for studies of the creation of molecular Bose–Einstein condensates (BECs) [1–3] as well as observation of the phase transition from the Bardeen–Cooper–Schrieffer (BCS) superconducting regime to the BEC regime at sufficiently low temperatures [4–7]. Conversely, measuring interactions as a function of controlled parameters can be used to characterize a Feshbach resonance.

A Feshbach resonance occurs when a diatomic molecular state energetically approaches the two-atom continuum [8, 9]. For a magnetic-field Feshbach resonance, a bias magnetic field defines the relative energy of the free atomic states in two hyperfine sublevels and the molecular state. Consequently, the Feshbach resonance can be accessed by changing the bias field. In cold atomic systems where only *s*-wave channels contribute to scattering, the interactions are entirely characterized by the scattering length *a*. In the simple case where there are no inelastic two-body channels, such as for the ⁴⁰K resonance discussed in this work, the effect of the resonance on the scattering length between two free atoms is [8]

$$a(B) = a_{\rm bg} \left(1 - \frac{\Delta}{B - B_0} \right),\tag{1}$$

where a_{bg} is the scattering length far from any resonance (background scattering length), Δ is the width of the resonance, and B_0 is the field value at which the resonance occurs.

The exact value of the resonant field B_0 is difficult to calculate analytically and is commonly computed via numerical models based on experimental input parameters [10–12] or determined experimentally [13, 14]. Many experimental techniques have been used to characterize Feshbach resonances. These include the observation of atom loss due to three-body inelastic scattering, measurement of re-thermalization timescales, and anisotropic expansion of a cloud upon release from a confining potential, which infer the elastic scattering cross section from collective behavior of the cloud [15–17]. More recently, precise measurements of the

molecular binding energy have been performed, which can be used in conjunction with theoretical modeling to extract the scattering cross section [18, 19].

Here direct scattering was the primary probe of the location and width of a Feshbach resonance. We collided pairs of DFGs and imaged the resulting *s*-wave scattered atoms as a function of bias magnetic field. This allowed us to observe the enhancement in scattering without relying on proxy effects. We measured the fraction of atoms scattered during the collision at different bias magnetic fields and deduced the location and width of the resonance.

In contrast to BECs, where scattering halos are readily imaged [20–22], the density of Fermi clouds is typically \approx 100 times less than that of BECs⁵, making it necessary to enhance the strength of inter-atomic interactions to directly detect the scattered atoms. In our dilute DFGs, even with the resonant enhancement of the scattering cross section, only a small fraction of the atoms was scattered. Using typical absorption imaging, direct detection of scattered atoms was difficult due to detection uncertainty that particularly affected regions of low atomic density. To optimize the signal-to-noise ratio (SNR) for low atom numbers, we absorption imaged with fairly long, high-intensity pulses—a non-standard regime—which imparted a non-negligible velocity and therefore Doppler shift to the atoms. Simulation of the absorption imaging process was necessary for an accurate interpretation of these images. Using the simulation-corrected images, we extracted the fraction of atoms scattered in our collision experiment.

This paper is divided into two parts. First, we study absorption imaging in the presence of a significant timedependent Doppler shift and show how we use our results to interpret data. Second, we describe our *s*-wave scattering experiment and extract a measure of the location and width of the Feshbach resonance in ⁴⁰K.

2. Absorption imaging in the presence of strong recoil induced detuning

Absorption imaging measures the shadow cast by an atomic ensemble in an illuminating probe laser beam with angular frequency ω_{L} . This imaging technique relies on optical transitions between ground and excited atomic states. Such atomic transitions have an energy difference $\hbar\omega_0$, and a natural transition linewidth Γ . When interacting with a laser field an atom scatters photons from the field into the vacuum modes. In the two-level atom approximation, the rate of scattering is [25]

$$\gamma_{\rm sc} = \frac{\Gamma}{2} \frac{\tilde{I}}{1+4\tilde{\delta}^2 + \tilde{I}},\tag{2}$$

where $\tilde{I} = I/I_{\text{sat}}$ is the laser intensity in units of the saturation intensity, and $\tilde{\delta} = \delta/\Gamma$ is the detuning $\delta = \omega_{\text{L}} - \omega_0$ in units of the natural linewidth.

An absorption image is obtained by shining an on- or near-resonant probe beam (generally $\delta \ll 1$) onto the atomic cloud. Some of the light is scattered by the atoms, and the shadow cast by the atoms in the probe beam, $\tilde{I}_f(x, y)$, is imaged onto a camera, as depicted in figure 1(a) (top). The probe light is reapplied with the atoms absent to calibrate the intensity $\tilde{I}_0(x, y)$ of light unaffected by the atoms (bottom).

Consider the light as it travels along the imaging axis e_z through a 3D atomic density profile $\rho(x, y, z)$. We focus on a single pixel of the camera: sensitive to a single column of atoms $\rho(z)$, integrated in x and y over the pixel, giving a single value of \tilde{I}_0 and \tilde{I}_f . Every atom scatters light according to equation (2). Therefore, the atoms further along the imaging axis e_z experience a reduced optical intensity due to attenuation of the laser field by the other atoms (figure 1(b)). The intensity change from scattering as a function of *z* is

$$\frac{\mathrm{d}\tilde{I}(z)}{\mathrm{d}z} = -\hbar\omega_{\mathrm{L}}\rho(z)\gamma_{\mathrm{sc}}(z) = -\rho(z)\sigma_{0}\frac{\tilde{I}(z)}{1+4\tilde{\delta}^{2}+\tilde{I}},\tag{3}$$

where σ_0 is the resonant scattering cross section. Integrating this equation [26] yields a straightforward relation between the observed intensities \tilde{I}_0 and \tilde{I}_f and the atomic column density $n = \int \rho(z) dz$:

$$\sigma_0 n = -\left(1 + 4\tilde{\delta}^2\right) \ln\left(I_f/I_0\right) + \tilde{I}_0 - \tilde{I}_f.$$
(4)

We call the column density deduced from this relation $\sigma_0 n^{(1)}$. When the probe intensity is much smaller than the saturation intensity, $\tilde{I}_0 \ll 1$, and the probe light is on resonance, $\tilde{\delta} = 0$, the right-hand side of equation (4) reduces to the optical depth, defined as $OD = -\ln (I_f/I_0)$ [26], giving the simple relationship $\sigma_0 n^{(0)} = OD$. In all other regimes, the optical depth is not constant and depends on the probe intensity and imaging time.

Equations (3), (4) neglect the atomic recoil momentum and the resulting Doppler shift [27]. When an atom absorbs a photon from the laser light field it acquires a momentum kick $\hbar k_r$ in the e_z direction. The associated recoil velocity is $v_r = \hbar k_r/m$, where *m* is the atomic mass and $\hbar k_r = h/\lambda$ is the recoil momentum from the laser with wavelength λ . Each re-emitted photon imparts a similar recoil momentum p_e , but over many scattering

⁵ This is not the case for recently realized erbium and dysprosium DFGs [23, 24], where strong dipolar interactions are present.







events this momentum distribution averages to zero. Therefore the atom will only acquire an average velocity per photon v_r along e_z . The variance of p_e , however, is not zero, allowing the atoms to acquire some momentum transverse to the laser field. While we ignore this correction, it results in the reduction of spatial resolution in the final image and its effect on the atomic cloud is pictured in figure 1(c).

The average atomic velocity parallel to the light field after scattering N photons is Nv_r and the laser frequency as seen by the atoms is Doppler shifted $\delta = k_r Nv_r$ from resonance. After an atom scatters $N_{\text{photons}} = \Gamma/2k_r v_r$, it gets Doppler detuned by half a linewidth. For a probe intensity of $\tilde{I}_0 = 1$, the time it takes a single atom to scatter on average that many photons is given by $t_{\text{recoil}} = N_{\text{photons}}/\gamma_{\text{sc}} = 2/k_r v_r$. For ⁴⁰K atoms imaged on the D2 transition, the case relevant for our experiment, $N_{\text{photons}} = 178$ and $t_{\text{recoil}} = 18.76 \ \mu\text{s}$ —for imaging times longer than that the recoil induced detuning correction cannot be neglected. Furthermore, this detuning varies both with imaging time t and with distance along the propagation direction e_z (figure 2). Thus, the laser's spatially varying intensity profile in the atomic cloud also depends on time:

$$\frac{d\tilde{I}(t,z)}{dz} = -\sigma_0 \rho(t,z) \frac{\tilde{I}(t,z)}{1+4\tilde{\delta}(t,z)^2 + \tilde{I}(t,z)}.$$
(5)

Assuming that the atoms do not move significantly during the imaging time (we will remove this assumption shortly), the dimensionless detuning is





$$\tilde{\delta}(t,z) = \frac{k_{\rm r} v_{\rm r}}{2\sigma_0 \rho(t,z)} \int_0^t \frac{\mathrm{d}\tilde{I}(z,\tau)}{\mathrm{d}z} \,\mathrm{d}\tau;\tag{6}$$

the relationship between the atomic density and the observed intensities is no longer straightforward. Peturbative treatments of these equations also prove insufficient (see appendix A.1).

2.1. Simulation

To obtain a relationship between the atomic density and the observed intensities in this non-standard regime, we numerically simulated the imaging process, including the recoil induced detuning. We used parameters relevant to our experiment—the D2 transition of ⁴⁰K, with $\lambda = 766.701$ nm, $\Gamma/2\pi = 6.035$ MHz, $v_r = 1.302$ cm s⁻¹ and $I_{sat} = 1.75$ mW cm⁻² [28]. The simulation obtained \tilde{I}_f as a function of imaging time *t*, atomic density $\sigma_0 n$, and probe intensity \tilde{I}_0 .

We performed two versions of this simulation. First, we took a simplistic approach where the spatial distribution of atoms did not change appreciably during the imaging time: $vt \ll I_0/(\hbar\omega_L\gamma_{sc}\rho)$ —the stationary assumption. Starting with a Gaussian density profile, we numerically integrated equations (5) and (6) and obtained a simulated optical depth for a range of input probe intensities and atomic column densities. We used the results of this simulation to check the self-consistency of the stationary atom assumption and found it to be invalid (see appendix A.2).

To account for the changing atomic distribution during the imaging pulse, we numerically simulated the classical kinetics of atoms subject to the recoil driven optical forces, and obtained a dynamics adjusted version of the simulated optical depth. We compared the optical depths predicted by each of the two simulations in the parameter range $0.3 \le t \le 100 \ \mu s$, $0.01 \le \tilde{I}_0 \le 50$ and $0.01 \le \sigma_0 n \le 2$ and found that the predicted optical depths were hardly changed by including the full time evolution (see appendix A.3). Thus, for the purposes of deducing the atomic density from experimental optical depths, the stationary atom simulation is sufficient in the experimentally relevant parameter regime we explored. However, since there is no *a priori* reason why this should be sufficient, it is possible that for some range of experimental optical depths or imaging times this correction is significant, or that it has some impact on the acquired transverse velocity of the atoms—an effect not considered in our simulations. Furthermore, we simulated a range of initial density profiles $\rho(z)$, and found their impact on the simulated OD to be negligible—the only observable is the integrated atomic density *n*, and 3D atomic densities cannot be reconstructed.

Figure 3 illustrates the effect of the recoil induced detuning correction as obtained from our simulations. In the limit of low probe intensity, $\tilde{I}_0 \ll 1$, the atomic velocities are hardly changed and the recoil induced detuning correction is negligible. In the limit of high probe intensity $\tilde{I}_0 \gg \tilde{\delta}$, even far detuned atoms will scatter light at their maximum rate and the overall absorption will again be unaffected by the correction. In the intermediate regime, there is a significant deviation between the optical depth predicted by equation (4) and the simulated optical depth, and this deviation becomes stronger with longer imaging times.



Figure 4. SNR for three different column densities after correcting for recoil induced detuning. (a) SNR as a function of imaging time for a probe intensity of $\tilde{I}_0 = 5.0$ and (b) SNR as a function of probe intensity for an imaging time of 50 μ s.

This simulation provided us with a correction procedure to interpret experimentally observed \tilde{I}_f and \tilde{I}_0 . For a given imaging time, the simulation predicted a final intensity as a function of probe intensity and atomic column density. We inverted this prediction to obtain an atomic column density given our observed intensities. For interpreting experimental images, we used the optical depths predicted by the traveling atom simulation, OD^{sim} .

2.2. SNR optimization

We added shot noise to our simulation and established optimal imaging parameters to maximize the SNR of this detection scheme. We considered Poisson noise on the detected arriving photons (i.e., photoelectrons) in both the initial and final images I_0 and I_f with standard deviation proportional to $\sqrt{q_e N_p}$, where q_e is the quantum efficiency of the camera (0.66 for our camera) and N_p is the photon number. We then propagated the shot noise in the two images using standard error propagation techniques through the correction scheme described in section 2.1 to obtain the uncertainty in a deduced column density, $\delta_{\sigma_0 n}$. We define the SNR as $\sigma_0 n/\delta_{\sigma_0 n}$.

As seen in figure 4(a), after about 40 μ s extending the imaging time no longer yields appreciable improvement in SNR. Imaging for 40 μ s as opposed to 10 μ s, where the uncorrected model is appropriate, improves the SNR by a factor of 1.5. We therefore performed the experiments described in the second section at 40 μ s imaging time. Figure 4(b) shows that the optimal probe intensity varies with the atomic column density. For low atom numbers, $\sigma_0 n \approx 0.1$, a probe intensity of $\tilde{I}_0 \approx 0.6$ is best. However, in our experiment the probe intensity had a Gaussian profile and was not uniform over the whole image. The typical probe intensities used in our experiments varied over the $\tilde{I}_0 = 0.1$ –0.7 range.

2.3. Calibration of saturation intensity

The calibration of the observed signal in units of the saturation intensity is crucial to our measurement of the column densities. Our absorption images were taken using a charge-coupled device camera. For each pixel, the camera returned an integer number of counts proportional to the radiant fluence seen by that pixel. However, the proportionality constant depended on many factors, such as the quantum efficiency of the camera, the electronic gain during the readout process and losses in the imaging system. One way to determine this proportionality constant is to experimentally calibrate the saturation intensity in counts per unit time.

To calibrate the saturation intensity in camera counts per unit time, we took absorption images of the atoms at three different imaging times (40, 100, and 200 μ s) with varying probe intensities. For each image we obtained \tilde{I}_0 and \tilde{I}_f in counts per microsecond by averaging over a few pixels in a region of constant atomic column density. We then simultaneously fit our simulated optical depth OD^{sim} to this full data set, with the atomic density $\sigma_0 n$ and I_{sat} in counts per microsecond as free parameters. As seen in figure 5, the model produced a good fit to the experimental data, and provided a calibration of the saturation intensity for our experiment.



Figure 5. The optical depth as a function of probe intensity for three imaging times: $t = 40 \ \mu s$ (black), $t = 75 \ \mu s$ (blue), $t = 100 \ \mu s$ (red). The dots represent experimental data and the curves represent the best fit of simulated data. The optimal fit parameters pictured are a $\sigma_0 n$ of 1.627(5) and saturation intensity of 29(7) counts/ μs . The dashed curve represents the theoretical prediction without recoil induced detuning corrections.

3. s-wave scattering experiment

For our measurement of the Feshbach resonance location and width, we collided two counter-propagating ⁴⁰K clouds in a spin mixture of |F = 9/2, $m_F = -9/2\rangle$ and |F = 9/2, $m_F = -7/2\rangle$ hyperfine states and observed the resulting *s*-wave halo of scattered atoms. We measured the dependence of the scattered atomic fraction on the bias magnetic field in the vicinity of the Feshbach resonance. We used this data to extract the location of the resonance at 20.206(15) mT with width 1.0(5) mT, consistent with the accepted values of 20.210(7) mT and 0.78 (6) mT [7].

3.1. Experimental procedure

Our experiment is a hybrid ⁴⁰K and ⁸⁷Rb apparatus, previously described in [29–31]. Initially, we prepared a spin polarized |F = 9/2, $m_F = -9/2$ DFG of $\approx 4 \times 10^5$ atoms of ⁴⁰K at a temperature of $T \approx 0.4 T_F$, where T_F is the Fermi temperature, in a crossed optical dipole trap with frequencies $(\omega_x, \omega_y, \omega_z)/2\pi = (39, 42, 124)$ Hz (see appendix B).

To map out the entire Feshbach resonance without the added losses associated with going through the resonance [8], we needed to create equal spin mixtures of |F = 9/2, $m_F = -9/2\rangle$ and |F = 9/2, $m_F = -7/2\rangle$ on either side of the resonance. We ramped the bias magnetic field to 19.05 mT (21.71 mT) and turned on a 42.42MHz (47.11 MHz) rf field⁶ resonant with the Zeeman splitting between the two states when preparing the mixture below (above) the Feshbach resonance. We then sinusoidally modulated the bias field at 125 Hz for 0.5 s, with a 0.14 mT amplitude, producing an equal mixture of the two hyperfine states. The depolarization allowed the fermions to re-thermalize, allowing us to further evaporate in the dipole trap [32]. These hyperfine states of ⁴⁰K were then used to study their Feshbach resonance.

After evaporation, we ramped the bias field in a two-step fashion to the desired field value near the Feshbach resonance. The two-step procedure was designed to allow us to approach the set-point quickly and avoid additional losses. This procedure used two sets of Helmholtz coils—large coils that provided the majority of the bias field but had a long inductive timescale, and smaller coils only capable of generating 0.59 mT of bias, but with a shorter inductive timescale. We approached the field using the large coils to bring the magnetic field to a set-point 0.59 mT above or below the intended bias field. We held the atoms at this field for 100 ms to allow the eddy currents induced by the large coils to settle, and then used the smaller coils to quickly change the bias field the remaining 0.59 mT. For all set-points, the data was taken approaching from both above and below the Feshbach resonance⁷.

Once at the intended bias field, we split the cloud into two spatially overlapping components with opposite momenta and observed scattering as they separated. These counterpropagating components were created using Kapitza-Dirac pulses of a 1D retro-reflected near-resonant optical lattice ($\lambda_L = 766.704 \text{ nm}$) with 8E_L depth, where $E_L = \hbar^2 k_L^2 / 2m_K$ is the lattice recoil energy and $\hbar k_L = 2\pi \hbar / \lambda$ is the recoil momentum. We rapidly pulsed

⁶ The rf intensity at these frequencies was too low for us to calibrate the coupling strength.

 $^{^7}$ An extra data point was taken on each side far from the resonance using only one approach.

this lattice on and off with a double-pulse protocol [33]. The pulse sequence was optimized to transfer most of the atoms into the $\pm 2\hbar k_{\rm L}$ momentum states. Since the initial Fermi gas had a wide momentum spread (here, $2\hbar k_{\rm L} \approx 2.5\hbar k_{\rm F}$, where $k_{\rm F}$ is the Fermi momentum), and the lattice pulsing is a momentum dependent process [34], not all the atoms were transferred into the target momentum states. We optimized our pulse times to minimize the atoms remaining in the zero momentum state.

We then released the atoms from the trap and allowed 1 ms for the two opposite momentum states to pass through each other while interacting at the magnetic field set-point. For data taken approaching the set-point from below, we then ramped down the field and imaged the atoms. For data taken approaching the set-point from above, molecules may have been created when crossing the Feshbach resonance. Therefore, we first ramped the field up to a point above the resonance to dissociate any molecules that were created and then quickly ramped the field back down and imaged the atoms. After a total time-of-flight $t_{\text{TOF}} = 6.8$ ms, we used a 40 μ s imaging pulse with $\tilde{I}_0 \approx 0.6$ at the center of the probe laser, chosen for SNR optimization as described in section 2.2.

3.2. Magnetic field calibration

The magnetic fields produced by our coils in the regime of interest were independently calibrated by rfspectroscopy on the |F = 9/2, $m_F = -9/2$ to |F = 9/2, $m_F = -7/2$ transition. We prepared a spin polarized state and ramped the large coils to variable set-points. We then illuminated the atoms with a rf field with frequency ν_{rf} and performed adiabatic rapid passage (ARP) by ramping the smaller coils 0.0284 mT in 250 ms. We applied a Stern–Gerlach pulse and imaged the atoms to measure the fractional population in the |F = 9/2, $m_F = -9/2$ and |F = 9/2, $m_F = -7/2$ states. We fit the fractional population as a function of current to a Gaussian function⁸. The center of the Gaussian corresponded to the resonant magnetic field, which was produced by the high inductance coil setpoint plus half the ARP, 0.0142 mT, with an uncertainty given by the Gaussian width. We used the Breit–Rabi formula to determine the resonant field value at ν_{rf} . We did this for five different rf frequencies, and acquired a field calibration with an uncertainty of 0.004 mT, which was included in the listed uncertainty on our measured value of B_0 .

3.3. Methods

We first processed the *s*-wave scattering images by comparing the observed OD to simulations taking into account the recoil induced detuning as described in section 2. An example of images before and after processing are shown in figure 6. The processing constituted a \approx 30% change in the column density.

We counted the fraction of atoms that experienced a single scattering event for each image. Single scattering events are easily identified, as two atoms that scatter elastically keep the same amplitude of momentum, but depart along an arbitrary direction. Therefore, an atom traveling at $2\hbar k_{\rm L}$ to the right that collides elastically with an atom traveling at $-2\hbar k_{\rm L}$ to the left will depart with equal and opposite momenta $2\hbar k_{\rm L}$ at an arbitrary angle, and after a time-of-flight sufficiently long to convert initial momentum into position, as ours was, such atoms will lie in a spherical shell, producing the scattering halo pictured in figure 7(a).

Absorption images captured the integrated column density along e_z , a projected 2D atomic distribution. To extract the radial dependence of the 3D distribution from the 2D image, we performed a standard inverse Abel transform [35]. The inverse Abel transform assumes cylindrical symmetry, which was present in our case, with the axis of symmetry along e_x , defined by the lattice. We thus obtained the atomic distribution $\rho(r, \theta)$ as a function of r, the radial distance from the scattering center, and θ , the angle between r and symmetry axis e_x , integrated over ϕ , the azimuthal angle around the x axis.

We then extracted the number of scattered atoms N_{scat} as a fraction of the total atom number N_{tot} for each image, as shown in figure 7(b). The unscattered atom number was the number of atoms in the two unscattered clouds. The number of atoms that underwent a single scattering event was the number of atoms outside the Fermi radius of the unscattered clouds, but inside the arc created by rotating the Fermi momentum k_F around the original center of the cloud (red arcs in figure 7(b)). For both the scattered and unscattered quantities, we extrapolated to include atoms that would fall outside the field of view of our camera. The atoms in the center region were not counted as they were originally in the zero momentum state and could not contribute to the scattering halo under study. We did not account for possible *s*-wave scattering between the atoms found in the zero momentum state and one of the $\pm 2k_L$ momentum states in this treatment.

Since we were in the low energy regime (the atomic momentum was much smaller than the momentum set by the van der Waals length $k_{\rm L} + k_{\rm F} \ll 1/l_{\rm vdW}$, and we were well below the *p*-wave threshold temperature [32]), the scattering cross-section was given by $\sigma = 4\pi a^2$. The scattering cross-section σ gives the probability $P_{\rm scat} = \sigma N/A$ that a single particle will scatter when incident on a cloud of atoms with a surface density of N/A, where *A* is the cross-sectional area of the cloud and *N* is the number of atoms in the cloud. In our case, each half

 $^{^8}$ Due to our low rf coupling and high noise, we did not fit to the traditional Loretzian model.











Figure 8. Normalized scattered population plotted versus bias field *B*. Green dots represent data taken coming from below the resonance, and blue dots represent the data taken coming from above the resonance. The red curve depicts the line of best fit. The regime where the scattering length is likely large enough for the atoms to behave hydrodynamically is shaded in gray, and data points in that area were excluded from the fit. Resonant field value *B*₀ from literature and as found in this work are indicated.

of the initial cloud, with atoms number $N_{tot}/2$, was incident on the other half. Thus, the number of expected scattering events was $N_{scat} = (N_{tot}/2) \sigma (N_{tot}/2) = \sigma N_{tot}^2/4A$. Assuming *A* was constant for all our data, we defined a fit parameter $b_0 = 4\pi a_{bg}^2/4A$, where a_{bg} is the background scattering length. We thus adapted equation (1) to obtain the fit function

$$\frac{N_{\text{scat}}}{N_{\text{tot}}^2} = b_0 \left(1 - \frac{\Delta}{B - B_0} \right)^2 + C,$$
(7)

where B_0 is the resonant field value and Δ is the width of the resonance, the parameters in equation (1), and the offset *C* accounts for any systematic difference in the initial and final intensity images with no atoms present.

For each value of the bias magnetic field, we took 15 nominally identical images, allowing us to compensate for shot-to-shot atom number fluctuations. We fit the fraction of scattered atoms $N_{\text{scat}}/N_{\text{tot}}$ versus the total atom number N_{tot} for each of these 15 images to a line. The slope of this fit was taken to be the value of $N_{\text{scat}}/N_{\text{tot}}^2$ at that bias magnetic field, and the variance of the fit gave the uncertainty on that data point.

3.4. Results

Our data is presented in figure 8. The red curve depicts a best fit of the model given in equation (7). The fit parameters we extracted were $\Delta = 1.0(5)$ mT and $B_0 = 20.206(15)$ mT. To obtain the fit, we used data taken by approaching the resonance from above for points above where we expected the resonance to be and data taken approaching the resonance from below for points below. We also excluded from the fit data points very near the resonance, as there the assumption $\sigma n \ll 1$ is no longer valid and the problem must be treated hydrodynamically [8]. Due to this, we could not obtain usable data very close to the resonance, explaining the large uncertainty on the resonance width.

The accepted values for the ⁴⁰K s-wave Feshbach resonance for the $|9/2, -9/2\rangle$ and $|9/2, -7/2\rangle$ states are $B_0 = 20.210(7)$ mT and $\Delta = 0.78(6)$ mT [7], which is in good agreement with our findings. Although the data without the recoil induced detuning correction were $\approx 30\%$ different from the corrected data, the optimal parameters from fitting the uncorrected data were within our uncertainties from the values listed above. The resonance location was largely unaffected by the correction, as the scattering cross-section diverges there, making it insensitive to the precise values. The width may have been affected, however the uncertainty on the fit is too large for the effect to be significant. Performing the correction was still necessary to ensure the values were proportional to the scattering cross section and to obtain a trustworthy result.

Some potential sources of systematic uncertainty that we did not account for include scattering with atoms that did not receive a momentum kick from the lattice pulsing and the impact of multiple scattering events.

4. Conclusion

We studied the effects of recoil-induced detuning on absorption images and found an imaging time that maximized SNR to be \approx 40 μ s for ⁴⁰K atoms. We used these results to directly image *s*-wave scattering halos of the Fermi gas around the \approx 20.2 mT Feshbach resonance and verify the resonance location and width. Our imaging analysis can be used in any absorption imaging application where SNR optimization is critical.

Acknowledgments

We thank Marcell Gall for helpful discussions. We thank William R McGehee and Elizabeth A Goldschmidt for a careful reading of the manuscript. This work was partially supported by the AROs Atomtronics MURI, by the AFOSRs Quantum Matter MURI, NIST, and the NSF through the PFC at the JQI. BKS acknowledges support from the National Research Council Research Associateship program. LMA acknowledges support from the National Science Foundation Graduate Research Fellowship program.

Appendix A

A.1. Peturbative treatment

By considering equations (5), (6) perturbatively in imaging time, we can obtain corrections to the column density due to recoil induced detuning to second order [36]:

$$\sigma_0 n^{(2)} = c_0 + c_1 t + c_2 t^2, \text{ where}$$
(A.1)

$$c_0 = \sigma_0 n^{(1)}, c_1 = 0, c_2 = \frac{\left(k_r v_r\right)^2}{3} \left[\frac{1}{\tilde{I}_f + 1} - \frac{1}{\tilde{I}_0 + 1} + \ln\left(\frac{\tilde{I}_f + 1}{\tilde{I}_0 + 1}\right) \right].$$
 (A.2)

However, as shown in figure A1(a), the perturbative treatment is only accurate to times up to the recoil time t_{recoil} —defined as the time it takes a single atom to get Doppler shifted from resonance by half a linewidth—after which this prediction begins to diverge. To adequately correct for the recoil induced detuning of the atoms, numerical simulation is necessary.

A.2. Stationary atom model

To solve equations (5), (6), we divided the cloud into spatial bins. In this approximation, the number of atoms in each bin was time-independent. The algorithm used is shown in algorithm 1, in which we took a Gaussian profile for our initial density distribution. We call the optical depth simulated by this algorithm the simulated optical depth OD^{sim1}.

Algorithm 1. Stationary atom model

$$\begin{split} \tilde{I}[n = 0, t] &= \tilde{I}_0 \{n \text{ is the bin index}, t \text{ is the time index}\}\\ \tilde{\delta}[n, t = 0] &= 0 \{\text{light initially resonant}\}\\ H_f &= 0 \{\text{Radiant fluence seen by camera after passing through cloud}\}\\ \text{for } t &= 0 \text{ to } t_f \text{ do } \{\text{loop over time steps}\}\\ \text{for } n &= 1 \text{ to } N \text{ do } \{\text{loop over bins}, N \text{ is total bin number}\}\\ A &= \sigma_0 \rho[n] dz \{ dz \text{ is the size of spatial step}\}\\ B &= v_t dt/(\hbar c \rho[n]) \{ dt \text{ is the size of the time step}\}\\ \tilde{I}[n, t] &= \tilde{I}[n - 1, t] - A \tilde{I}[n - 1, t]/(1 + 4 \tilde{\delta}[n, t - 1]^2 + \tilde{I}[n - 1, t]) \{\text{equation}(5)\}\\ \tilde{\delta}[n, t] &= \tilde{\delta}[n, t - 1] + B(\tilde{I}[n - 1, t] - \tilde{I}[n, t]) \{\text{equation}(6)\}\\ \text{end for}\\ H_f &= H_f + \tilde{I}[N, t] dt \{\text{collecting total fluence seen by the camera}\}\\ \text{end for}\\ \text{OD}^{\text{sim1}} &= -\ln(H_f/\tilde{I}_0 t_f) \end{split}$$

We checked the validity of our simulation in the limits where the problem is analytically solvable. In the limit where the probe intensity is much weaker than the saturation intensity, $\tilde{I}_0 \ll 1$, the atoms' velocities are hardly changed, and equation (5) reduces to

$$\frac{d\tilde{I}(z)}{dz} = -\rho\sigma_0\tilde{I}(z), \text{ from which we recover the analytic form,}$$
(A.3)
$$\sigma_0 n^{(0)} = -\ln\tilde{I}_0/\tilde{I}_f.$$
(A.4)

In the limit where the probe intensity is much larger than the saturation intensity, $\tilde{I}_0 \gg \tilde{\delta}$, even far detuned atoms will scatter light at their maximum rate. The time dependence of the detuning can thus be neglected, and equation (5) becomes

$$\frac{\mathrm{d}I(z)}{\mathrm{d}z} = -\rho\sigma_0, \text{ which integrates to}$$
(A.5)

$$\sigma_0 n = \tilde{I}_0 - \tilde{I}_f. \tag{A.6}$$

We recognize the right-hand sides of equation (A.4) and equation (A.6) as the two terms in equation (4). Thus, as shown in figure 3, OD^{sim1} coincides with the optical depth as predicted by equation (4) in both the small and large probe intensity limits.

We used the results of this simulation to check the self-consistency of the stationary atom assumption, i.e. the distance traveled by the atoms (as deduced from integrating the acquired recoil velocity over the imaging time) is less than the bin size. As shown in figure A1(b), not only do the atoms travel more than the bin size, but they travel far beyond the initial extent of the cloud. Moreover, owing to the higher initial scatter rate, the back of the cloud overtakes the front for long imaging times. Thus, the atomic distribution as a function of position changes dramatically during the imaging pulse, and the stationary assumption is invalid.

A.3. Traveling atom model

To account for the changing atomic distribution during the imaging pulse, we numerically simulated the classical kinetics of atoms subject to the recoil driven optical forces. To simulate large ensembles in a reasonable time, we modeled composite atoms, each describing the aggregate behavior of N_{ca} atoms. The amended algorithm is shown in algorithm 2.

Algorithm 2. Travelling atom model

 $z[n] = z_0, \tilde{\delta}[n] = 0$ { initialize position and detuning for each composite atom, labeled by index n } $O[i] = n \{ \text{make a list of composite atom indexes, ordered by position} \}$ $\tilde{I}[n = 0, t] = \tilde{I}_0 \{t \text{ is the time index}\}$ $H_f = 0$ {Radiant fluence seen by camera after passing through cloud} for t = 0 to t_f do {loop over time steps} for i = 1 to N_{ca} do {loop over composite atoms} n = O[i] {apply probe intensity to composite atoms in order of appearance} $A = \sigma_0 N_{ca} dz \{ dz \text{ is length over which atoms were grouped into single composite atom} \}$ $B = v_{\rm r} dt / (\hbar c N_{\rm sa}) \{ dt \text{ is the time step} \}$ $\tilde{I}[n, t] = \tilde{I}[n-1, t] - A\tilde{I}[n-1, t]/(1 + 4\tilde{\delta}[n]^2 + \tilde{I}[n-1, t]) \{\text{equation}(5)\}$ $\tilde{\delta}[n] + = B(\tilde{I}[n-1, t] - \tilde{I}[n, t]) \{\text{equation}(6)\}$ $z[n] + = dt\Gamma \tilde{\delta}/2k_r \{\Gamma \tilde{\delta}/2k_r \text{ is the velocity at } \tilde{\delta} \text{ detuning} \}$ end for O[i] = sort(n, key = z[n]) {sort composite atom indexes by current position} $H_f H_f + \tilde{I}[N, t] dt$ {collecting total fluence seen by the camera} end for $\mathrm{OD}^{\mathrm{sim2}} = -\mathrm{ln}(H_f/\tilde{I}_0 t_f)$

To validate our code, we again checked the velocity predicted in this model against known limits. One such limit is that of a single composite atom. In this case, there is no attenuation, and the intensity seen by the composite atom is constant at \tilde{I}_0 . Only the detuning evolves in time, and equations (5) and (6) give

$$\frac{\mathrm{d}\tilde{\delta}(t)}{\mathrm{d}t} = \frac{k_{\mathrm{r}}v_{\mathrm{r}}}{2}\frac{\tilde{I}}{1+4\tilde{\delta}^2+\tilde{I}}.$$
(A.7)

Equation (A.7) can be solved numerically, and is in agreement with our simulation, as seen in figure A2(a).

We used this model to study the time evolution of the cloud shape during imaging and visualized the phase space evolution of superatoms, shown in figure A3. The cloud shape is strongly distorted during imaging.

We compared the optical depths predicted by each of the two models, OD^{sim1} and OD^{sim2} . As seen figure A2(b), the predicted optical depths were hardly changed by including the full time evolution: $|OD^{sim1} - OD^{sim2}|/OD^{sim1} \leq 0.01$ for times up to 100 μ s, \tilde{I}_0 up to 50 and $\sigma_0 n$ up to 2.0.

Appendix B

We used a Zeeman slower to slow both ⁸⁷Rb and ⁴⁰K before capturing in a magneto-optical trap (MOT). After 7 seconds of MOT loading ⁴⁰K followed by 1.5 s of loading both ⁴⁰K and ⁸⁷Rb, we cooled both species in optical molasses for 2 ms. We optically pumped both species into their maximally stretched magnetically trappable states, |F = 9/2, $m_F = 9/2$ for ⁴⁰K and |F = 2, $m_F = 2$ for ⁸⁷Rb. Both species were then loaded into a quadrupole magnetic trap with a gradient of \approx 7.68 mT cm⁻¹ along e_z , and cooled evaporatively via forced rf evaporation, sweeping the rf frequency from 18 to 2 MHz in 10 s. The magnetic trap was plugged by a $\lambda = 532$ nm beam, tightly focused to \approx 30 μ m and \approx 5 W in power, providing a repulsive potential around the



Figure A1. (a) Column densities deduced from simulated optical depths of on-resonant imaging at probe intensity $\tilde{I}_0 = 0.8$. The input column density was $\sigma_0 n = 1.6$. $\sigma_0 n^1$ is the high probe intensity corrected column density given by equation (4). $\sigma_0 n^2$ is the column density as expanded to second order in time, equation (A.1). (b) Position of atoms as a function of imaging time for atoms in the first (solid green), middle (dashed red), and last (dotted blue) bins of the simulated density distribution for an initial cloud 50 μ m in extent. The probe intensity used in this calculation was 1.2 I_{sat} , and the column density was $\sigma_0 n = 1.6$.



Figure A2. (a) The velocity of a single composite atom as a function of probe intensity for various imaging times. Simulation data (dots) and numerical solutions of equation (A.7) (lines) are in agreement. (b) Top. Optical depth as a function of probe intensity for an imaging time $t = 100 \,\mu$ s. OD¹ and OD² are optical depths predicted from a given column density by equation (4) and (A.1) respectively. The two versions of simulated optical depth, OD^{sim1} (green curve) and OD^{sim2} (green dots) are plotted. Bottom. The fractional difference between two versions of the simulated OD, $|OD^{sim1} - OD^{sim2}|/OD^{sim1}$.

zero field point to prevent Majorana losses. Since the ⁴⁰K atoms were spin polarized and therefore only interacted by the strongly suppressed *p*-wave interactions, they re-thermalized largely due to sympathetic cooling with ⁸⁷Rb atoms.

We then loaded the atoms into a crossed optical dipole trap, provided by a 1064nm fiber laser, and continued evaporative cooling by slowly ramping down the dipole trap to trap frequencies of $(\omega_x, \omega_y, \omega_z)/2\pi$ = (39, 42, 124) Hz (for potassium atoms) in the three spatial directions, while also turning off the quadrupole field. We then used ARP to transfer the ⁸⁷Rb atoms from the |F = 2, $m_F = 2\rangle$ state to the |F = 1, $m_F = +1\rangle$ absolute ground state via 6.8556 GHz microwave coupling (20.02 MHz from the zero field resonance) followed by a magnetic field sweep from -0.469 to -0.486 mT in 50 ms. This state was chosen to minimize spin changing collisions with ⁴⁰K atoms during any further evaporation [37]. We then briefly applied an on-resonant probe laser, ejecting any remaining ⁸⁷Rb atoms in the F = 2 manifold from the trap. We again used ARP to transfer the ⁴⁰K atoms into the |F = 9/2, $m_F = -9/2\rangle$ state by using a 3.3 MHz rf field and sweeping the bias magnetic field from -0.518 to -0.601 mT in 150 ms.





References

- [1] Greiner M, Regal C A and Jin D S 2003 Emergence of a molecular Bose–Einstein condensate from a Fermi gas Nature 426 537–40
- [2] Zwierlein M W, Stan C A, Schunck C H, Raupach S M F, Gupta S, Hadzibabic Z and Ketterle W 2003 Observation of Bose–Einstein condensation of molecules *Phys. Rev. Lett.* 91 250401
- [3] Jochim S, Bartenstein M, Altmeyer A, Hendl G, Riedl S, Chin C, Hecker Denschlag J and Grimm R 2003 Bose–Einstein condensation of molecules Science 302 2101–3
- [4] Bartenstein M, Altmeyer A, Riedl S, Jochim S, Chin C, Hecker Denschlag J and Grimm R 2004 Collective excitations of a degenerate gas at the BEC-BCS crossover Phys. Rev. Lett. 92 203201
- [5] Bourdel T, Khaykovich L, Cubizolles J, Zhang J, Chevy F, Teichmann M, Tarruell L, Kokkelmans S J J M F and Salomon C 2004 Experimental study of the BEC-BCS crossover region in lithium 6 Phys. Rev. Lett. 93 050401
- [6] Zwierlein M W, Stan C A, Schunck C H, Raupach S M F, Kerman A J and Ketterle W 2004 Condensation of pairs of fermionic atoms near a Feshbach resonance Phys. Rev. Lett. 92 120403
- [7] Regal CA, Greiner M and Jin D S 2004 Observation of resonance condensation of fermionic atom pairs Phys. Rev. Lett. 92
- [8] Chin C, Grimm R, Julienne P and Tiesinga E 2010 Feshbach resonances in ultracold gases Rev. Mod. Phys. 82 1225-86
- [9] Timmermans E, Tommasini P, Hussein M S and Kerman A 1999 Feshbach resonances in atomic Bose–Einstein condensates Phys. Rep.-Rev. Section Phys. Lett. 315 199–230
- [10] Tiesinga E, Verhaar B J and Stoof H T C 1993 Threshold and resonance phenomena in ultracold ground-state collisions Phys. Rev. A 47 4114–22
- [11] Lysebo M and Veseth L 2009 Ab initio calculation of Feshbach resonances in cold atomic collisions: s- and p-wave Feshbach resonances in ⁶Li₂ Phys. Rev. A 79 062704
- [12] Gao B 2011 Analytic description of atomic interaction at ultracold temperatures: II. Scattering around a magnetic Feshbach resonance Phys. Rev. A 84 022706
- [13] Inouye S, Andrews M R, Stenger J, Miesner H J, Stamper-Kurn D M and Ketterle W 1998 Observation of Feshbach resonances in a Bose–Einstein condensate Nature 392 151–4
- [14] Cornish S L, Claussen N R, Roberts J L, Cornell E A and Wieman C E 2000 Stable Rb-85 Bose–Einstein condensates with widely tunable interactions Phys. Rev. Lett. 85 1795–8
- [15] Regal C A and Jin D S 2003 Measurement of positive and negative scattering lengths in a Fermi gas of atoms Phys. Rev. Lett. 90 230404
- [16] O'Hara K M, Hemmer S L, Gehm M E, Granade S R and Thomas J E 2002 Observation of a strongly interacting degenerate Fermi gas of atoms Science 298 2179–82
- [17] Monroe C, Cornell E, Sackett C, Myatt C and Wieman C 1993 Measurement of Cs–Cs elastic scattering at $T = 30 \ \mu$ K Phys. Rev. Lett. 70 414–7
- [18] Bartenstein M et al 2005 Precise determination of ⁶Li cold collision parameters by radio-frequency spectroscopy on weakly bound molecules Phys. Rev. Lett. 94 103201
- [19] Zürn G, Lompe T, Wenz A N, Jochim S, Julienne P S and Hutson J M 2013 Precise characterization of ⁶Li Feshbach resonances using trap-sideband-resolved RF spectroscopy of weakly bound molecules *Phys. Rev. Lett.* 110 135301
- [20] Chikkatur A P, Görlitz A, Stamper-Kurn D M, Inouye S, Gupta S and Ketterle W 2000 Suppression and enhancement of impurity scattering in a Bose–Einstein condensate Phys. Rev. Lett. 85 483–6
- [21] Thomas N R, Kjærgaard N, Julienne P S and Wilson A C 2004 Imaging of s and d partial-Wave interference in quantum scattering of identical Bosonic atoms Phys. Rev. Lett. 93 173201
- [22] Williams R A, LeBlanc L J, Jiménez-García K, Beeler M C, Perry A R, Phillips W D and Spielman I B 2012 Synthetic partial waves in ultracold atomic collisions *Science* 335 314–7

- [23] Aikawa K, Frisch A, Mark M, Baier S, Grimm R and Ferlaino F 2014 Reaching Fermi degeneracy via universal dipolar scattering Phys. Rev. Lett. 112 010404
- [24] Lu M, Burdick N Q and Lev B L 2012 Quantum degenerate dipolar Fermi gas Phys. Rev. Lett. 108 215301
- [25] Metcalf H J and van der Straten P 1999 Laser Cooling and Trapping (Graduate Texts in Contemporary Physics vol 26) (New York: Springer)
- [26] Reinaudi G, Lahaye T, Wang Z and Guéry-Odelin D 2007 Strong saturation absorption imaging of dense clouds of ultracold atoms Opt. Lett. 32 3143–5
- [27] Konstantinidis G O, Pappa M, Wikstroem G, Condylis P C, Sahagun D, Baker M, Morizot O and von Klitzing W 2012 Atom number calibration in absorption imaging at very small atom numbers Cent. Eur. J. Phys. 10 1054–8
- [28] Tiecke T G 2009 Feshbach resonances in ultracold mixtures of the fermionic quantum gases ⁶Li and ⁴⁰K PhD Thesis University of Amsterdam
- [29] Williams R A, Beeler M C, LeBlanc L J, Jiménez-García K and Spielman I B 2013 Raman-induced interactions in a single-component Fermi gas near an s-wave Feshbach resonance *Phys. Rev. Lett.* **111** 095301
- [30] Lin Y-J, Perry A R, Compton R L, Spielman I B and Porto J V 2009 Rapid production of ⁸⁷Rb Bose–Einstein condensates in a combined magnetic and optical potential *Phys. Rev.* A 79 063631
- [31] Jimenez-Garcia K 2012 Artificial gauge fields for ultracold neutral atoms *PhD Thesis* Joint Quantum Institute, National Institute of Standards and Technology, and the University of Maryland
- [32] DeMarco B, Bohn J L, Burke J P, Holland M and Jin D S 1999 Measurement of p-wave threshold law using evaporatively cooled fermionic atoms Phys. Rev. Lett. 82 4208–11
- [33] Wu S, Wang Y-J, Diot Q and Prentiss M 2005 Splitting matter waves using an optimized standing-wave light-pulse sequence *Phys. Rev.* A 71 043602
- [34] Edwards M, Benton B, Heward J and Clark C W 2010 Momentum-space engineering of gaseous Bose–Einstein condensates Phys. Rev. A 82 063613
- [35] Bracewell R 1965 The Fourier Transform and its Applications (New York: McGraw-Hill)
- [36] LeBlanc LJ 2011 Exploring many-body physics with ultracold atoms PhD Thesis University of Toronto
- [37] Best T 2010 Interacting Bose–Fermi mixtures in optical lattices PhD Thesis Johannes Gutenberg-Universitat