The C-terminus and Third Cytoplasmic Loop Cooperatively Activate Mouse Melanopsin
 Phototransduction

4	Juan C. Valdez-Lopez ¹ , Stephen T. Petr ^{1,2} , Matthew P. Donohue ^{3,4,5} , Robin J. Bailey ¹ , Meheret
5	Gebreeziabher ^{1,6} , Evan G. Cameron ^{1,7} , Julia B. Wolf ¹ , Veronika A. Szalai ² , Phyllis R. Robinson ^{1,*}
6	
7	¹ Department of Biological Sciences, University of Maryland, Baltimore County, Baltimore, MD, 21250;
8	² Current address: Eastern Virginia Medical School, Norfolk, VA, 23507; ³ Center for Nanoscale and
9	Technology, National Institutes of Standards and Technology, Gaithersburg, MD, 20899; ⁴ Maryland
10	NanoCenter, University of Maryland College Park, College Park, MD, 20742; ⁵ Current address: Salubris
11	Biotherapeutics, Gaithersburg, MD, 20878; ⁶ Current address: National Human Genome Research
12	Institute, National Institutes of Health, Bethesda, MD, 20892; ⁷ Current address: Department of
13	Ophthalmology, Byers Eye Institute, Mary M. and Sash A. Spencer Center for Vision Research, Stanford
14	University School of Medicine, Palo Alto, CA, 94034
15	
16	*To whom correspondence should be addressed: Prof. Phyllis R. Robinson, Department of Biological
17	Sciences, University of Maryland, Baltimore County, Baltimore, MD, 21250, USA. Email:
18	probinso@umbc.edu; Telephone: 410-455-2977
19	
20	Running Title: Melanopsin Structure and Activation
21	
22	
23	Keywords: G Protein-Coupled Receptor, Opsin, Visual Pigment, Phototransduction
24	
25	
26	

27 ABSTRACT

28 Melanopsin, an atypical vertebrate visual pigment, mediates non-image forming light responses 29 including circadian photoentrainment and pupillary light reflexes, and contrast detection for image 30 formation. Melanopsin-expressing intrinsically photosensitive retinal ganglion cells are characterized by 31 sluggish activation and deactivation of their light responses. The molecular determinants of mouse melanopsin's deactivation have been characterized (i.e. C-terminal phosphorylation and β -arrestin 32 33 binding), but a detailed analysis of melanopsin's activation is lacking. We propose that an extended 3rd cytoplasmic loop is adjacent to the proximal C-terminal region of mouse melanopsin in the inactive 34 conformation, which is stabilized by ionic interaction of these two regions. This model is supported by 35 site-directed spin labeling and electron paramagnetic resonance spectroscopy of melanopsin, the results of 36 which suggests a high degree of steric freedom at the 3rd cytoplasmic loop, which is increased upon C-37 38 terminus truncation, supporting the idea that these two regions are close in 3-dimensional space in wild-39 type melanopsin. To test for a functionally critical C-terminal conformation, calcium imaging of 40 melanopsin mutants including a proximal C-terminus truncation (at residue 365) and proline mutation of 41 this proximal region (H377P, L380P, Y382P) delayed melanopsin's activation rate. Mutation of all 42 potential phosphorylation sites, including a highly conserved tyrosine residue (Y382), into alanines also delayed the activation rate. A comparison of mouse melanopsin with armadillo melanopsin-which has 43 44 substitutions of various potential phosphorylation sites and a substitution of the conserved tyrosine— 45 indicates that substitution of these potential phosphorylation sites and the tyrosine residue result in dramatically slower activation kinetics, a finding that also supports the role of phosphorylation in 46 47 signaling activation. We therefore propose that melanopsin's C-terminus is proximal to intracellular loop 3 and C-terminal phosphorylation permits the ionic interaction between these two regions, thus forming a 48 stable structural conformation that is critical for initiating G protein signaling. 49

50

51 STATEMENT OF SIGNIFICANCE

52 Melanopsin is an important visual pigment in the mammalian retina that mediates non-image forming responses such as circadian photoentrainment and pupil constriction, and supports contrast 53 54 detection for image formation. In this study, we detail two critical structural features of mouse 55 melanopsin—its 3rd cytoplasmic loop and C-terminus—that are important in the activation of 56 melanopsin's light responses. Furthermore, we propose that these two regions directly participate in coupling mouse melanopsin to its G-protein. These findings contribute to further understanding of GPCR-57 G-protein coupling, and given recent findings suggesting flexibility of melanopsin signal transduction in 58 59 the retina (possibly by coupling to more than one G-protein type), these findings provide insight into the 60 molecular basis of melanopsin function in the retina.

61

62 INTRODUCTION

63 G-protein-coupled receptors (GPCRs) make up the largest family of integral membrane receptors 64 and are activated by a variety of biological stimuli, including hormones, odorants, small peptides, neurotransmitters and photons (1, 2). Upon activation, GPCRs undergo a series of conformational 65 66 changes that facilitate the binding and activation of intracellular heterotrimeric G-proteins, which initiate 67 a variety of signaling responses (3, 4). Visual pigments are specialized light-detecting GPCRs that are 68 comprised of an opsin protein covalently-attached to a chromophore, typically 11-cis-retinal in the 69 mammalian retina. The absorption of a photon by 11-cis-retinal results in its isomerization to all-trans-70 retinal. This conformational change in the chromophore results in the activation of visual pigment and the phototransduction cascade. In the mammalian retina, three distinct classes of photoreceptor cells detect 71 72 light. Rod and cone photoreceptor cells found in the outer retina express distinct specialized visual pigments that mediate photon absorption for image-forming vision. A less well-known third 73 photoreceptor class in the inner retina is composed of a small subset of intrinsically photosensitive 74 75 ganglion cells (ipRGCs) that express melanopsin, a rhabdomeric-type opsin (5, 6). In the mouse, ipRGCs 76 are divided into six subtypes (M1-M6) that are unique in their morphology, projections into the inner plexiform layer of the retina, amount of melanopsin expressed, transcription factors, and their projections 77

78	to the brain and thus, their functions (7, 8). For example, M1-ipRGCs exhibit the highest level of
79	melanopsin expression and project to the suprachiasmic nucleus (among other brain nuclei) and thus are
80	implicated in circadian photoentrainment (9) whereas M4-ipRGCs possess the largest soma of all ipRGC
81	subtypes, and project to the dorsal lateral geniculate nucleus and are implicated in image formation (10).
82	Light detected by melanopsin regulates non-image forming functions such as circadian photoentrainment,
83	pupillary light reflex, sleep, and melatonin synthesis (9, 11, 12, 13, 14). Unlike rods and cones,
84	melanopsin in M1-type ipRGCs signal through a G α q-mediated pathway that leads to the opening of
85	TRPC6/7 channels (15, 16, 17, 18). Recent studies analyzing the M4 ipRGC subtype suggest that
86	melanopsin in these cells signals either through the Gaq transduction cascade (19) or through a cyclic-
87	nucleotide cascade and the opening of HCN-channels (20).
88	Structural and molecular determinants of GPCR-G-protein complex formation have been
89	described in both a receptor specific (for review of rhodopsin activation see 21) and systematic manner
90	(22). However, it remains difficult to attribute specific receptor properties (e.g. amino acid sequence,
91	polarity, charge, steric properties) to their selectivity for their cognate G-protein. The notion of a singular
92	cognate G-protein for each receptor is also one that is challenged by evidence of receptor promiscuity to
93	several Ga classes (23, 24) including melanopsin promiscuously activating transducin <i>in vitro</i> (25).
94	GPCR regions important for G-protein binding selectivity include the 2 nd and 3 rd intracellular loops, and
95	cytoplasmic extensions of transmembrane helices 5 and 6, which form critical contacts with helices 4 and
96	5 on Ga (26). GPCR C-termini can regulate G-protein activation as well as serve as substrates for GPCR
97	kinase to mediate arrestin binding (for melanopsin C-terminal phosphorylation and deactivation, see 27,
98	28, 29, 30, 31, 32, 33). Structural analysis of rhodopsin—G-protein complexes reveals contacts between
99	helix 8 on rhodopsin's C-terminus and the C-terminal helix 5 on Gai (34) or transducin (35), and contacts
100	have been found on rhodopsin's C-terminus with $G\beta$ when in complex with $G\alpha$ i (36). Additionally,
101	rhodopsin C-terminal peptides bind and facilitate transducin signaling by enhancing phosphodiesterase
102	activity and inhibiting transducin's GTPase activity (37). The role of melanopsin's cytoplasmic loops or
103	C-terminus in stimulating G-protein activity remain to be determined.

104	In this study, we aimed to determine the critical cytoplasmic domains on mouse melanopsin that
105	contribute to phototransduction activation. Melanopsin-mediated behaviors are diverse and are both
106	image- and non-image-forming. Additionally, ipRGC light responses are sluggish (single flash responses
107	can persist across minutes) (38) compared to rapid photoresponses observed in rod and cone
108	photoreceptors (these cells can respond in a precise manner to millisecond-scale light flashes).
109	Melanopsin $G\alpha q$ signal transduction has been shown to be re-purposed to modulate leak potassium
110	channels (rather than activating phospholipase C) or could possibly couple to another G-protein to
111	stimulate cyclic nucleotide signal transduction, in M4-type ipRGCs (19, 20). Within the M1-type
112	population, there is a striking heterogeneity in the light responses, where there is a mixture of cells that
113	optimally respond to certain irradiances or respond in a linear fashion in response to increasing irradiance,
114	thus allowing for this M1 population to encode for a breadth of light intensities (39). Additionally,
115	melanopsin phosphorylation by Protein Kinase A, likely on the intracellular loops, can attenuate light
116	responses in a dopamine and cyclic AMP-dependent manner (28). It is therefore critical to examine
117	melanopsin—G-protein complex formation at a structural level to shed light on the molecular
118	determinants underlying the functional diversity observed in melanopsin signal transduction. Based on
119	amino acid sequence analysis and homology modeling using Todarodes pacificus rhodopsin as a
120	template, melanopsin is predicted to have extended 5 th and 6 th transmembrane helices and also a uniquely
121	long C-terminus (Figure 1 for 2D schematic of amino acids, Figure 2 for 3D figures), which is 171 amino
122	acids long (residues T350-L521). Given the functional significance of these regions in other GPCRs, we
123	synthesized a series of mutants designed to test if these regions contribute to phototransduction activation.
124	Our findings provide a detailed examination of the structural basis of melanopsin activation and,
125	furthermore, provide a more robust foundation to examine ipRGC activation and structure-function
126	relationships of other related visual pigments.
127	

128 EXPERIMENTAL PROCEDURES

129	Note that certain commercial equipment, instruments, and materials are identified in this paper to
130	specify an experimental procedure as completely as possible. In no case does the identification of
131	particular equipment or materials imply a recommendation or endorsement by NIST, nor does it imply
132	that the materials, instruments, or equipment are necessarily the best available for the purpose. The
133	opinions expressed in this article are the authors' own and do not necessarily represent the views of NIST.
134	Homology modeling of mouse melanopsin-Mouse melanopsin amino acid sequence (Uniprot ID:
135	Q9QXZ9-1; OPN4L) was used to produce a homology model using the online protein structure service
136	LOMETS, which ranks models derived from 11 servers using a Z-score as a metric of confidence for each
137	model (40). The melanopsin structural model that was generated used Todarodes pacificus rhodopsin
138	(PDB ID: 2ZIY) (48) as the template, which was the highest ranked model and was produced using the
139	HHpred threading program (41). To model the active conformation of mouse melanopsin, the LOMETS-
140	generated homology model was aligned to the crystal structure of the β_2 -adrenergic receptor in complex
141	with Gas (PDB ID: 3SN6) (46) and cryo-EM structure of rhodopsin in complex with Gai (PDB ID:
142	6CMO) (34) in PyMol (Schrödinger Inc, New York, NY) using the cealign tool, and predicted
143	melanopsin helical movements were modeled using the positions of the transmembrane helices of the β_2 -
144	adrenergic receptor or rhodopsin structures. Surface electrostatic potential of melanopsin was calculated
145	using vacuum electrostatics function on PyMol.
146	Synthesis of Melanopsin Mutants-All mutant genes were constructed using the mouse
147	melanopsin coding sequence (NCBI accession: NM_013887.2) cloned in the mammalian expression
148	plasmid PMT3 (42). Melanopsin Δ 365, H377A L380A Y382A, phosphonull + Y382S, melanopsin ICL3
149	Null, melanopsin 280-285 alanine, armadillo melanopsin constructs, and Mouse 1D4-Gaq (Mus musculus
150	Gnaq with N-terminal 1D4 tag—amino acid sequence TETSQVAPA, corresponding to the last 9 amino

acids of bovine rhodopsin, NCBI accession: NM_008139.5) were synthesized by cassette synthesis using

synthetically-synthesized gene fragments. Melanopsin C268 was constructed using Gibson assembly (43)

- 153 of synthetically-synthesized gene fragments to generate mutagenesis of residues C95, C271, C438, C469,
- 154 C493, C499 to alanine residues to eliminate non-specific spin label attachment. These constructs for

electron paramagnetic resonance (EPR) spectroscopy were also C-terminally tagged with the 1D4 epitope
to facilitate immunoprecipitation when expressed heterologously. Melanopsin C268 Δ365 was
synthesized through cassette synthesis using melanopsin C268 as a PCR template and with a reverse
primer to amplify from the start of the gene to the end of the codon encoding for L365. Melanopsin
H377P L380P Y382P, phosphonull + Y382A, and phosphonull + Y382F were generated using
Quickchange-based site-directed mutagenesis (44). All plasmids were sequence verified through Sanger
sequencing.

162 Cell Culture and Transfection of HEK293 Cells—HEK293 cells were grown in a monolayer on 163 10 cm culture dishes in Dulbecco's Modified Eagle Medium (DMEM) supplemented with fetal bovine serum (to a final total protein concentration of 3.6 g/L), 0.006 g/L penicillin, 0.01 g/L streptomycin, and 164 2.5 x 10⁻⁵ g/L Amphotericin B in a humid CO₂ incubator at 37 °C. Cells were passaged by disassociation 165 166 using 2.5 g/L trypsin-EDTA and seeded to a density of up to 5 000 000 cells per 10 cm dish and 240 000 167 cells per well in a 6 well dish. Transfections were done using Turbofect Transfection Reagent (ThermoFisher Scientific, Waltham, MA) as per manufacturer's protocol. Briefly, 4 or 10 µg DNA (6-168 169 well and 10 cm dish, respectively) was diluted in non-supplemented DMEM, and then the transfection 170 reagent was added to the mixture. Following a 20 min incubation at room temperature, the transfection 171 mixture was added to the cells, and the cells were incubated in a humid CO₂ incubator for 24 h to 48 h. 172 Calcium Imaging of Melanopsin-Transfected HEK293 Cells—24 h after transfection, HEK293 cells were trypsin-disassociated from the plates and seeded to 96-well plates at a density of 100 000 cells 173 174 per well. The 96-well plate is incubated overnight in a humid CO₂ incubator in a dark room for dark 175 adaptation. The cells are then incubated with equivalent volume of Fluo-4 AM Direct Calcium Dye supplemented with 20 mmol/L probenecid and 20 µmol/L 9-cis-retinal and are incubated for 1 hr in a 176 humid CO₂ incubator. Calcium kinetics were then measured on a commercial fluorescence plate reader by 177 178 exciting the sample at 487 nm and recording the emission fluorescence at 516 nm at a rate of 1 Hz. The 179 plate reader monochromator has a bandwidth of <9 nm for excitation light. Every melanopsin construct in 180 each transfection had six replicates.

181	Protein expression and preparation for EPR spectroscopy—Melanopsin C268 and C268 Δ 365
182	constructs were transfected in HEK293 cells in 60 to 80 10 cm plate batches, harvested 48 hr later, and
183	stored at -80 °C. Frozen cell pellets were then thawed on ice then immediately resuspended in PBS with
184	40 µmol/L 9-cis-retinal and incubated for at least an hour at 4 °C to reconstitute the visual pigment. The
185	sample was centrifuged at max speed on a clinical centrifuge and the cell pellet was then resuspended in
186	solubilization buffer composed of 1 g/L n-dodecyl-β-D-maltoside, 3 mmol/L MgCl2, 140 mmol/L NaCl,
187	1 mmol/L phenylmethylsulfonyl fluoride (PMSF), and 50 mmol/L HEPES, pH 7.5. After a minimum of 1
188	hr on a tube rotator at 4 °C, the solubilized cell lysate was centrifuged at max speed on a clinical
189	centrifuge and the supernatant was collected. The solubilized protein was then incubated with Sepharose-
190	4B resin conjugated with monoclonal α -1D4 antibody (provided generously by Prof. Daniel Oprian) for at
191	least an hour at 4 $^{\circ}$ C. The 1D4-resin was then washed by centrifuging 30 secs at 500 x g _n , removing the
192	supernatant, and resuspending the resin with solubilization buffer at least 10 times. After the final wash,
193	the 1D4-resin was resuspended in solubilization buffer containing 2 mmol/L MTSL to attach the spin
194	label at C268 for EPR spectroscopy, and was incubated overnight on a tube rotator at 4 °C. The sample
195	was then washed by centrifuging 30 secs at 500 x g_n , removing the supernatant, and resuspending the
196	resin with solubilization buffer at least 5 times. MTSL spin-labeled melanopsin was then eluted by
197	incubating the 1D4-resin in solubilization buffer containing 50 µmol/L 1D4 peptide (amino acid sequence
198	TETSQVAPA). Eluate was centrifuged using 10 000 g/mol (10 kDa) size-exclusion centrifugal filters to
199	concentrate the sample and filter out the eluting peptide.
200	EPR Spectroscopy of single spin-labeled melanopsin— CW EPR spectra at X-band
201	(approximately 9.8 GHz) were recorded using a commercial EPR spectrometer equipped with a
202	commercial microwave cavity in perpendicular mode. Samples were drawn, by capillary action, into

- and subsequently inserted into a critically coupled EPR cavity at ambient temperature. Spectra (88
- averages) were recorded using 0.1 mT modulation amplitude at 100 kHz, with an incident microwave
- 206 power of 1.5mW. Simulations of EPR data were done using EasySpin (45).

207 Calculation of Melanopsin Activation Rates—The activation phases of calcium imaging assay data (corresponding to the part of the data prior to the peak fluorescence level) for all melanopsin 208 209 constructs was fitted to a one-phase association function using GraphPad Prism software (GraphPad 210 Software, Inc., San Diego, CA). The following function was used: $Y=Y(0) + (Plateau-Y(0)) * (1-e^{(k*x)})$ 211 212 Where Y(0) is the value at t=0, Plateau is the peak activation value, and k is the rate constant. Data across 213 multiple transfections (six replicates per transfection) were pooled together and averaged, and the 214 activation rate of the averaged data was calculated. Standard error of the mean of the activation rates of 215 all melanopsin constructs and statistical analysis were calculated using GraphPad software. Statistical significance was determined by performing unpaired t-tests of all mutant melanopsin constructs with 216 respect to wild-type melanopsin, or between specific melanopsin constructs of interest. 217 218 219 RESULTS Three-dimensional homology modeling of mouse melanopsin predicts that an extended 3rd 220 221 cytoplasmic loop and the proximal C-terminus mediate G-protein binding—The structural basis of 222 melanopsin activation is not well understood; specifically, the unique molecular mechanisms underlying 223 its capability to couple its cognate G-protein have not been elucidated in detail. To shed light on this 224 problem, we constructed a homology model of mouse melanopsin using squid rhodopsin as a template. 225 The generated homology model had 68 % of melanopsin's amino acid sequence aligned to the template (Supplemental Figure 1), and among these amino acids, they shared 37 % sequence identity to the squid 226 227 (Todarodes pacificus) rhodopsin template. The sequence coverage of our model includes the entirety of the seven-transmembrane helical domain and the proximal portion of the C-terminus, up to S398. In the 228 229 inactive state (Figure 2A), the model predicts that melanopsin forms a seven-transmembrane visual pigment with an extended 3rd intracellular loop, which is comprised of cytoplasmic extensions of 230 231 transmembrane helices 5 and 6. In the active state, which we modeled based on helical movements of β_2 adrenergic receptor (46) and rhodopsin (34), melanopsin's 5th and 6th transmembrane helices are predicted 232

to swing away from the C-terminus (Figure 2A), thereby making way for the attachment of G-protein to
the newly formed binding pocket (Figure 2B-K). The generated homology model also predicts a Cterminal 9th helix (Figure 2A), but further experimental data is needed to test the existence of this
structure in melanopsin. However, the data presented in this study strongly support the predicted Cterminal conformation shown in the homology model.
The crystal structures obtained for squid rhodopsin reveal contacts between intracellular loop 3

239 (specifically, the cytoplasmic extension of transmembrane helix 6) and its C-terminal 9th helix (47, 48). Additionally, the presence of positively charged residues on squid rhodopsin's 3rd intracellular loop and 240 negatively charged residues on the C-terminus, may facilitate the formation of contacts between these 241 regions, or potentially to the Gaq C-terminus. Additionally, while most GPCRs display promiscuous 242 243 binding to several $G\alpha$ -subtypes, several molecular features on the GPCR determine the selectivity for 244 cognate G-proteins—the cognate G-protein being Gaq for melanopsin. Systematic analysis of GPCRs 245 suggest that selectivity to the Gag/11 family is determined primarily by the amino acid properties of the 246 receptor's C-terminus, specifically through enrichment of non-aromatic residues (22). Interestingly, neither the length of either the C-terminus nor the length of 3rd intracellular loop are positively correlated 247 248 with selectivity to Gag/11-subtypes and are only selectivity features in the binding of Ga12/13 (22). 249 Surface electrostatic analysis of our homology model of melanopsin suggests an enrichment of positively 250 charged residues on the intracellular loops, particularly intracellular loop 3 (Figure 2E & 2J) and 2 251 (Figure 2F & 2K). The proximal region of melanopsin's C-terminus is also positively charged (Figure 2E & 2J), and we predict it will make contacts with positively-charged residues on transmembrane helix 6 in 252 the inactive conformation (Figure 2A). Thus, the ionic contacts between the C-terminus and 3rd 253 254 intracellular loop would require amino acid modification (namely, C-terminal phosphorylation) to provide 255 the necessary charge for these ionic contacts. Comparison of both our active state melanopsin models 256 suggests that melanopsin would have to undergo different helical movements, specifically a greater outward swing of transmembrane helix 6 when melanopsin is modeled with Gas (Figure 2L & 2M). 257 Systematic analysis of GPCRs suggests no positive correlation with negative surface charge at these 258

259 critical cytoplasmic domains on the receptor and their capability to couple $G\alpha q$ (22). However, we propose that the amino acid ionic charges and the conformation of melanopsin's cytoplasmic domains 260 261 play critical roles in its ability to couple to G-protein(s) and initiate ipRGC phototransduction. Electron paramagnetic resonance (EPR) spectroscopy of single spin-labeled mouse melanopsin 262 263 supports the proposed C-terminus—intracellular loop 3 arrangement—To test our model that predicted 264 the C-terminus and cytoplasmic loop, we synthesized a site-directed spin labeling mutant melanopsin for 265 EPR spectroscopy analysis, Melanopsin C268. All solvent-accessible, non-specific cysteines were 266 mutated to alanine residues with the exception of C364, a predicted palmitoylation site, and residues 267 C142 and C220, which are predicted sites of a disulfide bridge, a common molecular feature of opsins necessary for proper folding and retention of covalently-linked retinal molecule (49, 50). This melanopsin 268 269 C268 mutant was designed for single labeling using the nitroxide spin label MTSL (2,2,5,5-tetramethyl-1-270 oxyl-3-methyl methanethiosulfonate) at C268 on intracellular loop 3 to analyze the steric freedom at this 271 position in the presence (Melanopsin C268) or absence (Melanopsin C268 Δ 365) of the C-terminus. The continuous wave (CW) EPR spectrum of single spin-labeled, dark-adapted melanopsin C268 272 273 (Figure 3A) in *n*-dodecyl- β -D-maltoside (DM) presents a sharper spectral lineshape compared to similar 274 EPR measurements on rhodopsin (51, 52), a finding that indicates a high degree of rotational mobility of 275 the MTSL spin label at this position. In addition to the qualitative observation of a sharper spectral 276 lineshape, mathematical fitting of the EPR spectrum of single spin-labeled melanopsin C268 to extract a 277 rotational correlation time (τ_{corr}) of the spin label supports the higher degree of steric freedom at this region compared to rhodopsin. The calculated τ_{corr} of melanopsin C268 was 9.36 ns (Supplemental Figure 278 279 2), which is lower than $\tau_{\rm corr}$ of both inactive and active forms of rhodopsin (77 ns and 26 ns, respectively) 280 when spin-labeled at the cytoplasmic end of transmembrane helix 6 (V250C) (53). One possible explanation for the increased spin label rotational mobility in the C268-labeled melanopsin is that the 3rd 281 282 intracellular loop is extended into the cytoplasm, as predicted by the structural model. The broader 283 lineshape and longer rotational correlation times for rhodopsin (51, 52, 54) suggest that the spin label's rotational mobility (MTSL attached at the cytoplasmic face of transmembrane helices) is hindered due to 284

closer proximity to the DM micelle compared to melanopsin's spin label. The CW EPR spectrum of single spin-labeled, dark-adapted C-terminal truncated melanopsin (Melanopsin C268 Δ 365) in DM (Figure 3B) possesses an even sharper spectral lineshape than full-length melanopsin and resembles the lineshape of the spectrum observed when measuring free MTSL in DM solution, without protein (Figure 3C). Together, these data indicate a higher degree of MTSL rotational mobility at the 3rd intracellular loop, which we attribute to the elimination of any steric hinderance caused by the C-terminus consistent with the C-terminal conformation proposed in Figure 2.

After measurement of the EPR spectra of full length and truncated Melanopsin C268, the samples were exposed to 30 s of 300 W white light to test for differences in the spectra upon light-induced conformational change of the visual pigment. Comparison of the dark and light EPR spectra indicates minimal differences between the two spectra of each sample (Figure 3A & 3B), indicating that the rotational mobility of the spin label at position C268 is minimally affected by light-induced conformational change.

298 The proximal region of mouse melanopsin's C-terminus is necessary for rapid phototransduction 299 activation—To test the hypothesis that melanopsin's C-terminus regulates signaling initiation, we 300 synthesized melanopsin C-terminal mutants (positions in Figure 1B) aimed to disrupt or ablate C-terminal 301 structure or configuration in its proximal region (prior to residue D396). To test for potential C-terminal 302 regulation of phototransduction activation, we synthesized a mouse melanopsin C-terminus truncation 303 mutant, Melanopsin $\Delta 365$, which is truncated at residue L365, and eliminates the C-terminal region predicted to form part of the G-protein binding pocket in our structural model. Calcium imaging of this 304 305 mutant reveals delayed activation kinetics (Figure 4A) compared to wildtype melanopsin, supporting a 306 functional role of this C-terminal region in mediating signaling activation. Previous studies that examined 307 melanopsin C-terminal truncations (29) suggest that the distal C-terminus (after D396) doesn't contribute 308 to the production of rapid signaling activation and can potentially sterically hinder the G-protein from 309 accessing the cytoplasmic side of the receptor (32). To disrupt the predicted proximal C-terminal conformation (C-terminus near cytoplasmic loop 3, depicted in Figure 2), we introduced proline 310

311	mutations (H377P L380P Y382P) in the critical region predicted to work synergistically with the
312	cytoplasmic loops to regulate phototransduction activation. These proline mutations introduced backbone
313	rigidity and thus will disrupt any important conformations or potential secondary structures found in the
314	wild-type protein (55). Calcium imaging of this triple proline mutant (Figure 4B) reveals that it has
315	delayed activation kinetics compared to wildtype melanopsin, indicating this region has a functionally
316	significant conformation, as suggested by the structural modeling. Mutation of those three residues
317	produces the most pronounced reduction in activation rate, as pairwise proline melanopsin mutants
318	H377P L380P and L380P Y382P show slight, but not statistically significant reductions in activation rate,
319	while the H377P Y382P and H377P L380P Y382P show significant activation rate reduction
320	(Supplemental Figure 3). To test that the activation defect observed in the proline mutant is attributable to
321	a proline-induced kink or disruption in structure, we synthesized and tested another mutant of melanopsin
322	with alanine mutations at the same sites of interest in the proline mutant (H377A L380A Y382A). We
323	predicted that these mutations would not disturb any functionally important C-terminal structure or
324	conformation, unlike the proline mutations. Calcium imaging of this mutant shows no reduction in the
325	activation rate, suggesting that the proline mutant's defect is indeed due to a disruption of C-terminal
326	structure or conformation.
327	C-terminal palmitoylation anchors and stabilizes GPCR C-termini (56) and in rhodopsin, it
328	regulates its proper expression and stability in the membrane (57, 58). Mutation of mouse melanopsin's
329	predicted palmitoylation site from a cysteine to serine residue and calcium imaging of this mutant,
330	melanopsin C364S, reveals a very modest increase in this mutant's time to reach peak activation, and a
331	slower deactivation compared to wild-type melanopsin (Supplemental Figure 4). These data suggest that

332 C364, if palmitoylated, might stabilize the proximal C-terminus for signal transduction, possibly in a

cooperative manner with the critical C-terminal conformation downstream of this site. Taken together, we

propose that the proximal region of the C-terminus, between residues L365 and D396, holds functional

335 significance that is dependent on an important structural conformation of these residues.

336	Proximal C-terminal phosphorylation regulates melanopsin phototransduction activation
337	kinetics-Relative to rod and cone opsins, mouse melanopsin has a uniquely long C-terminus that
338	possesses 38 serine and threonine residues (Figure 1B) that may serve as potential phosphorylation sites.
339	It is well established that melanopsin C-terminal phosphorylation is critical for signaling deactivation and
340	the lifetime of melanopsin-driven behaviors such as pupil constriction and jet-lag photoentrainment (28,
341	29, 31, 32). Given the enrichment of serine and threonine residues at the proximal region (from residues
342	H351-T385) of mouse melanopsin's C-terminus, we tested if phosphorylation of these sites contributes to
343	regulation of phototransduction activation, in addition to deactivation. To test this idea, we performed
344	calcium imaging of transiently-transfected HEK293 cells expressing melanopsin C-terminal
345	phosphorylation mutants: phosphonull melanopsin (28, 31), a mutant with all 38 C-terminal serine and
346	threonine residues mutated to alanine residues, and phosphonull + Y382A melanopsin, a mutant with an
347	evolutionarily conserved tyrosine residue (Figure 5A) mutated to an alanine residue in addition to the
348	phosphonull C-terminus mutations. Calcium imaging of these mutants (Figure 5B) suggests that
349	melanopsin C-terminal phosphorylation contributes to signaling activation, due to both mutants
350	displaying slower rates of activation (Figure 5C) compared to wildtype melanopsin. Phosphonull +
351	Y382A melanopsin displays a slower rate of activation than phosphonull melanopsin, similar to
352	melanopsin $\Delta 365$ and melanopsin H377P L380P Y382P mutants. This implies that Y382, an
353	evolutionarily conserved residue in mammalian melanopsins, holds key significance for regulating the
354	kinetics of signaling activation.

To elucidate the mechanism of this tyrosine residue, specifically, its potential phosphorylation to facilitate ionic interaction with intracellular loop 3, we synthesized additional point mutations at that position, specifically, phosphonull + Y382S and phosphonull + Y382F (Figure 5B). By mutating Y382 to serine and phenylalanine residues, we aimed to test if potential phosphorylation at this position (by means of the phosphorylatable residue, serine) contributes to activation kinetics. Specifically, if the elimination of phosphorylation, but retention of the atomic mass (phenylalanine has a similar aromatic structure as tyrosine but lacks the hydroxyl group for phosphorylation modification) produces a similar defect as

phosphonull + Y382A, then we can conclude that the addition of a phosphate group is important. Calcium imaging of these mutants reveals that phosphonull + Y382S has a higher rate of activation compared to phosphonull + Y382A, and has a similar rate as phosphonull and is not statistically lower than wild-type melanopsin's rate of activation (Figure 5C). This supports the importance of potential phosphorylation at this position. Analysis of the activation rate of phosphonull + Y382F shows that it has slow activation kinetics, slower than phosphonull + Y382A (Figure 5C), supporting the conclusion that charge and not atomic size is important in this region of melanopsin.

369 To further test the evolutionary basis of C-terminal phosphorylation and the proposed functional 370 tyrosine, Y382, in signaling activation, we transiently transfected armadillo melanopsin in HEK293 cells 371 and tested its signaling kinetics using calcium imaging. Armadillo melanopsin possesses a shorter C-372 terminus than mouse melanopsin, and it has fewer C-terminal serine and threonine residues than mouse 373 melanopsin (Armadillo melanopsin has 14 potential phosphorylation sites compared to 38 in mouse 374 melanopsin, Supplemental Figure 5). Additionally, armadillo melanopsin has a substitution of the 375 evolutionarily conserved tyrosine residue (P381 in armadillo melanopsin, Figure 5A), which may 376 potentially directly impact the activation kinetics. Calcium imaging of mouse and armadillo melanopsins 377 (Figure 5D) indicates a striking difference in signaling kinetics, most notably, armadillo melanopsin's 378 activation kinetics are significantly slower than mouse melanopsin (Figure 5E) and displays a brief time 379 period of inactivity before phototransduction onset. This suggests that the lack of C-terminal serine and 380 threonine residues, particularly the proximally-located (prior to residue D396) residues, and the lack of the tyrosine residue in armadillo melanopsin directly affects its capability to activate phototransduction. 381 382 Furthermore, point mutations of armadillo melanopsin to mimic mouse melanopsin residues (L371S P375S G378S P381Y A388T G391S A393S D412S E417T) resulted in faster activation kinetics (Figure 383 5D & 5E) than wild-type armadillo melanopsin, further supporting the importance of C-terminal 384 385 phosphorylation in producing rapid signaling activation.

Positively-charged residues on mouse melanopsin's 3rd intracellular loop play a critical role in
 G-protein activation—The functional importance of GPCR intracellular loop domains in the coupling and

activation of its cognate G-protein has been established in many receptors. We wanted to test what 388 389 molecular features of melanopsin's cytoplasmic loops confer its capability to activate G-protein signaling. We focused on the 3rd intracellular loop due to its extended nature and predicted interaction with the C-390 terminus. In addition, melanopsin's 3rd intracellular loop is rich with positively-charged residues, and we 391 392 wanted to test if this property has any role in phototransduction activation. First, we synthesized a melanopsin mutant where amino acids 280-285 (R280 O281 W282 O283 R284 L285) on the cytoplasmic 393 394 extension of transmembrane helix 6 were mutated to alanine residues (Melanopsin 280-285 Alanine, Figure 6A) to disrupt any hydrogen-bonding or ionic contacts with the C-terminus, as predicted by our 395 modeling. Calcium imaging of this mutant reveals sluggish activation kinetics (Figure 6D), similar to our 396 397 C-terminal activation mutants (Figure 4). This data, when analyzed with the structural model, suggests that this region of the 3rd cytoplasmic loop potentially forms functionally significant contacts between 398 399 intracellular loop 3 and the C-terminus. Furthermore, when this data is taken together with the data in 400 Figure 5, we propose that C-terminal phosphorylation could add the requisite negative charge to allow 401 ionic contacts to form with these positively-charged intracellular loop 3 residues. While we propose this 402 functionally important cytoplasmic conformation and a mechanism for its stabilization (i.e. intracellular 403 loop 3—C-terminus ionic interaction facilitated by C-terminal phosphorylation) further point-mutation analyses would be required to validate this interaction and to identify the required residues for this 404 405 interaction to occur.

To test for any functional significance of the positive charge localized at the 3rd cytoplasmic loop, we mutated all positively-charged residues on the 3rd intracellular loop (Figure 6B) to alanine residues (Melanopsin R259A R262A R266A R277A R279A R280A R284A K390A K293A or Melanopsin ICL3 Null, Figure 6C) to alter the loop's surface charge closer to neutral. Calcium imaging of this mutant shows a striking inability to activate phototransduction (Figure 6D), suggesting that the positive charge at the 3rd intracellular loop is critical for melanopsin to couple to G-protein in any capacity, thus we propose this feature as a contributor to melanopsin's capability to couple to its cognate G-protein.

413

414 **DISCUSSION**

415 The diversity of molecular features and amino acid properties in the cytoplasmic regions of GPCRs complicates the prediction of receptor-G-protein binding properties in an unstudied receptor. It is 416 417 evident in many class A GPCRs, including the prototypical receptor and visual pigment rhodopsin, that 418 cytoplasmic domains including the cytoplasmic loops and the C-terminus are critical for G-protein binding and selectivity. Melanopsin-expressing ipRGCs in the mammalian retina regulate a diversity of 419 420 non-visual and visual functions and functional heterogeneity is observed across ipRGC subtypes and even within the same subtypes. In this study, we propose our model of melanopsin consisting of a functionally 421 critical cytoplasmic conformation consisting of extended transmembrane helices 5 and 6 that form a 3rd 422 423 cytoplasmic loop domain that protrudes far into the cytoplasm. Additionally, in the dark inactive state, the 424 proximal region of the C-terminus (prior to residue D396, Figure 4A) takes on a conformation where, 425 after the predicted 8^{th} helix, it is adjacent to the cytoplasmic extension of transmembrane helix 6, and 426 close enough to form contacts. These contacts could potentially be ionic in nature, and our data support 427 the importance of C-terminal phosphorylation sites, and Y382-which may also be phosphorylated-in 428 phototransduction activation. Thus, C-terminal phosphorylation could provide the negative charge that is needed for the C-terminus to ionically interact with the arginine and lysine residue-rich 3rd cytoplasmic 429 loop, in the inactive state, prior to light-induced activation. Based on our homology modeling, we propose 430 that upon photon-mediated conformational change to active-state melanopsin, the 5th and 6th 431 432 transmembrane helices undergo an outward swing, moving the cytoplasmic ends of these helices—and the 3rd cytoplasmic loop—away from the C-terminus. We also propose that the C-terminus, together with 433 the 3rd cytoplasmic loop, form a critical part of the G-protein binding pocket and may play a role in G-434 435 protein specificity or stabilization of the melanopsin-G-protein complex. Previous studies have also generated homology models of melanopsin utilizing squid rhodopsin (59, 60) or bovine rhodopsin (61), 436 437 but these studies focused primarily on making computational predictions of melanopsin's chromophore 438 chemistry, and not focused on structural determinants of melanopsin—G-protein binding. However, these

- 439 previous models also generated a predicted structure containing the features we highlight here (i.e.
- 440

extended 5th and 6th transmembrane helices/3rd intracellular loop and C-terminal conformation).

Our EPR spectroscopy measurements on melanopsin support the cytoplasmic extension of the 3rd 441 442 cytoplasmic loop because the spectral lineshape that is indicative of high spin-label rotational mobility at 443 that region. Interestingly, while a clear difference was observed in full length and truncated melanopsin 444 C268 EPR spectra, we did not observe dramatic differences in the spectral lineshape after light exposure 445 in both samples. This result was unexpected, as we did predict to see light-induced conformational change reflected in the EPR spectra. Given the steric freedom at C268 and in the 3rd intracellular loop region, it is 446 possible that this region is unaffected by helical movement or conformational change due to its predicted 447 distance from the membrane and from other structures. However, our model and data—including our EPR 448 data of full length and C-terminal truncations—support the idea that the proximal C-terminus is adjacent 449 to the 3rd intracellular loop. Our model also predicts that upon light exposure, if melanopsin adopts a 450 451 similar active conformation as the β^2 adrenergic receptor, another family A GPCR, then we would expect 452 these regions to separate and form the G-protein binding pocket. This would lead to the prediction that the 453 EPR spectrum of the light-exposed full length melanopsin C268 sample should differ from the dark 454 spectrum. It is possible that this movement happens as predicted by our modeling but does not produce a significant difference in the EPR spectral lineshape. Alternatively, it is also possible that the predicted 455 interactions between these two regions, the C-terminus and the 3rd cytoplasmic loop, remain intact even 456 457 after light-induced activation. We observed no differences in the dark and light EPR spectra of the $\Delta 365$ 458 truncated melanopsin C268 because this mutant lacks this C-terminal region. We therefore propose that 459 our model of melanopsin activation is robust, but identification of structural and conformational changes 460 of melanopsin after light exposure remains an important question for further study.

461 Precise structural and molecular determinants of GPCR selectivity to its cognate G-protein(s) are 462 difficult to isolate, due the promiscuity of GPCRs for multiple G-proteins. While it would be erroneous to 463 suggest that, like melanopsin, all G α q-binding GPCRs require similar structural features (i.e. extended 5th 464 and 6th transmembrane helices/3rd cytoplasmic loop, positively charged cytoplasmic loops, and C-

465 terminus contacting G-protein) for their selectivity, we propose that similar features are shared amongst 466 rhabdomeric-type visual pigments. Of the available resolved protein structures of GPCRs, two are of 467 rhabdomeric visual pigments: two squid rhodopsin structures and jumping spider rhodopsin (47, 48, 62). Similar structural features are observed in melanopsin's predicted structure and the structures of the 468 resolved rhabdomeric-type visual pigments. Specifically, the extended 5th and 6th transmembrane helices 469 470 are a common feature amongst melanopsin and the invertebrate visual pigments, and like squid rhodopsin, 471 melanopsin also has an extended C-terminus, much longer than vertebrate, ciliary-type image-forming 472 visual pigments like rhodopsin. Melanopsin's intracellular loops are predicted to possess a more uniform positive charge compared to the surface charges observed on the loops of squid and jumping spider 473 rhodopsin, which are also positively-charged, but do possess regions of negative charge, particularly on 474 the 5th transmembrane portion of the 3rd cytoplasmic loop. It is evident that this positive charge hold 475 476 significance in melanopsin, but it remains unclear as to the extent of this feature's significance for $G\alpha q$ 477 interaction. Given the enrichment of potential phosphorylation sites on melanopsin's C-terminus and 478 intracellular loops, this provides an interesting mechanism to modify the electrostatic potential of these 479 regions, and thereby provides an interesting way to alter G-protein binding, and potentially specificity. 480 Thus, with these findings, we identified critical regions that regulate melanopsin's phototransduction 481 activation to shed light onto ipRGC function and provide targets for protein engineering for use as an 482 optogenetic tool or further study of its phototransduction.

483

484

485 AUTHOR CONTRIBUTIONS

JCVL, STP, MPD, MG, and RJB conducted experiments. JCVL, EGC, JBW, VAS, and PRR
intellectually contributed and conceived experiments. JCVL wrote the manuscript. PRR edited the
manuscript.

489

490 ACKNOWLEDGEMENTS

491	We thank and acknowledge NIH Training Grant T32 GM066706 awarded to J.C.V.L., generously
492	made possible by Prof. Katherine Seley-Radtke and NIH R01 EY027202-01A1 awarded to P.R.R. We
493	also thank Prof. Daniel D. Oprian for generously providing 1D4-antibody. M.P.D acknowledges support
494	under the Cooperative Research Agreement between the University of Maryland and the National
495	Institute of Standards and Technology Center for Nanoscale Science and Technology Award
496	70NANB10H193, through the University of Maryland. J.C.V.L also wishes to acknowledge and thank
497	Charles Thomas and Ramses Long for assistance in optimizing protein expression.
498	
499	CONFLICT OF INTEREST
500	The authors declare that they have no conflicts of interest with the contents of this article.
501	
502	
503	REFERENCES
504	1. Fredriksson, R., Lagerstrom, M.C., Lundin, L.G., and Schioth, H.B. (2003) The G-protein-
505	coupled receptors in the human genome form five main families. Phylogenetic analysis,
506	paralogon groups, and fingerprints. Molecular Pharmacology. 63:1256-1272.
507	
508	2. Takeda, S., Kadowaki, S., Haga, T., Takaesu, H., and Mitaku, S. (2002) Identification of G
509	protein-coupled receptor genes from the human genome sequence. FEBS Lett. 520:97-101.
510	
511	3. Flock, T., Ravarani, C.N., Sun, D., Venkatakrishnan, A.J., Kayikci, M., Tate, C.G., Veprintsev,
512	D.B., and Babu, M.M. (2015) Universal allosteric mechanism for $G\alpha$ activation by GPCRs.
513	Nature. 524:173-179.
514	

515	4.	Hofmann, K.P., Scheerer, P., Hildebrand, P.W., Choe, H.W., Park, J.H., Heck, M., and Ernst,
516		O.P. (2009) A G protein-coupled receptor at work: the rhodopsin model. Trends in Biochemical
517		Sciences. 34:540-552.
518		
519	5.	Provencio, I., Jiang, G., De Grip, W.J., Hayes, W.P., and Rollag, M.D. (1998) Melanopsin: An
520		opsin in melanophores, brain, and eye. Proceedings of the National Academy of Sciences.
521		95:340-345.
522		
523	6.	Provencio, I., Rodriguez, I.R., Jiang, G., Hayes, W.P., Moreira, E.F., and Rollag, M.D. (2000) A
524		novel human opsin in the inner retina. The Journal of Neuroscience. 20:600-605.
525		
526	7.	Schmidt, T.M., Chen, K.K., Hattar, S. (2011) Intrinsically photosensitive retinal ganglion cells:
527		many subtypes, diverse functions. Trends in Neurosciences. 34:572-580.
528		
529	8.	Quattrochi, L.E., Stabio, M.E., Kim, I., Ilardi, M.C., Fogerson, P.M., Leyrer, M.L., Berson, D.M.
530		(2019) The M6 cell: A small-field bistratified photosensitive retinal ganglion cell. The Journal of
531		Comparative Neurology. 527:297-311.
532		
533	9.	Hattar, S., Liao, H. W., Takao, M., Berson, D. M., and Yau, K. W. (2002) Melanopsin-containing
534		retinal ganglion cells: architecture, projections, and intrinsic photosensitivity. Science. 295:1065-
535		1070.
536		
537	10.	Estevez, M.E., Fogerson, P.M., Ilardi, M.C., Borghuis, B.G., Chan, E., Weng, S., Auferkorte,
538		O.N., Demb, J.B., and Berson, D.M. (2012) Form and Function of the M4 Cell, an Intrinsically
539		Photosensitive Retinal Ganglion Cell Type Contributing to Geniculocortical Vision. The Journal
540		of Neuroscience. 32:13608-13620.

541		
542	11.	Hattar, S., Lucas, R.J., Mrosovsky, N., Thompson, S., Douglas, R.H., Hankins, M.W., Lem, J.,
543		Biel, M., Hofmann, F., Foster, R.G., and Yau, K.W. (2003) Melanopsin and rod-cone
544		photoreceptive systems account for all major accessory visual functions in mice. Nature. 424:76-
545		81.
546		
547	12.	Peirson, S.N., Halford, S., and Foster, R.G. (2009) The evolution of irradiance detection:
548		melanopsin and the non-visual opsins. Philos Trans R Soc Lond B Biol Sci. 364:2849-2865.
549		
550	13.	Hattar, S., Kumar, M., Park, A., Tong, P., Tung, J., Yau, K. W., and Berson, D. M. (2006) Central
551		projections of melanopsin-expressing retinal ganglion cells in the mouse. J Comp Neurol.
552		497:326-349.
553		
554	14.	Rupp, A.C., Ren, M., Altimus, C.M., Fernandez, D.C., Richardson, M., Turek, F., Hattar, S., and
555		Schmidt, T.M. (2019) Distinct ipRGC subpopulations mediate light's acute and circadian effects
556		on body temperature and sleep. Elife. 23:e44358.
557		
558	15.	Lucas, R.J. (2013) Mammalian inner retinal photoreception. Current Biology. 23:R125-133.
559		
560	16.	Do, M.T. and Yau, K.W. (2010) Intrinsically photosensitive retinal ganglion cells. Physiol Rev
561		90:1547-1581.
562		
563	17.	Graham, D.M., Wong, K.Y., Shapiro, P., Frederick, C., Pattabiraman, K., and Berson, D.M.
564		(2008) Melanopsin Ganglion Cells Use a Membrane-Associated Rhabdomeric Phototransduction
565		Cascade. Journal of Neurophysiology. 99:2522-2532.
566		

567	18.	Panda, S., Nayak, S.K., Campo, B., Walker, J.R., Hogenesch, J.B., and Jegla, T. (2005)
568		Illumination of the Melanopsin Signaling Pathway. Science. 307:600-604.
569		
570	19.	Sonoda, T., Lee, S.K., Birnbaumer, L., and Schmidt, T.M. (2018) Melanopsin Phototransduction
571		Is Repurposed by ipRGC Subtypes to Shape the Function of Distinct Visual Circuits. Neuron.
572		99:754-767.
573		
574	20.	Jiang, Z., Yue, W.W.S., Chen, L., Sheng, Y., and Yau, K.W. (2018) Cyclic-Nucleotide- and
575		HCN-Channel Mediated Phototransduction in Intrinsically Photosensitive Retinal Ganglion Cells.
576		Cell. 175:652-664.
577		
578	21.	Zhou, X.E., Melcher, K., and Xu, H.E. (2012) Structure and activation of rhodopsin. Acta
579		Pharmacologica Sinica. 33:291-299.
580		
581	22.	Inoue, A., Raimondi, F., Kadji, F.M.N., Singh, G., Kishi, T., Uwamizu, A., Ono, Y., Shinjo, Y.,
582		Ishida, S., Arang, N., Kawakami, K., Gutkind, J.S., Aoki, J., and Russell, R.B. (2019)
583		Illuminating G-Protein-Coupling Selectivity of GPCRs. Cell. 177:1933-1947.
584		
585	23.	Masuho, I., Ostrovskaya, O., Kramer, G.M., Jones, C.D., Xie, K., and Martemyanov, K. (2015)
586		Distinct profiles of functional discrimination among G proteins determine the actions of G
587		protein-coupled receptors. Science Signaling. 8:ra123.
588		
589	24.	Okashah, N., Wan, Q., Ghosh, S., Sandhu, M., Inoue, A., Vaidehi, N., and Lambert, N.A. (2019)
590		Variable G protein determinants of GPCR coupling selectivity. PNAS. 116:12054-12059.
591		

592	25.	Newman, L.A., Walker, M.T., Brown, R.L., Cronin, T.W., and Robinson, P.R. (2003)
593		Melanopsin Forms a Functional Short-Wavelength Photopigment. Biochemistry. 42:12734-
594		12738.
595		
596	26.	Flock, T., Hauser, A.S., Lund, N., Gloriam, D.E., Balaji, S., and Babu, M.M. (2017) Selectivity
597		determinants of GPCR-G-protein binding. Nature. 545:317-322.
598		
599	27.	Blasic, Jr J.R., Brown, R.L., and Robinson, P.R. (2012) Light-dependent phosphorylation of the
600		carboxy tail of mouse melanopsin. Cellular and Molecular Life Sciences. 69:1551-1562.
601		
602	28.	Blasic, Jr, J.R., Brown, R.L., and Robinson, P.R. (2012) Phosphorylation of Mouse Melanopsin
603		by Protein Kinase A. PLOS One. 7:e45387.
604		
605	29.	Blasic, Jr J.R., Matos-Cruz, V., Ujla, D., Cameron, E.G., Hattar, S., Halpern, M.E., and Robinson,
606		P.R. (2014) Identification of critical phosphorylation sites on the Carboxy Tail of Melanopsin.
607		Biochemistry. 53:2644-2649.
608		
609	30.	Cameron, E.G. and Robinson, P.R. (2014) β -arrestin dependent deactivation of mouse
610		melanopsin. PLOS One. 9:e113138.
611		
612	31.	Somasundaram, P., Wyrick, G.R., Fernandez, D.C., Ghahari, A., Pinhal, C.M., Richardson, M.S.,
613		Rupp, A.C., Cui, L., Wu, Z., Brown, R.L., Badea, T.C., Hattar, S., and Robinson, P.R. (2017) C-
614		terminal phosphorylation regulates the kinetics of a subset of melanopsin-mediated behaviors in
615		mice. PNAS. 114:2741-2746.
616		

617	32.	Mure, L.S., Hatori, M., Zhu, Q., Demas, J., Kim, I.M., Nayak, S.K., and Panda, S. (2016)
618		Melanopsin-Encoded Response Properties of Intrinsically Photosensitive Retinal Ganglion Cells.
619		Neuron. 90:1016-1027.
620		
621	33.	Mure, L.S., Hatori, M., Ruda, K., Benegiamo, G., Demas, J., and Panda, S. (2018) Sustained
622		Melanopsin Photoresponse Is Supported by Specific Roles of β -arrestin 1 and 2 in Deactivation
623		and Regeneration of Photopigment. Cell Reports. 25:2497-2509.
624		
625	34.	Kang, Y., Kuybeda, O., de Waal P.W., Muhkerjee, S., Van Eps, N., Dutka, P., Zhou, X.E.,
626		Bartesaghi, A., Erramilli, S., Morizumi, T., Gu, X., Yin, Y., Liu, P., Jiang, Y., Meng, X., Zhao,
627		G., Melcher, K., Ernst, O.P., Kossiakoff, A.A., Subramaniam, S., and Xu, H.E. (2018) Cryo-EM
628		structure of human rhodopsin bound to an inhibitory G protein. Nature. 558:553-558.
629		
630	35.	Gao, Y., Hu, H., Ramachandran, S., Erickson, J.W., Cerione, R.A., and Skiniotis, G. (2019)
631		Structures of the Rhodopsin-Transducin Complex: Insights into G-Protein Activation. Molecular
632		Cell. 75:781-790.
633		
634	36.	Tsai, C.J., Marino, J., Adaixo, R., Pamula, F., Muehle, J., Maeda, S., Flock, T., Taylor, N.M.I.,
635		Mohammed, I., Matile, H., Dawson, R.J.P., Deupi, X., Stahlberg, H., and Schertler, G. (2019)
636		Cryo-EM structure of the rhodopsin-Gai- $\beta\gamma$ complex reveals binding of the rhodopsin C-terminal
637		tail to the Gβ subunit. Elife. 8:e46041.
638		
639	37.	Phillips, W.J. and Cerione, R.A. (1994) A C-terminal peptide of bovine rhodopsin binds to the
640		transducing alpha-subunit and facilitates its activation. Biochemical Journal. 299:351-357.
641		

642	38.	Emanuel, A.J. and Do, M.T. (2015) Melanopsin tristability for sustained and broadband
643		phototransduction. Neuron. 85:1043-1055.
644		
645	39.	Milner, E.S. and Do, M.T.H. (2017) A Population Representation of Absolute Light Intensity in
646		the Mammalian Retina. Cell. 171:865-876.
647		
648	40.	Wu, S. and Zhang, Y. (2007) LOMETS: A local meta-threading-server for protein structure
649		prediction. Nucleic Acids Research. 35:3375-3382.
650		
651	41.	Zimmermann, L., Stephens, A., Nam, S.Z., Rau, D., Kübler, J., Lozajic, M., Gabler, F., Söding,
652		J., Lupas, A.N., and Alva, V. (2018) A Completely Reimplemented MPI Bioinformatics Toolkit
653		with a New HHPred Server at its Core. Journal of Molecular Biology. 430:2237-2243.
654		
655	42.	Franke, R.R., Sakmar, T.P., Oprian, D.D., and Khorana, H.G. (1988) A Single Amino Acid
656		Subsitutition in Rhodopsin (Lysine 248Leucine) Prevents Activation of Transducin. The
657		Journal of Biological Chemistry. 263:2119-2122.
658		
659	43.	Gibson, D.G., Young, L., Chuang, R.Y., Venter, J.C., Hutchinson III, C.A., and Smith, H.O.
660		(2009) Enzymatic assembly of DNA molecules up to several hundred kilobases. Nature Methods.
661		6:343-345.
662		
663	44.	Wang, W. and Malcolm, B.A. (1999) Two-stage PCR protocol allowing introduction of multiple
664		mutations, deletions and inserstions using QuikChange Site-Directed Mutagenesis.
665		Biotechniques. 26:680-682.
666		

667	45.	Stoll, S. and Schweiger, A. (2006) EasySpin, a comprehensive software package for spectral
668		simulation and analysis in EPR. Journal of Magnetic Resonance. 178:42-55.
669		
670	46.	Rasmussen, S.G.F., DeVree, B.T., Zou, Y., Kruse, A.C., Chung, Y., Kobilka, T.S., Thian, F.S.,
671		Chae, P.S., Pardon, E., Calinski, D., Mathiesen, J.M., Shah, S.T.A., Lyons, J.A., Caffrey, M.,
672		Gellman, S.H., Steyaert, J., Skiniotis, G., Weis, W.I., Sunahara, R.K., and Kobilka, B.K. (2011)
673		Crystal structure of the β_2 adrenergic receptor-Gs protein complex. Nature. 477:549-555.
674		
675	47.	Murakami, M. and Kouyama, T. (2008) Crystal structure of squid rhodopsin. Nature. 453:363-
676		367.
677		
678	48.	Shimamura, T., Hiraki, K., Takahashi, N., Hori, T., Ago, H., Masuda, K., Takio, K., Ishiguro, M.,
679		and Miyano, M. (2008) Crystal Structure of Squid Rhodopsin with Intracellularly Extended
680		Cytoplasmic Region. The Journal of Biological Chemistry. 283:17753-17756.
681		
682	49.	Karnik, S.S. and Khorana, H.G. (1990) Assembly of Functional Rhodopsin Requires a Disulfide
683		Bond between Cysteine Residues 110 and 187. The Journal of Biological Chemistry. 265:17520-
684		17524.
685		
686	50.	Hwa, J., Reeves, P.J., Klein-Seetharaman, J., Davidson, F., and Khorana, H.G. (1999) Structure
687		and function in rhodopsin: Further elucidation of the role of the intradiscal cysteines, Cys-110, -
688		185, and -187, in rhodopsin folding and function. PNAS. 96:1932-1935.
689		
690	51.	Kim, J.M., Altenbach, C., Thurmond, R.L., Khorana, H.G., and Hubbell, W.L. (1997) Structure
691		and function in rhodopsin: Rhodopsin mutants with a neutral amino acid at E134 have a partially
692		activated conformation in the dark state. PNAS. 94:14273-14278.

693		
694	52.	Van Eps, N., Caro, L.N., Morizumi, T., Kusnetzow, A.K., Szczepek, M., Hofmann, K.P.,
695		Bayburt, T.H., Sligar, S.G., Ernst, O.P., and Hubbell, W.L. (2017) Conformational equilibria of
696		light-activated rhodopsin in nanodiscs. PNAS. 114:E3268-E3275.
697		
698	53.	Kusnetzow, A.K., Altenbach, C., and Hubbell, W.L. (2006) Conformational States and Dynamics
699		of Rhodopsin in Micelles and Bilayers. Biochemistry. 45:5538-5550.
700		
701	54.	Resek, J.F., Farahbakhsh, Z.T., Hubbell, W.L., Khorana, H.G. (1993) Formation of the Meta II
702		Photointermediate Is Accompanied by Conformational Changes in the Cytoplasmic Surface of
703		Rhodopsin. Biochemistry. 32:12025-12032.
704		
705	55.	Li, S.C., Goto, N.K., Williams, K.A., and Deber, C.M. (1996) α -Helical, but not β -sheet,
706		propensity of proline is determined by peptide environment. PNAS. 93:6676-6681.
707		
708	56.	Chini, B. and Parenti, M. (2009) G-protein-coupled receptors, cholesterol and palmitoylation:
709		facts about fats. Journal of Molecular Endocrinology. 42:371-379.
710		
711	57.	Seno, K. and Hayashi, F. (2017) Palmitoylation is a prerequisite for dimerization-dependent
712		raftophilicity of rhodopsin. The Journal of Biological Chemistry. 292:15321-15328.
713		
714	58.	Maeda, A., Okano, K., Park, P.S.H., Lem, J., Crouch, R.K., Maeda, T., and Palczewski, K. (2010)
715		Palmitoylation stabilizes unliganded rod opsin. PNAS. 107:8428-8433.
716		

717	59.	Rinaldi, S., Melaccio, F., Gozem, S., Fanelli, F., and Olivucci, M. (2014) Comparison of the
718		isomerization mechanisms of human melanopsin and invertebrate and vertebrate rhodopsins.
719		PNAS. 111:1714-1719.
720		
721	60.	Sekharan, S., Wei, J.N., and Batista, V.S. (2012) The Active Site of Melanopsin: The Biological
722		Clock Photoreceptor. Journal of the American Chemical Society. 134:19536-19539.
723		
724	61.	Hermann, R., Poppe, L., Pilbák, S., Boden, C., Maurer, J., Weber, S., and Lerchl, A. (2005)
725		Predicted 3D-structure of melanopsin, the non-rod, non-cone photopigment of the mammalian
726		circadian clock, from Djungarian hamsters (Phodopus sungorus). Neuroscience Letters. 376:76-
727		80.
728		
729	62.	Varma, N., Mutt, E., Mühle, J., Panneels, V., Terakita, A., Deupi, X., Nogly, P., Schertler,
730		G.F.X., Lesca, E. (2019) Crystal structure of jumping spider rhodopsin-1 as a light sensitive
731		GPCR. PNAS. 116:14547-14556.
732		
733		
734	63.	Kearse, M., Moir, R., Wilson, A., Stones-Havas, S., Cheung, M., Sturrock, S., Buxton, S.,
735		Cooper, A., Markowitz, S., Duran, C., Thierer, T., Ashton, B., Mentjies, P., and Drummond, A.
736		(2012) Geneious Basic: an integrated and extendable desktop software platform for the
737		organization and analysis of sequence data. Bioinformatics. 28:1647-1649.
738		
739	64.	Omasits, U., Ahrens, C.H., Muller, S., and Wollschied, B. (2014) Protter: interactive protein
740		feature visualization and integration with experimental proteomic data. Bioinformatics. 30:884-
741		886.
742		

- 743 65. Pándy-Szekeres, G., Munk, C., Tsonkov, T.M., Mordalski, S., Harpsøe, K., Hauser, A.S.,
- Bojarski, A.J., and Gloriam, D.E. (2017) GPCRdb in 2018: adding GPCR structure models and
 ligands. Nucleic Acids Research. 46:D440-D446.
- 746

747 FIGURE LEGENDS

748

749 Figure 1: Mouse melanopsin's amino acid sequence. Mouse melanopsin is 521 amino acids long, with a 750 long C-terminus domain that is 171 amino acids long. (A) Transmembrane and cytoplasmic helices are 751 annotated below the amino acid sequence. Amino acid property is denoted by the color of the residue: 752 Yellow: Nonpolar, Green: Polar, non-charged, Blue: Positively charged, Red: Negatively charged. 753 Hydrophobicity at each residue is plotted below the amino acid sequence, taller and red bars indicate a 754 high level of hydrophobicity at that position. Identities of residues possessing secondary structures are 755 based on prediction from the homology mode generated (See Figure 2). Figure generated using Geneious 756 software (63). (B) 2-dimensional schematic of mouse melanopsin C-terminus depicting functionally 757 significant residues. Figure made using Protter software (64). (C) 2D snakeplot of mouse melanopsin's 758 secondary structure. Figure from GPCRdb, using SWISSPROT for secondary structure prediction (65). 759 Note: the amino acids labeled as transmembrane helices in (A), not (C), correspond to identities of the 760 amino acids in the homology model in Figure 2.

761

Figure 2: *Homology modeling of mouse melanopsin*. (**A**) Inactive and active conformations of mouse melanopsin, generated using squid (*Todarodes pacificus*) rhodopsin as a template, and using β_2 adrenergic receptor or rhodopsin to model helical movements in the active melanopsin models. (**B**) Melanopsin in complex with Gas. Purple: melanopsin, Green: Ga, Yellow: G β , Dark Red: G γ . (**C & D**) Zoomed in views of mouse melanopsin's (purple) cytoplasmic domains when in complex with Gas C-terminal peptide (green). (**E & F**) Zoomed in views depicted in (C) and (D), depicted as a space-filling model with

768 melanopsin's electrostatic potentials plotted on the surface (Blue: positively charged, Red: negatively

charged, White: Neutral). (G) Melanopsin in complex with Gαi. Purple: melanopsin, Green: Gα, Yellow:

770 Gβ, Dark Red: Gγ. (H & I) Zoomed in views of mouse melanopsin's (purple) cytoplasmic domains when

in complex with Gαi C-terminal peptide (green). (J & K) Zoomed in views depicted in (H) and (I),

depicted as a space-filling model with melanopsin's electrostatic potentials plotted on the surface (Blue:

positively charged, Red: negatively charged, White: Neutral). (L & M) Overlay of Gas and Gai

774 complexed melanopsin structures.

775

776 Figure 3: Electron paramagnetic resonance (EPR) spectroscopy of site-directed spin-labeled mouse 777 *melanopsin in n-dodecyl-\beta-D-maltoside.* (A) Cartoon schematic of the EPR measurement of melanopsin. 778 Affinity-purified melanopsin in dodecyl maltoside solution was site-directed spin-labeled at C268, on the 3rd cytoplasmic loop. (**B**) EPR spectrum of MTSL in solution with no protein. (**C & D**) EPR spectra of 779 780 mouse melanopsin, MTSL-conjugated at C268. EPR spectra were recorded in darkness under dim red-781 light illumination at room temperature. (C) EPR spectrum of full length melanopsin C268 in darkness 782 (black trace) and after 30 sec light illumination (red trace). (D) EPR spectrum of C-terminally truncated 783 melanopsin C268 in darkness (black trace) and after 30 sec light illumination (red trace). 784 785 Figure 4: Calcium imaging of melanopsin C-terminal mutants. (A) Calcium imaging of HEK293 cells 786 expressing melanopsin $\Delta 365$, a C-terminus mutant truncated at residue L365 and (B) calcium imaging of 787 melanopsin H377P L380P Y382P, a C-terminus mutant with point mutations designed to disrupt the critical conformation at this region. (C) Calculated activation rates of melanopsin constructs tested in 788 789 these experiments. All error bars represent standard error of the mean (S.E.M.) of three independent 790 transfections. Statistical significance tested by using Students t-test, *, **, ***, **** represent P-values <0.05, 0.01, 0.001, and 0.0001, respectively. All constructs compared to wild-type melanopsin's rate, and 791 792 statistical significance is indicated over individual bars.

793

794 Figure 5: Calcium imaging of melanopsin C-terminal phosphorylation mutants. (A) Amino acid alignment of mammalian melanopsins focused on the proximal C-termini. Arrow depicts the conserved 795 796 tyrosine, Y382 in mouse melanopsin. Alignment generated using Geneious software (Kearse et al, 2012). 797 (B) Calcium imaging of HEK293 cells expressing melanopsin C-terminal phosphorylation mutants (C) 798 Calculated activation rates of melanopsin constructs depicted in (B). (D) Calcium imaging of HEK293 799 cells expressing mouse and armadillo melanopsins. (E) Calculated activation rates of melanopsin 800 constructs depicted in (D). All error bars represent standard error of the mean (S.E.M.) of three independent transfections. Statistical significance tested by using Students t-test, *, **, ***, **** 801 represent P-values <0.05, 0.01, 0.001, and 0.0001, respectively. All constructs compared to wild-type 802 melanopsin's rate, and statistical significance is indicated over individual bars. Additional statistical 803 804 analyses between constructs are indicated with the lines above the appropriate bars. 805 806 Figure 6: Calcium imaging of intracellular loop 3 mutants. (A) Homology model of melanopsin 807 depicting the region of mutations (darker purple color on the cytoplasmic extension of transmembrane helix 6) in the mutant, melanopsin 280-285 Alanine, where amino acids 280-285 are mutated to alanine 808 residues. (**B**) Homology model depicting the electrostatic charge of the 3rd cytoplasmic loop in wildtype 809 810 melanopsin and (C) the mutant melanopsin ICL3 Null, where all positively-charged residues were 811 mutated to alanines to neutralize the charge of this region (Blue: positively charged, Red: negatively 812 charged, White: Neutral). (**D**) Calcium imaging of melanopsin and the mutants depicted in (A) and (C). 813 Error bars represent S.E.M. of three independent transfections 814 Supplemental Figure 1: Template sequence coverage of mouse melanopsin model. (A & B) Mouse 815

melanopsin homology model depicting amino acids on melanopsin that were modeled using amino acids
on the *T. pacificus* template. Blue residues denote residues covered by template, pink residues are noncovered residues. (C) Template sequence coverage plotted on melanopsin's amino acid sequence.

820	Supplemental Figure 2: MATLAB/EasySpin simulation of experimental EPR data. (A) Simulation
821	spectrum fitting the full-length melanopsin C268 EPR spectrum or (B) melanopsin C268 Δ 265, C-
822	terminal truncated mutant. Rotational correlation times (τ_{corr}) calculated from the simulation, depicted
823	below the graph. Two components calculated for each construct, the 1 st and slower components represents
824	a more immobile component, derived from spin-label attached to melanopsin, while the 2 nd and faster
825	component represents a faster component, likely from excess, free spin-label in solution.
826	
827	Supplemental Figure 3: Calcium imaging of double and triple melanopsin C-terminus proline point
828	mutants. (A) Calcium imaging of HEK293 cells expressing melanopsin C-terminal mutants, synthesized
829	with combinations of point mutations at residues H377, L380, and Y382 to proline residues. (B)
830	Calculated activation rates of melanopsin constructs depicted in (A). All error bars represent standard
831	error of the mean (S.E.M.) of three independent transfections. Statistical significance tested by using
832	Students t-test, *, **, ***, **** represent P-values <0.05, 0.01, 0.001, and 0.0001, respectively. All
833	constructs compared to wild-type melanopsin's rate, and statistical significance is indicated over
834	individual bars.
835	
836	Supplemental Figure 4: Calcium imaging of melanopsin palmitoylation mutant. Representative calcium
837	imaging of melanopsin C364S, with mutation of melanopsin's predicted C-terminus palmitoylation site,
838	C364 to a serine residue. Error bars depict standard deviation (S.D.).
839	
840	Supplemental Figure 5: 2-dimensional schematics comparing mouse and armadillo melanopsin C-

841 *termini*.

and a second THE FOLD MADE TO THE CONTROL OF A CONTROL OF A CONTROL OF A CONSTRAINT OF A CONSTRAINT OF A CONTROL OF A en elemente de la complete de la complete de la construction de la construction de la complete de la complete d -----NET THE WEST CONTRACTOR OF A DESCRIPTION OF A STATE OF A DESCRIPTION OF A Received and the second s STR (0.000 (0.000) (0.







- Untransfected
 Melanopsin
- Melanopsin H377P L380P Y382P
- Melanopsin H377A L380A Y382A

A

HIKYRXA			V S G Q X X 3	IS YRIS THRIS T	S S		S	S RRRQ S
H KYR			SRRHSR	Y SYRSTERST	TS	HT S N	S	S R R R O I S
HKYR	OH		S ORSH	S SYRSTHRST	SS	0550	5	S RKROIS
HKYR				M SYRSEIL RST	SS	[0] [1]	5	S RRRO S
HKYR			R OBRI	IS ISVIRISHIERS	ISIS.	OSSD		
H KYR			S S OR NR	S SVESHERSE	55	0550		K BOLS
	N			VTČVŘČTL ŘČT				S PPPO S
	O H		SSOPTE		रॅटॅ			
	ion-1			R BYBETHBET	र्ट्ट			
			E BURD	3 8 8 8 8 8 7 8 8 7 8 7 8 7 8 7 8 7 1 8 7 1	हर			2 888372
			SPDUGU	V EVDETUDET	T R	PID S N		S D D D D T S
				VTSVDSTUDST	IN S			
H								
		1-1			T-T-T	ATSI		
		IM						
			SKUHSK	Y SYRSIAKSI			S	S KKKQES
			SROHSR	Y SYRSTHRST				SI KKKQISI
H KYR			<u>S QRSR</u>	S SYRSTHRST			2	<u>S</u> <u>H</u> <u>K</u> <u>K</u> <u>Q</u> <u>F</u> <u>S</u>
H KYK	Q		SKRHSR	M SYRSTHRST			S	
HKYR				Y I S YRS THRS T		O PD		SKKKKRQ S
			A PLOKE A	M SYRSTINGS		0 190		ISIR R R R R O IS
H KYR			A PLOKING	MISYR DHRSD	5.5		12	TERRROSS
H KYR A	QH		- SDOR			O SD	5	IS OKRO S



Phosphonull + Y082A
 Phosphonull + Y082S
 Phosphonull + Y082F





- Untransfected
 Iduse Melanopsin
 Amadillo Melanopsin
- Armadillo Malanopain + P9 (L3713, P3758, G3785, P381Y, A388T, 03918, A3938, D4128, E417T)





♦ Melanopsin 280-285 Alanine

С	A		B						
1 MDSPSGPRVLS 70	SLTQDPSFTT	30 SPALQGIWNGT 90	40 QNVSVRAQLL 100	50 S V S P T T S Á H Q 110	60 A A A W V P F P T 120				
V D V P D H A H Ý T L 130	. G T V I L L V Ġ L T 140	GMLGNLTVIYT 150	FCRNRGLRTP 160	ANMFIINLAV 170	SDFLMSVTQ 180				
APVFFASSLYK 190	KWLFGETGCE 200	FYAFCGAVFGI 210	TSMITLTAIAI 220	MDRYLVITRP 230	LATIGRGSK 240				
RRTALVLLGVW 250	260	FFGWSAYVPEG 270	L L T S C S W D Y M	TFTPQVRAYT 290	MLLFCFVFF 300				
	FIFRAIREIG 320	RACEGCGESPL 330	340	350 350	360				
WAPYSTVALVA 370 VLLGVSGQRSH 430 AAQQASGQSFC 490 RLAVCTGLSEC	380 IPSLSYRSTHR 440 SQNLEDGELK 500 IPHPHTSQFPL	390 390 S TLS SQSSDLS 450 ASSSPQVQRSK 510 AFLEDDVTLRH	400 WISGRKRQES 460 TPKVPGPSTCI 521 L	410 LGSESEVGWT 70 RPMKGQGARP	420 DTETTAAWG 80 SSLRGDQKG				



Mouse Melanopsin 38 Possible Phosphorylation Sites Armadillo Melanopsin Long Isoform 14 Possible Phosphorylation Sites