

JGR Atmospheres

RESEARCH ARTICLE

10.1029/2020JD034362

Special Section:

Carbon Weather: Toward the next generation of regional greenhouse gas inversion systems

Key Points:

- A new terrestrial ecosystem respiration parameterization is implemented in WRF-VPRM to improve the simulation of both respiration and GPP
- The daytime bands of elevated CO₂ mole fractions ahead of cold fronts form in part due to the accumulation of nighttime respiration
- Cloud shading induced perturbation of photosynthesis cannot explain the CO₂ band immediately ahead of the surface cold front on August 4, 2016

Supporting Information:

Supporting Information may be found in the online version of this article.

Correspondence to:

X.-M. Hu,
xhu@ou.edu

Citation:

Hu, X.-M., Gourdji, S. M., Davis, K. J., Wang, Q., Zhang, Y., Xue, M., et al. (2021). Implementation of improved parameterization of terrestrial flux in WRF-VPRM improves the simulation of nighttime CO₂ peaks and a daytime CO₂ band ahead of a cold front. *Journal of Geophysical Research: Atmospheres*, 126, e2020JD034362. <https://doi.org/10.1029/2020JD034362>

Received 4 DEC 2020
Accepted 21 MAR 2021

Implementation of Improved Parameterization of Terrestrial Flux in WRF-VPRM Improves the Simulation of Nighttime CO₂ Peaks and a Daytime CO₂ Band Ahead of a Cold Front

Xiao-Ming Hu¹ , Sharon M. Gourdji² , Kenneth J. Davis^{3,4} , Qingyu Wang⁵, Yao Zhang⁶ , Ming Xue¹ , Sha Feng³ , Berrien Moore⁷ , and Sean M. R. Crowell⁷ 

¹Center for Analysis and Prediction of Storms, University of Oklahoma, Norman, OK, USA, ²National Institute of Standards and Technology, Gaithersburg, MD, USA, ³Department of Meteorology and Atmospheric Science, Pennsylvania State University, University Park, PA, USA, ⁴Earth and Environmental Systems Institute, Pennsylvania State University, University Park, PA, USA, ⁵School of Meteorology, University of Oklahoma, Norman, OK, USA, ⁶Department of Environmental Engineering, Columbia University, New York, NY, USA, ⁷GeoCarb Mission, University of Oklahoma, Norman, OK, USA

Abstract Enhanced CO₂ mole fraction bands were often observed immediately ahead of cold front during the Atmospheric Carbon and Transport (ACT)-America mission and their formation mechanism is undetermined. Improved understanding and correct simulation of these CO₂ bands are needed for unbiased inverse CO₂ flux estimation. Such CO₂ bands are hypothesized to be related to nighttime CO₂ respiration and investigated in this study using WRF-VPRM, a weather-biosphere-online-coupled model, in which the biogenic fluxes are handled by the Vegetation Photosynthesis and Respiration Model (VPRM). While the default VPRM satisfactorily parameterizes gross ecosystem exchange, its treatment of terrestrial respiration as a linear function of temperature was inadequate as respiration is a nonlinear function of temperature and also depends on the amount of biomass and soil wetness. An improved ecosystem respiration parameterization including enhanced vegetation index, a water stress factor, and a quadratic temperature dependence is incorporated into WRF-VPRM and evaluated in a year-long simulation before applied to the investigation of the frontal CO₂ band on August 4, 2016. The evaluation shows that the modified WRF-VPRM increases ecosystem respiration during the growing season, and improves model skill in reproducing nighttime near-surface CO₂ peaks. A nested-domain WRF-VPRM simulation is able to capture the main characteristics of the August 4 CO₂ band and informs its formation mechanism. Nighttime terrestrial respiration leads to accumulation of near-surface CO₂ in the region. As the cold front carrying low-CO₂ air moves southeastward, and strong photosynthesis depletes CO₂ further southeast of the front, a CO₂ band develops immediately ahead of the front.

1. Introduction

An enhanced CO₂ mole fraction band was often observed immediately ahead of cold fronts during the Atmospheric Carbon and Transport (ACT)-America mission (Pal et al., 2020). Correct simulation of the gradients of CO₂ across cold fronts and understanding of the formation mechanisms are needed for accurate estimation of regional CO₂ budget (Chan et al., 2004; Hurwitz et al., 2004; Lee et al., 2012; Pal et al., 2020; Parazoo et al., 2008). Improved understanding of atmospheric constituents including CO₂ ahead of fronts may also help explain air pollution events sometimes found ahead of fronts (Hu, Xue, Kong, et al., 2019) since CO₂ and pollutants are sometimes coemitted (Brioude et al., 2012, 2013; Konovalov et al., 2014; Lindenmaier et al., 2014; Reuter et al., 2019; Turnbull et al., 2011; Wunch et al., 2009; Yang, Duan, et al., 2019). These CO₂ bands were previously briefly examined using WRF-VPRM, a weather-biosphere-online-coupled model, in which the terrestrial fluxes are handled by the Vegetation Photosynthesis and Respiration Model (VPRM) while the Weather Research and Forecast (WRF) model (Skamarock et al., 2008; Skamarock & Klemp, 2008) is used to simulate atmospheric flows and CO₂ transport and diffusion. These previous WRF-VPRM simulations were able to capture the existence of the CO₂ bands during the ACT-America mission (Hu, Crowell, Wang, Feng, et al., 2020). However substantial bias in terms of CO₂ magnitude existed, which appeared related to underestimation of diurnal variation of CO₂, particularly nighttime CO₂

peaks (Li et al., 2020; Wang et al., 2020). Other studies also implied that VPRM and WRF-VPRM tended to underestimate nighttime CO₂ fluxes (Dayalu et al., 2018; Lopez-Coto et al., 2017) and concentration (Park et al., 2020). Nighttime CO₂ biases were not extensively examined previously. Many previous regional CO₂ studies focused on daytime boundary layer CO₂, while nighttime CO₂ was less studied (Parazoo et al., 2008), likely because of two factors: 1, nighttime CO₂ simulations are less trusted because of uncertainties in the parameterization of the nighttime boundary layer and terrestrial respiration fluxes; 2, many remote-sensing and in-situ (such as aircraft) CO₂ observations are only available during daytime (Crevoisier et al., 2010; Pal et al., 2020; Sweeney et al., 2015).

Terrestrial ecosystem respiration in simple empirical modeling systems such as VPRM is often parameterized as a function of temperature, thus not accounting for factors such as soil moisture and substrate available for heterotrophic respiration (Ai et al., 2020; Buchmann, 2000; Fang & Moncrieff, 2001; Lloyd & Taylor, 1994; Raich et al., 2002; Schubert et al., 2010). VPRM simulates the Net Ecosystem Exchange (NEE) of CO₂ through separate parameterizations for ecosystem respiration (ER) and Gross Ecosystem Exchange (GEE) (Mahadevan et al., 2008):

$$NEE = ER + GEE \quad (1)$$

$$ER = \alpha \times T + \beta \quad (2)$$

$$GEE = \lambda \times T_{scale} \times W_{scale} \times P_{scale} \times FAPAR_{PAV} \times PAR \times \frac{1}{1 + \frac{PAR}{PAR_0}} \quad (3)$$

where α and β are parameters used to model ecosystem respiration as a function of temperature T , λ is the maximum light-use efficiency, T_{scale} , W_{scale} , and P_{scale} account for effects of temperature, water stress, and leaf age on photosynthesis, respectively, $FAPAR_{PAV}$ is the Fraction of Photosynthetically Active Radiation (PAR , $\mu\text{mol m}^{-2} \text{s}^{-1}$) absorbed by the photosynthetically active portion of the vegetation, which roughly equals the Enhanced Vegetation Index (EVI), and PAR_0 is the half-saturation value. EVI is derived from the MOD09A1 C6 500 m land surface reflectance data set (Zhang et al., 2016a, 2016b, 2017, 2018) and aggregated to model grids. This version of VPRM (Mahadevan et al., 2008) and the corresponding parameters previously calibrated off-line using eddy covariance tower data over North America (Hilton et al., 2016) were implemented into WRF-VPRM to examine spatiotemporal variation of CO₂ over the contiguous U.S. (Hu, Crowell, Wang, Zhang, et al., 2020) and China (Li et al., 2020; Dong et al., 2021). While Equation 3 is a light-use efficiency algorithm commonly used to simulate GEE (Dong et al., 2015; Wagle et al., 2014; Zhang et al., 2016a, 2016b, 2017, 2018), respiration (Equation 2), is considerably simpler than parameterizations of ecosystem respiration in more complex, process-based models. Ecosystem respiration depends not only on temperature but also on other factors such as soil moisture (Davidson et al., 1998; Murayama et al., 2003; Rey et al., 2002) and the mass of leaf and stem of biomes (proxy of plant productivity) that can be roughly represented by EVI (Guan et al., 2006). The leaf and stem respiration (which is the dominant contributor to autotrophic respiration) was observed to account for about 27–52% of the total ecosystem respiration on annual average, with a higher percentage during summer (Guan et al., 2006; Tang et al., 2008). Biogeochemical models like Carnegie Ames Stanford Approach (CASA) (Zhou et al., 2020) and Simple Biosphere Model (SiB4) (Haynes et al., 2019) track carbon pools, and simulate respiration as a function of the carbon pool size, along with pool-specific litter decay rates and environmental controlling factors. Other studies (Lorant et al., 2011) attempted to include plant productivity in ecosystem respiration parameterization by making respiration a function of leaf area index. A better respiration parameterization is warranted for improving WRF-VPRM.

Recently an improved terrestrial respiration parameterization in VPRM has been developed by incorporating EVI, water stress scaling factor (W_{scale}) and its interactions with temperature to capture soil moisture effects, and a quadratic dependence on surface air temperature (T in °C) (Gourdji et al., 2020; Gourdji et al., 2021):

$$ER = \beta + \alpha_1 \cdot T + \alpha_2 \cdot T^2 + \gamma \cdot EVI + k_1 \cdot W_{scale} + k_2 \cdot W_{scale} \cdot T + k_3 \cdot W_{scale} \cdot T^2 \quad (4)$$




Figure 1. (a) Enhanced vegetation index (EVI) on August 4, 2016, and (b) dominant vegetation types including water, evergreen forest, deciduous forest, mixed forest, shrubland, savanna, cropland, grassland, and urban derived from MODIS data in the simulation domain. The second domain with a 4 km grid spacing over the Northern Great Plains is marked. Six NOAA tower sites (LEF [WI], AMT [ME], WBI [IA], SCT [SC], BAO [CO], and WKT [TX]) are marked with hollow diamonds. The ACT-America B200 flight track is also marked. (c) MODIS cloud image on this day overlaid with OCO-2 XCO₂ observation. LEF, WLEF Tower; AMT, Argyle, Maine Tower; WBI, West Branch, Iowa; SCT, South Carolina Tower; BAO, Boulder Atmospheric Observatory; WKT, WKT Tower.

Linear dependence on EVI in Equation 4 represents linear dependence of terrestrial respiration on the amount of biomass. A quadratic dependence on T better captures the quick increase of respiration fluxes with increase of temperature (Fang & Moncrieff, 2001). The parameters in this improved VPRM (Equations 3 and 4) were calibrated with comprehensive flux tower data for the most abundant vegetation types in the United States, i.e., evergreen forest, deciduous forest, mixed forest, shrub, savanna, crop, and grass. Preliminary evaluation showed that this improved parameterization leads to better prediction of diurnal and monthly variation of CO₂ fluxes and concentration (Gourdji et al., 2020; Gourdji et al., 2021). This improved respiration parameterization is implemented into WRF-VPRM in this study and its performance over the contiguous United States (CONUS) is evaluated in 2016, particularly in terms of reproducing nighttime boundary layer CO₂ at six NOAA towers (see their locations in Figure 1), where hourly in situ boundary layer CO₂ data are available in both daytime and nighttime. High-resolution simulations with this improved WRF-VPRM are further used to examine the CO₂ band ahead of the cold front on 4 August 2016 observed during the ACT-America aircraft field campaign, a case well documented previously using observations (Pal et al., 2020) and simulations.

2. Modeling Approach and Evaluation Data

2.1. Brief Description of VPRM Improvement

The respiration parameterization of Equation 4 is implemented into WRF-VPRM. While in the GEE calculation (Equation 3), W_{scale} considers water stress effect using a function of Land Surface Water Index (LSWI):

$$W_{scale}(GEE) = \frac{1 + LSWI}{1 + LSWI_{max}} \quad (5)$$

following earlier VPRM development studies (Mahadevan et al., 2008; Xiao et al., 2004), W_{scale} used for ER calculation adopts

$$W_{scale}(ER) = \frac{LSWI - LSWI_{min}}{LSWI_{max} - LSWI_{min}} \quad (6)$$

proposed by Matross et al. (2006) for relatively more water-limited region, because Equation 6 leads to a larger variation of W_{scale} between different seasons with different water stresses. Equation 6 was previously used for GEE calculation in grass and shrubs, but we change it to Equation 5 to better match early growing season uptake in these ecosystems (Gourdji et al., 2020, 2021).

To maintain minimal terrestrial respiration in presence of low temperature in winter months and account for the fact that soil temperature is higher than air temperature during winter, the temperature (in °C) used in Equation 4 is adjusted to

$$T = (T_a - T_{crit}) \cdot T_{scale2} + T_{crit} \quad (7)$$

when the air temperature T_a is lower than T_{crit} . The VPRM parameters (GEE and respiration parameters in Equations 3 and 4, and T_{scale2} and T_{crit} in Equation 7) for each vegetation type are summarized in Table 1, and were calibrated by Gourdji et al., (2020) using AmeriFlux data (<https://ameriflux.lbl.gov/sites>). Respiration parameters were first calibrated using nighttime data, and then daytime respiration were computed,

Table 1
Parameter Values Used in this Study

	Evergreen forest	Deciduous forest	Mixed forest	Shrub	Savanna	Crop	Grass
T_{crit}	1	13	2	12	11	−1	3
T_{scale2}	0.05	0.1	0.05	0	0.2	0.05	0.2
T_{min}	0	0	0	0	0	2	2
T_{opt}	20	20	20	20	20	25	18
T_{max}	40	40	40	40	40	40	40
PAR_0	477.3	594	453	320	722	2,782	900
λ	−0.122	−0.096	−0.121	−0.074	−0.10	−0.078	−0.124
β	0.560	−14.048	0.077	0.922	−4.124	−0.156	−1.650
α_1	0.120	1.766	0.132	−0.114	0.68	0.072	0.180
α_2	0.003	−0.048	0.001	0.002	−0.018	−0.001	−0.006
γ	1.430	5.349	3.20	5.944	3.269	5.501	9.050
k_1	−0.531	9.008	−0.799	0.570	2.698	0.145	0.944
k_2	0.181	−1.207	0.182	−0.109	−0.382	−0.152	−0.260
k_3	−0.005	0.038	−0.002	0.004	0.016	0.016	0.013

Note. Units for parameters are as follows: λ : $\mu\text{Mol CO}_2 \text{ m}^{-2} \text{ s}^{-1} / \mu\text{mol PAR m}^{-2} \text{ s}^{-1}$; β , λ , γ , k_1 : $\mu\text{Mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$; PAR_0 : $\mu\text{Mol PAR m}^{-2} \text{ s}^{-1}$; α_1 , k_2 : $\mu\text{Mol CO}_2 \text{ m}^{-2} \text{ s}^{-1} \text{ }^\circ\text{C}^{-1}$; α_2 , k_3 : $\mu\text{Mol CO}_2 \text{ m}^{-2} \text{ s}^{-1} \text{ }^\circ\text{C}^{-2}$; temperature and the temperature parameters are in $^\circ\text{C}$. Parameters for shrub and grass are specifically optimized for this study, but other parameters are taken from Gourdji et al. (2020).

and daytime GEE were separated out, which was subsequently used to optimize GEE parameters. This is a different optimization procedure than originally formulated by Mahadevan et al. (2008), where 24 h of hourly NEE are used to simultaneously optimize GEE and respiration parameters. The new optimization procedure results in a larger magnitude of both respiration and GEE, and hence the diurnal cycle of NEE, by separately fitting respiration and GEE parameters to night and daytime observations respectively. A more detailed description of the improved VPRM and parameter calibration can be found in Gourdji et al. (2020), although parameters for western shrubs and grasslands were optimized separately for the simulations presented in this study, given that Gourdji et al. (2020) focuses on estimating fluxes in eastern North America.

2.2. Configurations of WRF-VPRM Simulations and Episode Description

Two kinds of WRF-VPRM simulations, referred to as year-long and episodic, are conducted. A single-domain, 1 year WRF-VPRM downscaling simulation is conducted over the CONUS domain at a 12-km grid spacing in 2016 to examine the updated VPRM performance over a full year. Following the year-long run, nested-domain episodic WRF-VPRM simulations are conducted for August 4, 2016 to examine the formation mechanism of the CO_2 band ahead of the cold front at a higher horizontal resolution, with an inner nested-domain covering the northern Great Plains at a 4 km grid spacing (see domain configuration in Figure 1).

The year-long WRF-VPRM simulation follows the configuration of our previous downscaling study (Hu, Crowell, Wang, Zhang, et al., 2020), with the NCEP/DOE R2 data (Kanamitsu et al., 2002) providing meteorological initial and boundary conditions and the CT2017 global simulation $3^\circ \times 2^\circ$ outputs (Peters et al., 2007) providing CO_2 initial and boundary conditions. Anthropogenic CO_2 emissions are taken from the $0.1^\circ \times 0.1^\circ$ Open-Data Inventory for Anthropogenic Carbon dioxide (ODIAC) (Oda et al., 2018) version 2018 and ocean CO_2 fluxes are taken from climatological monthly values derived by Takahashi et al. (2009).

Table 2
Configurations for Nested-Domain Sensitivity Simulations

Simulations	Configuration	Purpose
Control	Fully coupled	To diagnose the formation mechanism of the CO ₂ band ahead of the cold front
Clear-sky-photosynthesis	Use clear-sky radiation to drive VPRM	The difference between clear-sky-photosynthesis and control runs shows the impact of clouds shading on terrestrial CO ₂ fluxes, and subsequent impact on frontal CO ₂ structures
Clear-sky-surface-energy	Use clear-sky radiation to drive the Noah land surface model	The difference between clear-sky-surface-energy and control runs shows the impact of clouds on surface-energy balance, and subsequent impact on frontal CO ₂ structures

Using ODIAC led to a better CO₂ simulation than using Emission Database for Global Atmospheric Research (EDGAR) in Hu, Crowell, Wang, Zhang, et al. (2020). Selected major physics parameterization schemes within WRF include the Dudhia shortwave radiation scheme (Dudhia, 1989), the rapid radiative transfer model (RRTM) (Mlawer et al., 1997) for longwave radiation, the Noah land surface model (Chen & Dudhia, 2001), the Grell-3 cumulus scheme (Grell & Dévényi, 2002), the Morrison microphysics scheme (Morrison et al., 2009), and the Yonsei University (YSU) planetary boundary layer (PBL) scheme (Hong et al., 2006). Selection of PBL schemes is critical for accurate simulation of lower tropospheric CO₂ vertical distribution (Ballav et al., 2016; Diaz-Isaac et al., 2018) and CO₂ accumulation ahead of cold fronts. The YSU scheme has been shown to perform well for both daytime and nighttime at the 12 and 4 km grid spacings used in this study (e.g., Hu, Nielsen-Gammon, et al., 2010; Hu, Doughty, et al., 2012; Hu et al., 2013; Hu, Xue, & Li, 2019; Yang, Hu, et al., 2019). The YSU scheme is a nonlocal scheme with explicit treatment of entrainment fluxes, which was shown to be critical to reproducing convective boundary layer structures (Hu et al., 2010b, 2019b) and achieve a better performance than some local schemes such as the Mellor-Yamada-Janjic (MYJ) scheme (Wang et al., 2016). For stable boundary layer, an update in stability function in 2013 led to a better YSU performance in terms of reproducing nighttime profiles of both meteorological and chemical variables, particularly over the Great Plains (Hu et al., 2013). The model domain has 47 vertical layers extending from the surface to 10 hPa with 21 layers in the lower 2 km above the ground to resolve boundary layer processes. The WRF-VPRM downscaling simulation ran continuously from January 1 to December 31, 2016, adopting the spectral nudging technique (Gómez & Míguez-Macho, 2017; Otte et al., 2012; Vincent & Hahmann, 2015) to ensure that the large scale flows do not deviate from the driving NCEP/DOE R2 data, following our previous downscaling studies (Hu, Crowell, Wang, Zhang, et al., 2020; Hu et al., 2017, 2018). Such long-term regional simulations refine global model simulation (CT2017 in this case) and improve the general accuracy of simulation by better resolving local forcing and atmospheric processes, and are commonly referred to as dynamic downscaling.

The nested-domain simulations for August 4, 2016 examine the formation mechanism of the CO₂ band ahead of the cold front on the nested 4 km grid. During the selected episode, a surface cold front that formed on August 3, moved southeastward quickly to central Nebraska until 12 CST (18 UTC) 4 August when it slowed down (Pal et al., 2020). Along the front, scattered light clouds were present and at some isolated points/time there were large thunderstorms, e.g., at Lincoln after landing of ACT research flight. OCO-2 XCO₂ data indicate a transition of low XCO₂ to high XCO₂ across the front (Figure 1c). But due to the interference of clouds, OCO-2 could not detect the detailed gradient across the front (Bell et al., 2020; O'Dell et al., 2018). Lower troposphere measurements from the ACT-America flight campaign reveal an enhanced CO₂ mole fraction band (~50 km wide) immediately ahead of the cold front with the cross-frontal CO₂ contrast as high as 30 ppm (Pal et al., 2020). Note ppm is equivalent to μmol/mol. These CO₂ bands formed likely due to a few factors including horizontal and vertical transport, and reduced photosynthetic uptake caused by clouds (Chan et al., 2004; Hu, Crowell, Wang, Feng, et al. 2020; Lee et al., 2012). The finer horizontal and vertical resolutions used in this study can better resolve small-scale processes and shallow surface layers than earlier studies using coarse resolutions (e.g., Chan et al., 2004), which is critical for reproducing the formation of the narrow CO₂ bands. To diagnose the CO₂ band, in addition to the fully coupled control run, two sensitivity simulations (summarized in Table 2) are conducted, in which clear-sky ra-

diation is used to drive VPRM and the Noah land surface model, respectively. These sensitivity simulations are designed to isolate the impact of cloud shading on terrestrial CO₂ fluxes and surface-energy balance, respectively, and the subsequent impact on frontal CO₂ structures.

2.3. Description of Observations

2.3.1. NOAA Tall Towers

Hourly data from six NOAA tall towers over central and eastern half of US, i.e., WLEF Tower (LEF), Argyle, Maine Tower (AMT), West Branch, Iowa (WBI), South Carolina Tower (SCT), Boulder Atmospheric Observatory (BAO), and WKT Tower (WKT), are used to evaluate simulated CO₂ profiles in the continental boundary layer during both daytime and nighttime. The measurements at BAO were discontinued on July 26, 2016. More detailed description of those towers can be found at <https://www.esrl.noaa.gov/gmd/ccgg/insitu/>.

2.3.2. Atmospheric Carbon and Transport (ACT)-America Data

The ACT-America mission was conducted to improve the understanding of sources and sinks of CO₂ at regional scale ($\sim 10^6$ km²) and transport of greenhouse gases by weather systems in midlatitudes. During the mission, two aircraft (NASA Wallops C130—Hercules and NASA Langley B200—King Air) were deployed over the eastern half of the continental U.S. during 2016–2019, to measure meteorological variables and gas species including CO₂ and CH₄ across a variety of weather conditions in both the boundary layer and the free troposphere (Davis et al., 2018; Digangi et al., 2018). Airborne CO₂ were measured using a PICARRO 2401-m analyzer calibrated with gas standards traceable to the WMO scale. The ACT-America flight data on August 4, 2016 (see flight track of B200 in Figure 1) are used in this study to evaluate the CO₂ simulation with the updated WRF-VPRM and to examine the CO₂ band ahead of the cold front.

3. Results

3.1. Evaluation of Year-Long CO₂ Downscaling Simulation with Tower Data

The year-long WRF-VPRM downscaling simulation is conducted mainly to examine the performance of the updated VPRM, particularly in terms of reproducing nighttime boundary layer CO₂ peaks, because nighttime CO₂ was found to contribute substantially to the daytime CO₂ bands ahead of cold fronts (Hu, Crowell, Wang, Feng, et al., 2020). Note that the effect of simulated CO₂ variations on radiation and subsequent weather is not considered in current WRF-VPRM (such effect should be much smaller than meteorological conditions such as clouds, atmospheric temperature, and water vapor distributions, Houghton & Moir, 1977). Instead, a climatological CO₂ mole fraction (379 ppm) is used in the radiation scheme. Thus, updates in VPRM do not affect the simulated meteorological fields, which are essentially the same as those reported in Hu, Crowell, Wang, Zhang, et al. (2020). Comprehensive evaluation conducted in Hu, Crowell, Wang, Zhang, et al. (2020) concluded that the downscaled meteorological fields are sufficiently accurate to drive online-coupled VPRM and subsequent CO₂ simulation.

The updated VPRM predicts enhanced respiration during growing season, partially due to added dependence on EVI. While a more detailed evaluation of VPRM fluxes can be found in a companion paper (Gourdji et al., 2020, 2021), CO₂ fluxes at an AmeriFlux site (US-Ne3, see its location in Figure 1) near the B200 flight track are evaluated in Figure 2. During nongrowing season, both model and observations show minimal NEE except observations show occasional spikes. During growing season at this crop site (July, August, September), both negative and positive CO₂ fluxes are enhanced, indicating daytime NEE and nighttime ER, respectively. The previous version of VPRM with Equation 2 underestimates growing season NEE and ER (Figures 2a and 2c). With the modified ER parameterization (Equation 4) and newly calibrated parameters, the updated VPRM (referred to as VPRM_G in figure legends) predicts enhanced productivity with enhanced NEE during both daytime and nighttime during growing season, showing a better agreement with observations (Figures 2b and 2d). The growing season mean bias of nighttime CO₂ fluxes is reduced from -4.2 to -0.2 $\mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$ at this site, which is a critical improvement to reproducing the daytime CO₂ bands ahead of cold front as will




Figure 2. Evaluation of (a, b) hourly CO_2 flux and (c, d) its median diurnal variation during growing season simulated by (left) VPRM with respiration parametrization using Equation 2 (Mahadevan et al., 2008) and (right) VPRM_G using Equation 4 (Gourdji et al., 2020) comparing to the NEE measurements at US-Ne3, a crop site. Growing season mean bias (MB) of day and nighttime fluxes are marked in (a, b). VPRM, Vegetation Photosynthesis and Respiration Model. NEE, Net Ecosystem Exchange.

be discussed in Section 3.2. In July, EVI already increased substantially at US-Ne3 (Figure S1 in the supporting information), thus VPRM simulates enhanced NEE; while observed NEE increased much slower. The inconsistency between VPRM-simulated flux and tower flux data may indicate poor representation of model grid EVI for the US-Ne3 site in July and/or other model uncertainties in VPRM. Note that MODIS EVI not only captures the cropland, but also nearby natural vegetation in the model grid. Usually natural vegetation has an earlier growing season than crop, which might have led to earlier enhanced EVI in July than crop's actual growth at US-Ne3. Different from simulated fluxes, which represent the average flux over a $12 \times 12 \text{ km}^2$ grid cell, observed fluxes represent the flux at the site and observed nighttime fluxes often show positive spikes (Figures 2a and 2b) presumably due to instantaneous intermittent turbulence particularly during growing season (Aubinet, 2008). The median diurnal variation of CO_2 fluxes during growing season is evaluated in Figures 2c and 2d, which further confirms that the new VPRM better captures the diurnal variation of fluxes relative to the old VPRM formulation. Examination of CO_2 fluxes at a grass site in Kansas close to the 4 August B200 flight track also indicates a better performance of the new VPRM (figure not shown).

Fluxes together with transport dictate the atmospheric CO_2 spatiotemporal variation. Simulated hourly CO_2 values at six NOAA tower sites are evaluated. Among the six sites, three northern sites, WBI (crop site) and LEF and AMT (mixed forest sites), show more pronounced seasonal variation of CO_2 (Figure 3), indicating more pronounced seasonal cycle of productivity. The strong variation of hourly values in the growing season (Figure 4) is due to enhanced diurnal variation of NEE relative to nongrowing season. Detailed




Figure 3. Near-surface CO₂ in 2016, observed at six NOAA tower sites and simulated by WRF-VPRM_G. Note that CO₂ is measured at different heights AGL at different towers, simulated values are extracted at the nearest model layer. Nighttime and daytime mean bias (MB) of CO₂ at each site is marked. WRF, Weather Research and Forecast; VPRM, Vegetation Photosynthesis and Respiration Model.

hourly variation of CO₂ in first half of July is shown in Figure 5, illustrating intense diurnal variation with nighttime CO₂ peaks (gray shading) and afternoon CO₂ minima (no shading). Afternoon minimum CO₂ shows a prominent seasonal cycle with minimum in July-August at WBI crop site and in June-September at LEF, AMT mixed forest sites (Figure 3), consistent with previous studies (Bakwin et al., 1998; Parazoo et al., 2008), indicating active terrestrial uptake in growing season during daytime. Note that mixed forest typically has a longer growing season than crop (Li et al., 2020). Such seasonal variation of afternoon minimum CO₂ also shows up in the aircraft boundary layer CO₂ data from NOAA Global Greenhouse Gas




Figure 4. Median diurnal variation of near-surface CO_2 during growing season (June–September) in 2016, observed at six NOAA tower sites and simulated by WRF-VPRM_G. Note that CO_2 is measured at different heights AGL at different towers, simulated values are extracted at the corresponding model height. WRF, Weather Research and Forecast; VPRM, Vegetation Photosynthesis and Respiration Model.

Reference Network (<https://www.esrl.noaa.gov/gmd/ccgg/aircraft/>) and the XCO_2 data recorded at Total Carbon Column Observing Network [TCCON] (Hu, Crowell, Wang, Zhang, et al., 2020), both of which only have daytime observations. Relatively less-reported is the seasonal variation of nighttime CO_2 peaks/spikes (e.g., Bakwin et al., 1998; Davis et al., 2003), which increase significantly during growing seasons (Figure 3). The increasing magnitude of nighttime CO_2 peaks/spikes during growing season is even larger than the decreasing magnitude of afternoon CO_2 minimum (Figure 3). Since magnitude of nighttime CO_2 fluxes is smaller than daytime CO_2 fluxes during growing season (e.g., Figure 2), stronger nighttime increase of CO_2 concentration indicates that diurnal variation of boundary layer structure must play an important role in modulating diurnal variation of CO_2 concentration.

For SCT and two grass sites, BAO, and WKT, the contrast between the nongrowing season and growing season identified from time series of hourly CO_2 is not as clear as the crop and mixed forest sites, which is due to a smaller seasonal variation of local fluxes and location in the continent (Sweeney et al., 2015).




Figure 5. Near-surface CO₂ in first half of July 2016 (growing season for most sites), observed at six NOAA tower sites and simulated by WRF-VPRM_G in this study and WRF-VPRM in Hu, Crowell, Wang, Zhang, et al. (2020). Nighttime is shaded. WRF, Weather Research and Forecast; VPRM, Vegetation Photosynthesis and Respiration Model.

The updated VPRM showed improved performance compared to the previous version, particularly in terms of reproducing nighttime CO₂ peaks (Figure 5). Overall, the underestimation of surface CO₂ is reduced, with 24-h mean bias reducing from -5.3 to -1.0 ppm at the six NOAA sites in 2016 (Table 3). During nighttime (22–06 CST) the mean bias is reduced from -9.4 to -4.0 ppm (Table 5) while during the day (10–18 CST) the mean bias is changed from -1.1 to 1.6 ppm (Table 4). The nighttime mean bias remains most prominent at two mixed forest sites, AMT (-10.4 ppm) and LEF (-4.4 ppm) with the updated VPRM (Figure 3 and Table 5). At the WBI crop site, nighttime mean bias reduced from -11.46 to -4.2 ppm. The remaining bias at night could be due to overestimated nighttime turbulent mixing (Hu, Nielsen-Gammon, et al., 2010), as well as underestimated nighttime respiration due to drainage and loss of measured flux at towers (Aubinet, 2008). These evaluations illustrate that the updated terrestrial respiration parameterization (i.e., Equation 4) and the newly calibrated parameters (Table 1) have a much-improved skill to reproduce the nighttime CO₂ peaks, as well as diurnal variation of near-surface CO₂, particularly during growing season. Note that among the six NOAA towers, WBI is closest to the B200 flight track on August 4, 2016 and

Table 3

Overall Statistics for Simulated CO₂ Over Six NOAA Towers During Both Daytime and Nighttime by WRF-VPRM_G in this Study and WRF-VPRM in Hu, Crowell, Wang, Zhang, et al. (2020)

Sites	Day and night											
	AMT		BAO		LEF		SCT		WBI		WKT	
Simulations	VPRM_G	VPRM	VPRM_G	VPRM	VPRM_G	VPRM	VPRM_G	VPRM	VPRM_G	VPRM	VPRM_G	VPRM
Mean obs, ppm	415.12	415.12	417.12	417.12	411.33	411.33	416.47	416.47	416.19	416.19	414.32	414.32
Mean sim, ppm	410.84	406.16	414.72	412.66	410.05	405.63	417.67	414.42	415.42	409.39	415.59	410.72
Number	8153	8153	4742	4742	8124	8124	7940	7940	7754	7754	8024	8024
R	0.52	0.34	0.51	0.45	0.65	0.57	0.75	0.71	0.63	0.46	0.71	0.57
MB, ppm	−4.28	−8.96	−2.39	−4.45	−1.29	−5.70	1.20	−2.05	−0.77	−6.80	1.27	−3.61
MAGE, ppm	7.91	10.19	6.89	7.41	5.31	6.77	5.84	6.07	7.04	8.81	5.07	5.74
RMSE, ppm	15.54	19.00	10.84	11.80	9.05	11.52	8.03	8.74	11.58	15.48	7.01	8.84
NMB, %	−1.0	−2.2	−0.6	−1.1	−0.3	−1.4	0.3	−0.5	−0.2	−1.6	0.3	−0.9

Note. The metrics include correlation coefficient *R*, mean bias (MB), mean absolute gross error (MAGE), root mean-square error (RMSE), and normalized mean bias (NMB). Their formula can be found in Seigneur et al. (2000).

Table 4

Evaluation Statistics for Simulated CO₂ Over Six NOAA Towers During Daytime Between 10 and 18 CST

Sites	Late day (10–18 CST)											
	AMT		BAO		LEF		SCT		WBI		WKT	
Simulations	VPRM_G	VPRM	VPRM_G	VPRM	VPRM_G	VPRM	VPRM_G	VPRM	VPRM_G	VPRM	VPRM_G	VPRM
Mean obs, ppm	405.57	405.57	412.52	412.52	405.62	405.62	407.39	407.39	408.43	408.43	408.25	408.25
Mean sim, ppm	406.5	403.43	410.96	410.3	407.36	403.98	410.57	408.4	411.04	406.81	410.95	408.44
Number	2836	2836	1642	1642	2826	2826	2771	2771	2676	2676	2782	2782
R	0.86	0.84	0.51	0.46	0.90	0.91	0.75	0.75	0.86	0.86	0.78	0.76
MB, ppm	0.92	−2.15	−1.56	−2.23	1.74	−1.64	3.19	1.02	2.61	−1.62	2.70	0.19
MAGE, ppm	3.98	4.28	5.62	5.88	3.68	3.48	4.86	4.13	5.32	4.85	4.14	3.28
RMSE, ppm	5.75	6.4	10.41	10.84	5.22	5.11	6.52	5.70	7.38	7.19	5.44	4.56
NMB, %	0.2	−0.5	−0.4	−0.5	0.4	−0.4	0.8	0.2	0.6	−0.4	0.7	0

Table 5

Evaluation Statistics for Simulated CO₂ Over Six NOAA Towers During Nighttime Between 22–06 CST

Sites	Late night (22–06 CST)											
	AMT		BAO		LEF		SCT		WBI		WKT	
Simulations	VPRM_G	VPRM	VPRM_G	VPRM	VPRM_G	VPRM	VPRM_G	VPRM	VPRM_G	VPRM	VPRM_G	VPRM
Mean obs, ppm	424.99	424.99	421.1	421.1	416.99	416.99	424.73	424.73	423.13	423.13	419.43	419.43
Mean sim, ppm	414.62	408.58	417.85	414.65	412.58	407.26	423.64	419.52	418.95	411.67	419.22	412.53
Number	3197	3197	1865	1865	3182	3182	3100	3100	3055	3055	3152	3152
R	0.21	−0.12	0.39	0.34	0.24	0.11	0.57	0.50	0.31	−0.02	0.53	0.32
MB, ppm	−10.38	−16.42	−3.25	−6.45	−4.41	−9.72	−1.08	−5.20	−4.19	−11.46	−0.21	−6.9
MAGE, ppm	12.69	16.88	7.93	8.81	7.10	10.16	6.52	7.82	8.86	12.50	5.75	8.06
RMSE, ppm	22.79	27.79	11.29	12.79	12.02	15.91	9.10	10.89	14.67	20.66	8.11	11.54
NMB, %	−2.4	−3.9	−0.8	−1.5	−1.1	−2.3	−0.3	−1.2	−1.	−2.7	0	−1.6




Figure 6. Mean CO_2 profiles in growing season (June–September) of 2016 (first row) observed at (left) LEF and (right) AMT NOAA towers and simulated by (second row) WRF-VPRM_G in this study and (third row) WRF-VPRM in Hu, Crowell, Wang, Zhang, et al. (2020). WRF, Weather Research and Forecast; VPRM, Vegetation Photosynthesis and Respiration Model.

the flight track is mostly over the crop land category (Figures 1a and 1b). Thus, significant improvement of WRF-VPRM performance over the WBI crop site is likely to be particularly beneficial to diagnose formation mechanism of the CO_2 band observed by B200 on August 4.

In addition to terrestrial fluxes, the surface CO_2 concentration is dictated by boundary layer vertical distribution. Boundary layer CO_2 profiles are examined in Figures 6–8 to further diagnose the CO_2 biases. During daytime (10, 14, 18 CST), CO_2 in the boundary layer is nearly well mixed, except at 18 CST when CO_2 starts to accumulate near the surface at eastern sites, e.g., AMT and SCT because these sites enter evening earlier comparing to western sites. The biases of these well-mixed daytime profiles are minor, consistent with previous evaluation using daytime measurements of boundary layer CO_2 from NOAA small aircraft and XCO₂ from TCCON (Hu, Crowell, Wang, Zhang, et al., 2020).

Comparing to daytime, biases of nighttime boundary layer CO_2 are relatively less-reported. Through the night and the following early morning (22, 2, 06 CST), CO_2 from terrestrial respiration keeps accumulating




Figure 7. Same as Figure 6, but at (left) WBI and (right) SCT NOAA towers. WBI, West Branch, Iowa; SCT, South Carolina Tower.

near the surface in presence of a stable boundary layer and near-surface vertical gradients of CO_2 increase. Comparing to daytime, WRF-VPRM has more challenge to reproduce the nighttime CO_2 peaks. The bias of near-surface CO_2 becomes prominent during nighttime. With increased nighttime respiration in the updated VPRM (Equation 4), the nighttime bias is alleviated. However, the bias of early morning CO_2 at certain sites are still large. For example, at AMT, the bias reaches as large as >20 ppm at 02 CST and the bias quickly reduces to <10 ppm at 06 CST as the convective boundary layer starts to grow and photosynthesis becomes active at this eastern site, reducing surface-layer CO_2 peaks quickly (Figure 6). At 06 CST, the bias of surface CO_2 at WBI and LEF reaches ~ 16 and 10 ppm, and the biases at three other south sites, SCT, BAO, WKT are smaller. The median diurnal variations of CO_2 during growing season (Figure 4) also illustrate the largest nighttime bias remains at AMT, WBI, and LEF. Future further improvement of simulating boundary layer CO_2 diurnal variation may focus on the biologically very productive sites such as WBI crop site and AMT, LEF mixed forest sites, and may benefit from colocated measurements of fluxes, mole fraction, and boundary layer meteorology (Davis et al., 2003; Yi et al., 2001, 2004).




Figure 8. Same as Figure 6, but at (left) BAO and (right) WKT NOAA towers. BAO, Boulder Atmospheric Observatory; WKT, WKT Tower.

3.2. Examination of the CO₂ Band ahead of Cold Front on August 4, 2016

3.2.1. Analysis of ACT-America Data and Fully Coupled WRF-VPRM Control Simulation

An important purpose of this study is to examine the cause of elevated CO₂ bands ahead of cold fronts, which appeared to be related to nighttime accumulation of near-surface CO₂ (Hu, Crowell, Wang, Feng, et al., 2020). With improved capability to reproduce the nighttime CO₂ surface flux and mole fraction peaks with the updated VPRM, we seek to better diagnose the formation mechanism of the CO₂ bands. The CO₂ band ahead of the cold front on 4 August 2016 is examined with nested-domain WRF-VPRM simulations (see domain configuration in Figure 1). In early morning (06 CST), the model captures the stable boundary layer structure, as verified by routine sounding data at KLB, KOAX, and KTOP in the vicinity of the ACT-America flight track on this day (Figure 9). At 10:00 CST when C130 took off, the convective boundary layer grew up to ~700 m AGL, as indicated by the potential temperature profile measured by C130 (Figure 9b). Nighttime accumulated CO₂ in the stable boundary layer was mixed in this early shallow convective boundary layer with a vertically constant mole fraction of ~418 ppm, about 14 ppm higher than the CO₂




Figure 9. (first row) Profiles of CO₂, potential temperature (θ), and water vapor mixing ratio simulated at 10:00 CST and observed during the C130 ascending time between 10:00 and 10:15 CST on August 4 over the Lincoln airport (40.8367°N, 96.7619°W), and profiles of (left to right) wind speed, θ , and water vapor mixing ratio at 06:00 CST on August 4 simulated and observed at sounding sites: (second-fourth rows) KLB, KOAX, and KTOP, which are marked on Figure 10.

mole fraction in the free troposphere above (Figure 9a). The control WRF-VPRM simulation captures these boundary layer structures even though certain biases exist, including overestimation of boundary layer potential temperature by 2–3 K and underestimation of boundary layer CO₂ with a mean bias of ~5 ppm. During the morning, the cold front kept pushing southeastward bringing the low-CO₂ air mass (indicated by the OCO-2 XCO₂ data in Figure 1c and aircraft data in Figures 10a and 10b) into the region. Meanwhile, enhanced GEE over cropland around Lincoln and area east of Lincoln (partially due to large EVI in the region, Figure 1a) reduced the boundary layer CO₂, leading to formation of a high CO₂ mole fraction band between west of Lincoln and leading edge of the cold front, which was detected by B200 at ~12:30 CST when it flew westward and crossed the front (Figure 10). The WRF-VPRM simulation captures the CO₂ gradient across the front in both the boundary layer and free troposphere (Figure 10b) and the horizontal variation of boundary layer CO₂ mole fraction along the B200 flight track, including the CO₂ band immediately ahead of the cold front and west of Lincoln, albeit the band position is a little off (Figures 10a and 10b), which




Figure 10. (a, c) Horizontal (at ~995 m above sea level) and (b, d) vertical cross sections of (left) CO₂ and (right) equivalent potential temperature (θ_e) through the B200 flight track at 1200 CST on August 4 simulated by WRF-VPRM_G, overlaid with B200 aircraft CO₂ data. Note that the flight path and CO₂ data are marked in panel (a). The CST time is marked along the flight path. WRF, Weather Research and Forecast; VPRM, Vegetation Photosynthesis and Respiration Model.

is partially due to the time difference between observation and simulation. The CO₂ band was observed at ~12:30 CST while the simulation is taken at 12:00 CST in Figure 10 (WRF outputted hourly).

Simulated individual components of CO₂ in the higher-resolution inner domain are examined to further diagnose the CO₂ band attribution. Total CO₂ are decomposed into background, biogenic, anthropogenic, and oceanic CO₂, which are tracked by individual tracer in WRF-VPRM. While biogenic, anthropogenic, and oceanic CO₂ are CO₂ accumulated from the terrestrial biogenic fluxes, anthropogenic emissions and oceanic fluxes, background CO₂ is the CO₂ transported into the domain from the lateral boundary CO₂, which is derived from CT2017. Biogenic CO₂ can be further decomposed into respiration CO₂ and photosynthesis CO₂, which are CO₂ accumulated from transported ER and GEE, i.e., Equations 4 and 3. Among these components, oceanic CO₂ is negligible over the continent (Hu, Crowell, Wang, Zhang, et al., 2020) and therefore not shown here. The background CO₂ transported by the cold front has its origins from the north-




Figure 11. (a) Vertical cross sections of total CO₂ through the B200 flight track at 12:00 CST on August 4 simulated by WRF-VPRM_G in the nested inner domain, overlaid with B200 aircraft CO₂ data and (b)–(f) CO₂ components including background, biogenic, anthropogenic, respiration, and photosynthesis CO₂. Location of Lincoln is marked using a star on X axis. WRF, Weather Research and Forecast; VPRM, Vegetation Photosynthesis and Respiration Model.

ern boundary where CO₂ is low in summer (Hu, Crowell, Wang, Zhang, et al., 2020; Sweeney et al., 2015), thus southeastward penetration of background CO₂ carried by the cold front creates a step-function-like gradient (~6 ppm difference) across the front, with a nearly constant CO₂ (~402 ppm) ahead of the cold front (Figures 11b and 12b). The anthropogenic CO₂ includes a small contribution to CO₂ to the far east side along the B200 flight track, while it does not contribute to the CO₂ band immediately ahead of the cold front (Figures 11d and 12d). The CO₂ enhancement band is mostly comprised of biogenic CO₂ (Figure 11c), that is, accumulated from terrestrial biogenic fluxes, which contributes a ~15 ppm cross-front difference (Figure 12c). The horizontal variation of biogenic CO₂ (Figures 11c and 12c) matches the horizontal variation of total CO₂ (Figures 11a and 12a) in terms of general trend.

Figure 12 also illustrates that updated VPRM significantly enhances the model capability to capture the CO₂ band ahead of the cold front. The previous version of VPRM, with a much weaker respiration given by Equation 2, barely simulates any contribution to the cross-front CO₂ gradient from biogenic CO₂ at 12 CST,




Figure 12. (a) Total CO_2 through the B200 flight track at 12:00 CST on August 4 in the nested inner domain simulated by WRF-VPRM_G and WRF-VPRM, along with B200 aircraft CO_2 data in the boundary layer during 11:20–13:10 CST and (b–f) simulated CO_2 components including background, biogenic, anthropogenic, respiration, and photosynthesis CO_2 . Location of Lincoln is marked using a star on X axis. WRF, Weather Research and Forecast; VPRM, Vegetation Photosynthesis and Respiration Model.

with only background CO_2 contributing a ~ 6 ppm CO_2 gradient across the front. With a stronger respiration, updated WRF-VPRM simulates a dominating contribution from biogenic CO_2 (~ 15 ppm) to the cross-front CO_2 gradient, with certain biases still remaining including underestimating the peak concentration of the CO_2 band by ~ 8 ppm and overestimating the width of the CO_2 band.

To further delineate the effects of biogenic CO_2 fluxes, individual contribution history from respiration CO_2 and photosynthesis CO_2 during the morning of August 4, is examined in Figure 13. In the early morning at 08 CST, respiration CO_2 ahead of the cold front is significant (~ 40 ppm) and does not show much variation along the B200 flight track (Figure 13b). Photosynthesis CO_2 are enhanced (absolute value) over the cropland region with enhanced EVI (Figure 1a) around Lincoln and area east of Lincoln (Figure 13c). As a result of collective contribution of respiration and photosynthesis CO_2 , biogenic CO_2 shows a band of enhanced mole fraction immediately ahead of the cold front (Figure 13a). Through time, this CO_2 band is squeezed and pushed southeastward by the front, which was detected by B200 $\sim 12:30$ CST (Figure 13d). When the CO_2 band is pushed further into high GEE region east of Lincoln, stronger GEE (negative photosynthesis




Figure 13. (top to bottom) Surface biogenic CO₂, respiration, and photosynthesis CO₂ at (a, b, c) 08 and (d, e, f) 12 CST. B200 aircraft CO₂ data in the boundary layer is overlaid for location reference.

CO₂ contribution) gradually weakens the narrow CO₂ band immediately ahead of the cold front in the afternoon until it merges into the CO₂ background in the warm sector of the front (not shown).

The morning and noontime CO₂ band extends from the surface into the free troposphere (Figures 10b and 11a) ahead of the cold front, which is related to nighttime convective transport of accumulated CO₂ from the nighttime stable boundary layer into the free troposphere (Figure 14). The snapshots at 02:00 CST shows significant updrafts and downdrafts ahead of the front associated with moist convection, which ejects the near-surface CO₂-rich and water vapor-rich air mass into the free troposphere. As a result, a weaker CO₂ band extends at least 3–4 km AGL (Figure 11a). This may explain other CO₂ bands observed in the free troposphere by ACT-America flights. The moist convection in the region is more active during nighttime and weakens during the day on August 4, which might be related to the diurnal variation of low-level jet (LLJ) over the Great Plains. On the night of August 3–4, a strong LLJ is formed ahead of the cold front (not shown). The nighttime convection is presumably enhanced by convergence at the northern end of the LLJ over the Northern Great Plains. Moist convective transport of chemical species has been reported to have significant impacts on air quality (Barth et al., 2007; Hu, Fuentes, et al., 2010; Jiang et al., 2007; Lang-




Figure 14. (a) Vertical cross sections of total CO₂ through the B200 flight track at 02:00 CST on August 4 simulated by WRF-VPRM_G in the nested inner domain, overlaid with B200 aircraft CO₂ data and (b–d) CO₂ components including background, biogenic, anthropogenic CO₂, (e–f) vertical velocity, and hydrometer mixing ratio. WRF, Weather Research and Forecast; VPRM, Vegetation Photosynthesis and Respiration Model.

ford et al., 2010); it similarly impacts CO₂ transport and its budget in this case, as also shown by Penn State simulations.

3.2.2. Analysis of WRF-VPRM Sensitivity Simulations to Examine Cloud Impacts

Clouds associated with moist convection block shortwave radiation during the day, thus affecting photosynthesis, and the land surface-energy balance and eventually terrestrial CO₂ fluxes. Two sensitivity simulations (summarized in Table 2) are conducted to examine the impact of clouds on terrestrial fluxes and the frontal CO₂ band. In the clear-sky-photosynthesis simulation, clear-sky radiation is used to drive VPRM. The difference between clear-sky-photosynthesis and control runs shows the impact of clouds shading on terrestrial CO₂ fluxes (through Equation 3), and subsequent impact on frontal CO₂ structures. During the night of August 3–4, moist convection initially develops along the frontal boundary due to convergence. Through time, the convection lags behind the surface cold front convergence line along the southwest end of the convection line (Figure 15). In the control simulation, reduced shortwave radiation due to cloud shad-




Figure 15. (top to bottom) Surface shortwave radiation, GEE, and biogenic CO₂ simulated by (left) control and (middle) the clear-sky-photosynthesis sensitivity simulation, and (right) their difference at 10 CST. B200 aircraft CO₂ data in the boundary layer at ~12 CST is overlaid. GEE, Gross Ecosystem Exchange.

ing suppressed GEE, thus leading to enhanced biogenic CO₂. However, the enhanced biogenic CO₂ due to cloud shading (difference between control and clear-sky-photosynthesis) is minor (~2 ppm) comparing to the total enhancement in biogenic CO₂ ahead of the leading edge of the front (~20 ppm) and their position is not matching along the B200 flight track (Figure 15). The storm weakens/shrinks from 10 to 12 CST and does not move while the surface front keeps moving southeastward (Figure 16). So the biogenic CO₂ due to cloud shading at the time remains lagging behind the front while biogenic CO₂ during previous hours is transported southwesterly by the front, and the magnitude remains ~2 ppm (Figure 16). Thus, cloud shading barely contributes to the CO₂ band ahead of the front observed by B200 in this case (Figure 17).

At the northeast end of the convection line in the domain, the convection is at the leading edge of the front and the contribution from cloud shading to CO₂ concentration enhancement is more significant because biogenic activity is more vigorous (higher EVI), and convection is stronger in the region. Thus, in this region, cloud shading contributes more to the CO₂ band ahead of the front (Figures 15 and 16).

The impacts of clouds on CO₂ flux and mole fraction through the surface-energy balance is more complex. Since the nested-domain WRF-VPRM simulations start at 00:00 UTC August 4 (18:00 CST), there is still daylight (shortwave radiation) in the first 2 h of simulation during early evening. Using clear-sky shortwave radiation to drive surface-energy balance in the clear-sky-surface-energy simulation initially leads to slight-




Figure 16. Difference of (top to bottom) surface short wave radiation, GEE, and biogenic CO₂ simulated by the control and the clear-sky-photosynthesis sensitivity simulation, at (left) 11 CST and (right) 12 CST. B200 aircraft CO₂ data in the boundary layer at ~12 CST is overlaid. GEE, Gross Ecosystem Exchange.

ly higher temperature around clouds, which later on provides perturbation for the strong moist convection during nighttime (not shown), and the subsequent perturbation on CO₂ fields becomes quite nonlinear, which can propagate through the rest of simulation. The deep moist convection weakens during the day and cloud shading on surface temperature becomes more linear. In the control simulation, cloud shading leads to lower surface temperature compared to the clear-sky-surface-energy simulation (Figure 18g). Despite a nearly linear temperature response, its subsequent impact on CO₂ fluxes and mole fraction is complex. Compared to the clear-sky-surface-energy simulation, lower temperature in the control simulation results in lower respiration according to Equation 4 and lower boundary layer height. Lower respiration leads to lower biogenic CO₂, but lower boundary layer on the other hand leads to more accumulation of respiration CO₂. The impact of lower surface temperature on photosynthesis is also mixed depending on whether the ambient temperature is lower or higher than the optimal temperature (T_{opt}) because optimal temperature leads to the strongest GEE through the highest T_{scale} (Mahadevan et al., 2008). In the morning of August 4, temperature in the nested 4 km domain varies from 5 to 35 °C, spanning both sides of the T_{opt} (25 and 18 °C)




Figure 17. Same as Figure 12, but simulated by the Sensitivity1 (clear-sky-photosynthesis) and Sensitivity2 (clear-sky-surface-energy) simulations.

of two dominant vegetation types in the domain, i.e., crop and grass. Thus, the impact of reduced temperature due to cloud shading on NEE is mixed (Figure 18h). Another likely impact of lower temperature behind the cold front is to slightly enhance the front penetration, which might explain the negative CO_2 difference between control and clear-sky-surface-energy along some portion of the front. Thus, with all these impacts, CO_2 difference between the control and clear-sky-surface-energy simulation has mixed results with both positive and negative values (Figure 18i). Given the relatively small magnitude of these CO_2 difference (<4 ppm), the elevated CO_2 mole fraction band ahead of the front is not alleviated by the impact of clouds on surface-energy balance (Figure 17).

4. Conclusions and Discussion

An enhanced CO_2 mole fraction band was often observed immediately ahead of cold fronts during the Atmospheric Carbon and Transport (ACT)-America mission and its cause is undetermined (Pal et al., 2020). Improved understanding and correct simulation of these CO_2 bands are needed for unbiased inverse CO_2




Figure 18. (top to bottom) Temperature at 2 m AGL (T_2), NEE, and biogenic CO_2 simulated by (left) control and (middle) the clear-sky-surface-energy sensitivity simulation, and (right) their difference at 10 CST. B200 aircraft CO_2 data in the boundary layer at ~ 12 CST is overlaid. NEE, Net Ecosystem Exchange.

flux estimation. Such CO_2 bands are hypothesized to be related to nighttime CO_2 respiration and investigated in this study using WRF-VPRM, an online-coupled weather-biosphere model. While the original version of VPRM is mechanistically realistic and satisfactorily parameterizes gross ecosystem exchange with appropriate calibration, its treatment for terrestrial respiration, a linear dependence on temperature only, is inadequate to simulate ecosystem respiration that is nonlinear and also depends on biomass and soil wetness. In this study, an improved terrestrial respiration parameterization that adds enhanced vegetation index (EVI), water stress scaling factor (W_{scale}) and its interactions with temperature to capture soil moisture effects, and a quadratic function of temperature (Gourdji et al., 2020, 2021) is incorporated into the WRF-coupled VPRM and evaluated first in a year-long dynamic downscaling simulation before being applied to investigate the CO_2 band ahead of a cold front on August 4, 2016.

The year-long downscaling simulation for 2016 is conducted over the contiguous United States at a 12 km grid spacing with spectral nudging to help maintain consistency of large scale flows with the driving reanalyses. Evaluations show that the improved WRF-VPRM simulates a stronger respiration (and stronger GEE also), thus showing a better skill to reproduce nighttime near-surface CO_2 peaks measured at six NOAA towers.

Nested-domain WRF-VPRM simulations including a 4-km inner domain covering the Northern Great Plains are conducted for August 4, 2016 to examine the CO₂ band formation mechanism. The control simulation is shown to be able to capture main characteristics of the CO₂ bands, albeit with certain biases in magnitudes and/or locations. Nighttime terrestrial respiration leads to accumulation of near-surface CO₂ in the region. As the cold front with low-CO₂ air mass moves southeastward, together with strong photosynthesis depleting CO₂ further southeast of the front, an enhanced CO₂ mole fraction band develops immediately ahead of the front. Two sensitivity simulations are conducted to examine the impact of cloud shading on CO₂ attributions, using clear-sky radiation (i.e., neglecting cloud effects on radiation parameterization) to drive VPRM and land surface-energy balance, respectively. Cloud shading is not found to enhance or alleviate the CO₂ band observed by the ACT-America flight in this case because the cloud shading induced perturbation of biogenic CO₂ through radiative and surface temperature effects is not significant and its location lags behind the surface cold front.

Though not shown here, a few other CO₂ bands ahead of cold front including the ones on October 18, 2017, and May 2, 2018, were also examined using WRF-VPRM simulations. Nighttime accumulation of respiration CO₂ is found to contribute substantially to these CO₂ bands, as in the case of August 4, 2016 (Hu, Crowell, Wang, Feng, et al., 2020).

Parameterization of impact of radiation on photosynthesis in VPRM through Equation 3 may be overly simplified. Some studies reported enhanced photosynthesis in presence of diffuse radiation. Such a process is not currently considered in VPRM. Nonetheless, this study illustrates the first-order effect of cloud shading on terrestrial CO₂ fluxes and CO₂ attribution around cold fronts.

This study finds that respiration fluxes contribute significantly to the CO₂ bands ahead of cold fronts. Further improvement of terrestrial respiration parameterization, particularly over biologically very-active land covers, is warranted to improve CO₂ simulations in the vicinity of mesoscale/synoptic weather systems, which is critical to accurately estimate regional CO₂ budgets (Chan et al., 2004; Pal et al., 2020). In addition, transport/dispersion, particularly during nighttime, could also be improved to reduce CO₂ biases (Hu et al., 2013).

Data Availability Statement

The reanalysis data set was downloaded from <https://rda.ucar.edu/>, and the CT2017 data were provided by NOAA ESRL, Boulder, Colorado, USA from the website at <http://carbontracker.noaa.gov>. Tomohiro Oda provided the 0.1° × 0.1° Open-Data Inventory for Anthropogenic Carbon dioxide (ODIAC) emissions. Monthly ocean CO₂ fluxes are downloaded from https://www.ldeo.columbia.edu/res/pi/CO2/carbondioxide/pages/air_sea_flux_2000.html. ACT-America data (DOI 10.3334/ORNLDAAAC/1593) is downloaded from <https://doi.org/10.3334/ORNLDAAAC/1593>. AmeriFlux data are downloaded from <https://ameriflux.lbl.gov/sites/site-search/>. Model data produced from this study have been archived at the Luster NSF projects data server at the San Diego Super computer Center, /expanseluster/projects/uok114/xhu2.

References

- Ai, J., Xiao, S., Feng, H., Wang, H., Jia, G., & Hu, Y. (2020). A global terrestrial ecosystem respiration dataset (2001–2010) estimated with MODIS land surface temperature and vegetation indices. *Big Earth Data*, 4(2), 142–152. <https://doi.org/10.1080/20964471.2020.1768001>
- Aubinet, M. (2008). Eddy covariance CO₂ flux measurements in nocturnal conditions: An analysis of the problem. *Ecological Applications*, 18(6), 1368–1378. <https://doi.org/10.1890/06-1336.1>
- Bakwin, P. S., Tans, P. P., Hurst, D. F., & Zhao, C. (1998). Measurements of carbon dioxide on very tall towers: Results of the NOAA/CMDL program. *Tellus B: Chemical and Physical Meteorology*, 50(5), 401–415. <https://doi.org/10.3402/tellusb.v50i5.1621610.1034/j.1600-0889.1998.t01-4-00001.x>
- Ballav, S., Patra, P. K., Sawa, Y., Matsueda, H., Adachi, A., Onogi, S., et al. (2016). Simulation of CO₂ concentrations at Tsukuba tall tower using WRF-CO₂ tracer transport model. *Journal of Earth System Science*, 125(1), 47–64. <https://doi.org/10.1007/s12040-015-0653-y>
- Barth, M. C., Kim, S.-W., Wang, C., Pickering, K. E., Ott, L. E., Stenchikov, G., et al. (2007). Cloud-scale model intercomparison of chemical constituent transport in deep convection. *Atmospheric Chemistry and Physics*, 7(18), 4709–4731. <https://doi.org/10.5194/acp-7-4709-2007>
- Bell, E., O'Dell, C. W., Davis, K. J., Campbell, J., Browell, E., Scott Denning, A., et al. (2020). Evaluation of OCO-2 x variability at local and synoptic scales using lidar and in situ observations from the ACT-America Campaigns. *Journal of Geophysical Research: Atmospheres*, 125, e2019JD031400. <https://doi.org/10.1029/2019JD031400>
- Brioude, J., Angevine, W. M., Ahmadov, R., Kim, S.-W., Evan, S., McKeen, S. A., et al. (2013). Top-down estimate of surface flux in the Los Angeles Basin using a mesoscale inverse modeling technique: Assessing anthropogenic emissions of CO, NO_x and CO₂ and their impacts. *Atmospheric Chemistry and Physics*, 13(7), 3661–3677. <https://doi.org/10.5194/acp-13-3661-2013>

Acknowledgments

This study was supported by the Atmospheric Carbon and Transport (ACT)-America project through Grant NNX17AG11G at OU and Grant NNX15AG76G at Penn State. The ACT-America project is a NASA Earth Venture Suborbital 2 project funded by NASA's Earth Science Division. The last author was supported by NASA OCO-2 program under Grant 80NSSC18K0896. Computations were performed at the San Diego Supercomputer Center (SDSC) through XSEDE allocation Grant TG-ATM160014. References are made to certain commercially available products in this paper to adequately specify the experimental procedures involved. Such identification does not imply recommendation or endorsement by the National Institute of Standards and Technology, nor does it imply that these products are the best for the purpose specified.

- Brioude, J., Petron, G., Frost, G. J., Ahmadv, R., Angevine, W. M., Hsie, E.-Y., et al. (2012). A new inversion method to calculate emission inventories without a prior at mesoscale: Application to the anthropogenic CO₂ emission from Houston, Texas. *Journal of Geophysical Research*, 117, D05312. <https://doi.org/10.1029/2011JD016918>
- Buchmann, N. (2000). Biotic and abiotic factors controlling soil respiration rates in *Picea abies* stands. *Soil Biology and Biochemistry*, 32(11), 1625–1635. [https://doi.org/10.1016/S0038-0717\(00\)00077-8](https://doi.org/10.1016/S0038-0717(00)00077-8)
- Chan, D., Yuen, C. W., Higuchi, K., Shashkov, A., LIU, J., Chen, J., & Worthy, D. (2004). On the CO₂ exchange between the atmosphere and the biosphere: The role of synoptic and mesoscale processes. *Tellus B: Chemical and Physical Meteorology*, 56(3), 194–212. <https://doi.org/10.1111/j.1600-0889.2004.00104.x10.3402/tellusb.v56i3.16424>
- Chen, F., & Dudhia, J. (2001). Coupling an advanced land surface-hydrology model with the Penn State-NCAR MM5 modeling system. Part I: Model implementation and sensitivity. *Monthly Weather Review*, 129(4), 569–585. [https://doi.org/10.1175/1520-0493\(2001\)129<0569:Caalsh>2.0.Co;2](https://doi.org/10.1175/1520-0493(2001)129<0569:Caalsh>2.0.Co;2)
- Crevoisier, C., Sweeney, C., Gloor, M., Sarmiento, J. L., & Tans, P. P. (2010). Regional US carbon sinks from three-dimensional atmospheric CO₂ sampling. *Proceedings of the National Academy of Sciences of the United States of America*, 107(43), 18348–18353. <https://doi.org/10.1073/pnas.0900062107>
- Davidson, E. A., Belk, E., & Boone, R. D. (1998). Soil water content and temperature as independent or confounded factors controlling soil respiration in a temperate mixed hardwood forest. *Global Change Biology*, 4(2), 217–227. <https://doi.org/10.1046/j.1365-2486.1998.00128.x>
- Davis, K. J., Bakwin, P. S., Yi, C., Berger, B. W., Zhao, C., Teclaw, R. M., & Isebrands, J. G. (2003). The annual cycles of CO₂ and H₂O exchange over a northern mixed forest as observed from a very tall tower. *Global Change Biology*, 9(9), 1278–1293. <https://doi.org/10.1046/j.1365-2486.2003.00672.x>
- Davis, K. J., Obland, M. D., Lin, B., Lauvaux, T., O'Dell, C., Meadows, B., et al. (2018). ACT-America: L3 merged in situ atmospheric Trace gases and Flask data. ORNL Distributed Active Archive Center. <https://doi.org/10.3334/ORNLDAAAC/1593>
- Dayalu, A., Munger, J. W., Wofsy, S. C., Wang, Y., Nehrkorn, T., Zhao, Y., et al. (2018). Assessing biotic contributions to CO₂ fluxes in northern China using the Vegetation, Photosynthesis and Respiration Model (VPRM-CHINA) and observations from 2005 to 2009. *Biogeosciences*, 15(21), 6713–6729. <https://doi.org/10.5194/bg-15-6713-2018>
- Diaz-Isaac, L. I., Lauvaux, T., & Davis, K. J. (2018). Impact of physical parameterizations and initial conditions on simulated atmospheric transport and CO₂ mole fractions in the US Midwest. *Atmospheric Chemistry and Physics*, 18(20), 14813–14835. <https://doi.org/10.5194/acp-18-14813-2018>
- Digangi, J. P., Choi, Y., Nowak, J. B., Halliday, H. S., & Yang, M. M. (2018). ACT-America: L2 in situ atmospheric CO₂, CO, CH₄, and O₃ concentrations. USA. Retrieved from https://daac.ornl.gov/cgi-bin/dsviewer.pl?ds_id=1556
- Dong, J., Xiao, X., Wagle, P., Zhang, G., Zhou, Y., Jin, C., et al. (2015). Comparison of four EVI-based models for estimating gross primary production of maize and soybean croplands and tallgrass prairie under severe drought. *Remote Sensing of Environment*, 162, 154–168. <https://doi.org/10.1016/j.rse.2015.02.022>
- Dong, X., Yue, M., Jiang, Y., Hu, X.-M., Ma, Q., Pu, J., & Zhou, G. (2021). Analysis of CO₂ spatiotemporal variations in China using tower data and a weather-biosphere-online-coupled model, WRF-VPRM. *Atmospheric Chemistry and Physics*, <https://doi.org/10.5194/acp-2020-1128>
- Dudhia, J. (1989). Numerical study of convection observed during the winter monsoon experiment using a mesoscale two-dimensional model. *Journal of the Atmospheric Sciences*, 46(20), 3077–3107. [https://doi.org/10.1175/1520-0469\(1989\)046<3077:NSOCOD>2.0.CO;2](https://doi.org/10.1175/1520-0469(1989)046<3077:NSOCOD>2.0.CO;2)
- Fang, C., & Moncrieff, J. B. (2001). The dependence of soil CO₂ efflux on temperature. *Soil Biology and Biochemistry*, 33(2), 155–165. [https://doi.org/10.1016/S0038-0717\(00\)00125-5](https://doi.org/10.1016/S0038-0717(00)00125-5)
- Gómez, B., & Míguez-Macho, G. (2017). The impact of wave number selection and spin-up time in spectral nudging. *Quarterly Journal of the Royal Meteorological Society*, 143(705), 1772–1786. <https://doi.org/10.1002/qj.3032>
- Gourdji, S., Karion, A., Lopez-Coto, I., Ghosh, S., Mueller, K. L., & Zhou, Y. (2021). A modified Vegetation Photosynthesis and Respiration Model (VPRM) for the eastern USA and Canada, evaluated with comparison to atmospheric observations and other biospheric models. preprint available at ESSOAr. <https://www.essoar.org/doi/pdf/10.1002/essoar.10506768.1>
- Gourdji, S. M., Lopez-Coto, I., Karion, A., Mueller, K., Ghosh, S., Baker, I., et al. (2020). Modeling (and evaluating) the upwind biosphere for urban CO₂ inversions in the Northeast Corridor. Paper presented at the ACT-America Community Workshop. Retrieved from <https://drive.google.com/drive/folders/1hm0e0QuIp3g4TAQ8KWejSfc7VW9ZXoUA>
- Grell, G. A., & Dévényi, D. (2002). A generalized approach to parameterizing convection combining ensemble and data assimilation techniques. *Geophysical Research Letters*, 29(14), 1693. <https://doi.org/10.1029/2002GL015311>
- Guan, D.-X., Wu, J.-B., Zhao, X.-S., Han, S.-J., Yu, G.-R., Sun, X.-M., & Jin, C.-J. (2006). CO₂ fluxes over an old, temperate mixed forest in northeastern China. *Agricultural and Forest Meteorology*, 137(3), 138–149. <https://doi.org/10.1016/j.agrformet.2006.02.003>
- Haynes, K. D., Baker, I. T., Denning, A. S., Stöckli, R., Schaefer, K., Lokupitiya, E. Y., & Haynes, J. M. (2019). Representing grasslands using dynamic prognostic phenology based on biological growth stages: 1. Implementation in the simple biosphere model (SiB4). *Journal of Advances in Modeling Earth Systems*, 11, 4423–4439. <https://doi.org/10.1029/2018MS001540>
- Hilton, T. W., Davis, K. J., Keller, K., & Urban, N. M. (2016). NACP VPRM NEE parameters optimized to North American flux tower sites, 2000–2006. Retrieved from http://daac.ornl.gov/cgi-bin/dsviewer.pl?ds_id=1349
- Hong, S.-Y., Noh, Y., & Dudhia, J. (2006). A new vertical diffusion package with an explicit treatment of entrainment processes. *Monthly Weather Review*, 134(9), 2318–2341. <https://doi.org/10.1175/MWR3199.1>
- Houghton, J. T., & Moir, D. W. (1977). *The physics of atmospheres*. Cambridge, UK: Cambridge University Press.
- Hu, X.-M., Crowell, S., Wang, Q., Feng, S., & Davis, K. (2020). Examine CO₂ bands along cold fronts observed during ACT-America using WRF-VPRM. Paper presented at the ACT-America Community Workshop. Retrieved from <https://drive.google.com/drive/folders/1hm0e0QuIp3g4TAQ8KWejSfc7VW9ZXoUA>
- Hu, X.-M., Crowell, S., Wang, Q., Zhang, Y., Davis, K. J., Xue, M., et al. (2020). Dynamical downscaling of CO₂ in 2016 over the contiguous United States using WRF-VPRM, a weather-biosphere-online-coupled model. *Journal of Advances in Modeling Earth Systems*, 12, e2019MS001875. <https://doi.org/10.1029/2019MS001875>
- Hu, X.-M., Doughty, D. C., Sanchez, K. J., Joseph, E., & Fuentes, J. D. (2012). Ozone variability in the atmospheric boundary layer in Maryland and its implications for vertical transport model. *Atmospheric Environment*, 46, 354–364. <https://doi.org/10.1016/j.atmosenv.2011.09.054>
- Hu, X.-M., Fuentes, J. D., & Zhang, F. (2010). Downward transport and modification of tropospheric ozone through moist convection. *Journal of Atmospheric Chemistry*, 65(1), 13–35. <https://doi.org/10.1007/s10874-010-9179-5>
- Hu, X.-M., Klein, P. M., & Xue, M. (2013). Evaluation of the updated YSU planetary boundary layer scheme within WRF for wind resource and air quality assessments. *Journal of Geophysical Research: Atmospheres*, 118, 490–510. <https://doi.org/10.1002/jgrd.50823>

- Hu, X.-M., Nielsen-Gammon, J. W., & Zhang, F. (2010). Evaluation of three planetary boundary layer schemes in the WRF model. *Journal of Applied Meteorology and Climatology*, 49(9), 1831–1844. <https://doi.org/10.1175/2010JAMC2432.1>
- Hu, X.-M., Xue, M., Kong, F., & Zhang, H. (2019). Meteorological conditions during an ozone episode in Dallas-Fort Worth, Texas, and impact of their modeling uncertainties on air quality prediction. *Journal of Geophysical Research: Atmospheres*, 124, 1941–1961. <https://doi.org/10.1029/2018JD029791>
- Hu, X.-M., Xue, M., & Li, X. (2019). The use of high-resolution sounding data to evaluate and optimize nonlocal PBL schemes for simulating the slightly stable upper convective boundary layer. *Monthly Weather Review*, 147(10), 3825–3841. <https://doi.org/10.1175/mwr-d-19-0085.1>
- Hu, X.-M., Xue, M., & McPherson, R. A. (2017). The importance of soil-type contrast in modulating august precipitation distribution near the edwards plateau and Balcones escarpment in Texas. *Journal of Geophysical Research: Atmospheres*, 122, 711–810. <https://doi.org/10.1002/2017JD027035>
- Hu, X.-M., Xue, M., McPherson, R. A., Martin, E., Rosendahl, D. H., & Qiao, L. (2018). Precipitation dynamical downscaling over the great plains. *Journal of Advances in Modeling Earth Systems*, 10(2), 421–447. <https://doi.org/10.1002/2017ms001154>
- Hurwitz, M. D., Ricciuto, D. M., Bakwin, P. S., Davis, K. J., Wang, W., Yi, C., & Butler, M. P. (2004). Transport of carbon dioxide in the presence of storm systems over a northern Wisconsin forest. *Journal of the Atmospheric Sciences*, 61(5), 607–618. [https://doi.org/10.1175/1520-0469\(2004\)061<0607:TOCDIT>2.0.CO;2](https://doi.org/10.1175/1520-0469(2004)061<0607:TOCDIT>2.0.CO;2)
- Jiang, J. H., Livesey, N. J., Su, H., Neary, L., McConnell, J. C., & Richards, N. A. D. (2007). Connecting surface emissions, convective uplifting, and long-range transport of carbon monoxide in the upper troposphere: New observations from the Aura Microwave Limb Sounder. *Geophysical Research Letters*, 34, L18812. <https://doi.org/10.1029/2007GL030638>
- Kanamitsu, M., Ebisuzaki, W., Woollen, J., Yang, S.-K., Hnilo, J. J., Fiorino, M., & Potter, G. L. (2002). NCEP-DOE AMIP-II reanalysis (R-2). *Bulletin of the American Meteorological Society*, 83(11), 1631–1643. <https://doi.org/10.1175/Bams-83-11-1631>
- Konovalov, I. B., Berezin, E. V., Ciaia, P., Broquet, G., Beekmann, M., Hadji-Lazarou, J., et al. (2014). Constraining CO₂ emissions from open biomass burning by satellite observations of co-emitted species: A method and its application to wildfires in Siberia. *Atmospheric Chemistry and Physics*, 14(19), 10383–10410. <https://doi.org/10.5194/acp-14-10383-2014>
- Langford, A. O., Tucker, S. C., Senff, C. J., Banta, R. M., Brewer, W. A., Alvarez, R. J., II, et al. (2010). Convective venting and surface ozone in Houston during TexAQS 2006. *Journal of Geophysical Research*, 115, D16306. <https://doi.org/10.1029/2009JD013301>
- Lee, T. R., De Wekker, S. F. J., Andrews, A. E., Kofler, J., & Williams, J. (2012). Carbon dioxide variability during cold front passages and fair weather days at a forested mountaintop site. *Atmospheric Environment*, 46, 405–416. <https://doi.org/10.1016/j.atmosenv.2011.09.068>
- Li, X., Hu, X.-M., Cai, C., Jia, Q., Zhang, Y., Liu, J., et al. (2020). Terrestrial CO₂ fluxes, concentrations, sources and budget in North-east China: Observational and modeling studies. *Journal of Geophysical Research: Atmospheres*, 125, e2019JD031686. <https://doi.org/10.1029/2019JD031686>
- Lindenmaier, R., Dubey, M. K., Henderson, B. G., Butterfield, Z. T., Herman, J. R., Rahn, T., & Lee, S.-H. (2014). Multiscale observations of CO₂, ¹³CO₂, and pollutants at Four Corners for emission verification and attribution. *Proceedings of the National Academy of Sciences of the United States of America*, 111(23), 8386–8391. <https://doi.org/10.1073/pnas.1321883111>
- Lloyd, J., & Taylor, J. A. (1994). On the temperature dependence of soil respiration. *Functional Ecology*, 8(3), 315–323. <https://doi.org/10.2307/2389824>
- Lopez-Coto, I., Prasad, K., & Whetstone, J. R. (2017). *Carbon dioxide biogenic vs anthropogenic sectoral contribution for the Indianapolis flux experiment (INFLUX)*. Gaithersburg, MD: National Institute of Standards and Technology. <https://doi.org/10.6028/nist.sp.1237>
- Lorant, M. M., Goetz, S. J., Rastetter, E. B., Rocha, A. V., Shaver, G. R., Humphreys, E. R., & Laflaur, P. M. (2011). Scaling an instantaneous model of Tundra NEE to the Arctic Landscape. *Ecosystems*, 14(1), 76–93. <https://doi.org/10.1007/s10021-010-9396-4>
- Mahadevan, P., Wofsy, S. C., Matross, D. M., Xiao, X., Dunn, A. L., Lin, J. C., et al. (2008). A satellite-based biosphere parameterization for net ecosystem CO₂ exchange: Vegetation Photosynthesis and Respiration Model (VPRM). *Global Biogeochemical Cycles*, 22, GB2005. <https://doi.org/10.1029/2006GB002735>
- Matross, D. M., Andrews, A., Pathmathevan, M., Gerbig, C., Lin, J. C., Wofsy, S. C., et al. (2006). Estimating regional carbon exchange in New England and Quebec by combining atmospheric, ground-based and satellite data. *Tellus B: Chemical and Physical Meteorology*, 58(5), 344–358. <https://doi.org/10.1111/j.1600-0889.2006.00206.x>
- Mlawer, E. J., Taubman, S. J., Brown, P. D., Iacono, M. J., & Clough, S. A. (1997). Radiative transfer for inhomogeneous atmospheres: RRTM, a validated correlated-k model for the longwave. *Journal of Geophysical Research*, 102(D14), 16663–16682. <https://doi.org/10.1029/97JD00237>
- Morrison, H., Thompson, G., & Tatarskii, V. (2009). Impact of cloud microphysics on the development of trailing stratiform precipitation in a simulated Squall Line: Comparison of one- and two-moment schemes. *Monthly Weather Review*, 137(3), 991–1007. <https://doi.org/10.1175/2008mwr2556.1>
- Murayama, S., Saigusa, N., Chan, D., Yamamoto, S., Kondo, H., & Eguchi, Y. (2003). Temporal variations of atmospheric CO₂ concentration in a temperate deciduous forest in central Japan. *Tellus B: Chemical and Physical Meteorology*, 55(2), 232–243. <https://doi.org/10.3402/tellusb.v55i2.16751>
- Oda, T., Maksyutov, S., & Andres, R. J. (2018). The Open-source Data Inventory for Anthropogenic CO₂, version 2016 (ODIAC2016): A global monthly fossil fuel CO₂ gridded emissions data product for tracer transport simulations and surface flux inversions. *Earth System Science Data*, 10(1), 87–107. <https://doi.org/10.5194/essd-10-87-2018>
- O'Dell, C. W., Eldering, A., Wennberg, P. O., Crisp, D., Gunson, M. R., Fisher, B., et al. (2018). Improved retrievals of carbon dioxide from Orbiting Carbon Observatory-2 with the version 8 ACOS algorithm. *Atmospheric Measurement Techniques*, 11(12), 6539–6576. <https://doi.org/10.5194/amt-11-6539-2018>
- Otte, T. L., Nolte, C. G., Otte, M. J., & Bowden, J. H. (2012). Does Nudging Quell the extremes in regional climate modeling? *Journal of Climate*, 25(20), 7046–7066. <https://doi.org/10.1175/JCLI-D-12-00048.1>
- Pal, S., Davis, K. J., Lauvaux, T., Browell, E. V., Gaudet, B. J., Stauffer, D. R., et al. (2020). Observations of greenhouse gas changes across summer frontal boundaries in the Eastern United States. *Journal of Geophysical Research: Atmospheres*, 125, e2019JD030526. <https://doi.org/10.1029/2019JD030526>
- Parazoo, N. C., Denning, A. S., Kawa, S. R., Corbin, K. D., Lokupitiya, R. S., & Baker, I. T. (2008). Mechanisms for synoptic variations of atmospheric CO₂ in North America, South America and Europe. *Atmospheric Chemistry and Physics*, 8(23), 7239–7254. <https://doi.org/10.5194/acp-8-7239-2008>

- Park, C., Park, S.-Y., Gurney, K. R., Gerbig, C., DiGangi, J. P., Choi, Y., & Lee, H. W. (2020). Numerical simulation of atmospheric CO₂ concentration and flux over the Korean Peninsula using WRF-VPRM model during Korus-AQ 2016 campaign. *PloS One*, 15(1), e0228106. <https://doi.org/10.1371/journal.pone.0228106>
- Peters, W., Jacobson, A. R., Sweeney, C., Andrews, A. E., Conway, T. J., Masarie, K., et al. (2007). An atmospheric perspective on North American carbon dioxide exchange: CarbonTracker. *Proceedings of the National Academy of Sciences of the United States of America*, 104(48), 18925–18930. <https://doi.org/10.1073/pnas.0708986104>
- Raich, J. W., Potter, C. S., & Bhagawati, D. (2002). Interannual variability in global soil respiration, 1980–94. *Global Change Biology*, 8(8), 800–812. <https://doi.org/10.1046/j.1365-2486.2002.00511.x>
- Reuter, M., Buchwitz, M., Schneising, O., Krautwurst, S., O'Dell, C. W., Richter, A., et al. (2019). Toward monitoring localized CO₂ emissions from space: Co-located regional CO₂ and NO₂ enhancements observed by the OCO-2 and S5P satellites. *Atmospheric Chemistry and Physics*, 19(14), 9371–9383. <https://doi.org/10.5194/acp-19-9371-2019>
- Rey, A., Pegoraro, E., Tedeschi, V., De Parri, I., Jarvis, P. G., & Valentini, R. (2002). Annual variation in soil respiration and its components in a coppice oak forest in Central Italy. *Global Change Biology*, 8(9), 851–866. <https://doi.org/10.1046/j.1365-2486.2002.00521.x>
- Schubert, P., Eklundh, L., Lund, M., & Nilsson, M. (2010). Estimating northern peatland CO₂ exchange from MODIS time series data. *Remote Sensing of Environment*, 114(6), 1178–1189.
- Seigneur, C., Pun, B., Pai, P., Louis, J.-F., Solomon, P., Emery, C., et al. (2000). Guidance for the performance evaluation of three-dimensional air quality modeling systems for particulate matter and visibility. *Journal of the Air & Waste Management Association*, 50(4), 588–599. <https://doi.org/10.1080/10473289.2000.10464036>
- Skamarock, W. C., & Klemp, J. B. (2008). A time-split nonhydrostatic atmospheric model for weather research and forecasting applications. *Journal of Computational Physics*, 227(7), 3465–3485. <https://doi.org/10.1016/j.jcp.2007.01.037>
- Skamarock, W. C., Klemp, J., Dudhia, J., Gill, D., Barker, M., Duda, K., et al. (2008). *A description of the advanced research WRF version 3*. (NCAR Tech. Note TN-475_STR). Boulder, CO: University Corporation for Atmospheric Research.
- Sweeney, C., Karion, A., Wolter, S., Newberger, T., Guenther, D., Higgs, J. A., et al. (2015). Seasonal climatology of CO₂ across North America from aircraft measurements in the NOAA/ESRL Global Greenhouse Gas Reference Network. *Journal of Geophysical Research: Atmospheres*, 120, 5155–5190. <https://doi.org/10.1002/2014JD022591>
- Takahashi, T., Sutherland, S. C., Wanninkhof, R., Sweeney, C., Feely, R. A., Chipman, D. W., et al. (2009). Climatological mean and decadal change in surface ocean pCO₂, and net sea-air CO₂ flux over the global oceans. *Deep-Sea Research Part II-Topical Studies in Oceanography*, 56(8–10), 554–577. <https://doi.org/10.1016/j.dsr2.2008.12.009>
- Tang, J., Bolstad, P. V., Desai, A. R., Martin, J. G., Cook, B. D., Davis, K. J., & Carey, E. V. (2008). Ecosystem respiration and its components in an old-growth forest in the Great Lakes region of the United States. *Agricultural and Forest Meteorology*, 148(2), 171–185. <https://doi.org/10.1016/j.agrformet.2007.08.008>
- Turnbull, J. C., Karion, A., Fischer, M. L., Faloona, I., Guilderson, T., Lehman, S. J., et al. (2011). Assessment of fossil fuel carbon dioxide and other anthropogenic trace gas emissions from airborne measurements over Sacramento, California in spring 2009. *Atmospheric Chemistry and Physics*, 11(2), 705–721. <https://doi.org/10.5194/acp-11-705-2011>
- Vincent, C. L., & Hahmann, A. N. (2015). The impact of grid and spectral nudging on the variance of the near-surface wind speed. *Journal of Applied Meteorology and Climatology*, 54(5), 1021–1038. <https://doi.org/10.1175/Jamc-D-14-0047.1>
- Wagle, P., Xiao, X., Torn, M. S., Cook, D. R., Matamala, R., Fischer, M. L., et al. (2014). Sensitivity of vegetation indices and gross primary production of tallgrass prairie to severe drought. *Remote Sensing of Environment*, 152, 1–14. <https://doi.org/10.1016/j.rse.2014.05.010>
- Wang, Q., Crowell, S., Hu, X.-M., & Davis, K. (2020). *Evaluation of online and Offline regional modeled CO₂ transport with INFLUX observations*. Boston. Paper presented at the 100th AMS Annual Meeting. Retrieved from <https://ams.confex.com/ams/2020Annual/meetingapp.cgi/Paper/364572>
- Wang, W., Shen, X., & Huang, W. (2016). A comparison of boundary-layer characteristics simulated using different parametrization schemes. *Boundary-Layer Meteorology*, 161(2), 375–403. <https://doi.org/10.1007/s10546-016-0175-4>
- Wunch, D., Wennberg, P. O., Toon, G. C., Keppel-Aleks, G., & Yavin, Y. G. (2009). Emissions of greenhouse gases from a North American megacity. *Geophysical Research Letters*, 36, L15810. <https://doi.org/10.1029/2009GL039825>
- Xiao, X., Hollinger, D., Aber, J., Goltz, M., Davidson, E. A., Zhang, Q., & Moore, B. (2004). Satellite-based modeling of gross primary production in an evergreen needleleaf forest. *Remote Sensing of Environment*, 89(4), 519–534. <https://doi.org/10.1016/j.rse.2003.11.008>
- Yang, S., Duan, F., Ma, Y., He, K., Zhu, L., Ma, T., et al. (2019). Haze formation indicator based on observation of critical carbonaceous species in the atmosphere. *Environmental Pollution*, 244, 84–92. <https://doi.org/10.1016/j.envpol.2018.10.006>
- Yang, Y., Hu, X.-M., Gao, S., & Wang, Y. (2019). Sensitivity of WRF simulations with the YSU PBL scheme to the lowest model level height for a sea fog event over the Yellow Sea. *Atmospheric Research*, 215, 253–267. <https://doi.org/10.1016/j.atmosres.2018.09.004>
- Yi, C., Davis, K. J., Bakwin, P. S., Denning, A. S., Zhang, N., Desai, A., et al. (2004). Observed covariance between ecosystem carbon exchange and atmospheric boundary layer dynamics at a site in northern Wisconsin. *Journal of Geophysical Research*, 109, D08302. <https://doi.org/10.1029/2003JD004164>
- Yi, C., Davis, K. J., Berger, B. W., & Bakwin, P. S. (2001). Long-term observations of the dynamics of the continental planetary boundary layer. *Journal of the Atmospheric Sciences*, 58(10), 1288–1299. [https://doi.org/10.1175/1520-0469\(2001\)058<1288:LTOOTD>2.0.CO;2](https://doi.org/10.1175/1520-0469(2001)058<1288:LTOOTD>2.0.CO;2)
- Zhang, Y., Xiao, X., Guanter, L., Zhou, S., Ciais, P., Joiner, J., et al. (2016). Precipitation and carbon-water coupling jointly control the inter-annual variability of global land gross primary production. *Scientific Reports*, 6, 39748. <https://doi.org/10.1038/srep39748>
- Zhang, Y., Xiao, X., Jin, C., Dong, J., Zhou, S., Wagle, P., et al. (2016). Consistency between sun-induced chlorophyll fluorescence and gross primary production of vegetation in North America. *Remote Sensing of Environment*, 183, 154–169. <https://doi.org/10.1016/j.rse.2016.05.015>
- Zhang, Y., Xiao, X., Wolf, S., Wu, J., Wu, X., Gioli, B., et al. (2018). Spatio-temporal convergence of maximum daily light-use efficiency based on radiation absorption by canopy chlorophyll. *Geophysical Research Letters*, 45, 3508–3519. <https://doi.org/10.1029/2017GL076354>
- Zhang, Y., Xiao, X., Wu, X., Zhou, S., Zhang, G., Qin, Y., & Dong, J. (2017). A global moderate resolution dataset of gross primary production of vegetation for 2000–2016. *Scientific Data*, 4, 170165. <https://doi.org/10.1038/sdata.2017.165>
- Zhou, Y., Williams, C. A., Lauvaux, T., Davis, K. J., Feng, S., Baker, I., et al. (2020). A multiyear gridded data ensemble of surface biogenic carbon fluxes for north America: Evaluation and analysis of results. *Journal of Geophysical Research: Biogeosciences*, 125, e2019JG005314. <https://doi.org/10.1029/2019JG005314>