

Simulating Decay Energy Spectra Using Geant4

Timothy R. Ockrin¹, Katrina E. Koehler¹, Ryan P. Fitzgerald²

¹ Houghton University, 1 Willard Ave, Houghton, NY 14774


² National Institute of Standards and Technology, 100 Bureau Dr, Gaithersburg, MD 20899

Nuclear Safeguards and Metrology




Trustworthy measurements are essential to managing risk and informing decisions

Alpha Decay




Beta Decay




Examples of decay modes measurable with Decay Energy Spectroscopy

Simulating Energy Transport



Compton Scattering



Photoelectric Effect



Pair Production



Simulation Assumptions


All decays measured by detector (e.g. no deadtime)



All energies measured perfectly (e.g. linear response)



Simulation Parameters




Convolving with Detector Response




Gaussian detector response convolved with Geant4 energy histogram output


Decay Energy Spectroscopy SIMulation for Absolute Total Efficiency



Total spectrum with all simulated events




Measurement Time: determined by time it takes the highest activity isotope to reach total simulated events



Total spectrum with scaled number of simulated events

Next Steps

Linearly fit experimental data with simulated spectra



Run Geant4 directly from DESSIMATE instead of command prompt

```
C:\windows\system32\cmd.exe
C:\g4work\Geant4.10.7.2\TrueBq01-build>
```